


# New Zealand Threat Classification System lists

2005

Rod Hitchmough, Leigh Bull and Pam Cromarty (compilers)

Published by  
Science & Technical Publishing  
Department of Conservation  
PO Box 10420  
The Terrace  
Wellington 6143, New Zealand



Cover: Archey's frog on fern, Whareorino, 2003. *Photo: Paul Schilov.*

© Copyright January 2007, New Zealand Department of Conservation

ISBN 0-478-14128-9

This report was prepared for publication by Science & Technical Publishing; editing and layout by Amanda Todd. Publication was approved by the Chief Scientist (Research, Development & Improvement Division), Department of Conservation, Wellington, New Zealand. Individual copies are printed, and are also available from the departmental website in pdf form. Titles are listed in our catalogue on the website, refer [www.doc.govt.nz](http://www.doc.govt.nz) under *Publications*, then *Science & technical*.

In the interest of forest conservation, we support paperless electronic publishing. When printing, recycled paper is used wherever possible.

# Foreword

New Zealand has some of the most ancient and fascinating species in the world (e.g. the tuatara). Many of our plants and animals are found nowhere else. However, since the arrival of humans about 1000 years ago, we have also been world leaders in rates of extinctions (particularly of land and freshwater birds, where nearly one third have been lost), and in our levels of threatened species—a legacy of a history of unsustainable harvest, habitat destruction and alien species introductions.

Preventing the extinction of New Zealand's unique plant and animal species is a critical element in the Government's New Zealand Biodiversity Strategy; it is a responsibility we owe to the rest of the world, but this is not a small task.

A vital step in doing this is to identify those species that are at risk of extinction, and to measure the level and nature of that risk. This information will allow us to focus our resources on the highest priority actions necessary to prevent extinction.

In 2002, the Department of Conservation (DOC) published a document written by a group that was led by Janice Molloy, which set out a system for classifying species according to threat of extinction using criteria that had specifically been developed for New Zealand conditions. At the same time, a companion volume compiled by Rod Hitchmough presented the results of using that system to classify all those species for which sufficient information was available. For the first time, it was possible to provide an accurate threat classification for all those taxa for which information was available, ensuring that all those that were threatened received the necessary attention to secure their future.

This document, which has been compiled by Rod Hitchmough, Leigh Bull and Pam Cromarty, updates the earlier lists, recording both actual changes in status and changes in our understanding as more knowledge is gathered, with the inclusion of additional species in many groups. I greatly appreciate the willingness of experts from outside DOC to share their knowledge and devote often substantial amounts of their valuable time to compiling this information, and for their positive attitude to the threat classification system.

John Omblér  
General Manager Research, Development & Improvement



## CONTENTS

Foreword	3
Abstract	7
Introduction	8
Interpreting these lists—the importance of qualifiers	8
The process	9
Additional information available in electronic but not published form	10
Coverage	10
Listing at one taxonomic level only	11
Distribution of listings	12
Conservancy distributions	12
Changes in the threatened category status of indigenous species since the previous listing exercise	14
Extinction	23
Updating listings	24
Acknowledgements	25
References	25
Appendix 1	
Taxonomic lists by threat category	26
Extinct	26
Nationally Critical	28
Nationally Endangered	38
Nationally Vulnerable	45
Serious Decline	47
Gradual Decline	49
Sparse	55
Range Restricted	64
Data Deficient	99

---

**Appendix 2**

---

**Threat-listing information and questionnaire****135**

---

**Appendix 3**

---

**New Zealand Threat Classification System expert panel membership****145**

---

**Index**

---

**146**

# New Zealand Threat Classification System lists

2005

Rod Hitchmough, Leigh Bull and Pam Cromarty (compilers)

Research, Development & Improvement Division, Department of Conservation,  
PO Box 10420, The Terrace, Wellington 6143, New Zealand

Email: [rhitchmough@doc.govt.nz](mailto:rhitchmough@doc.govt.nz)

## A B S T R A C T

New Zealand's threatened and potentially threatened species of animals (marine mammals, bats, birds, reptiles, amphibians, freshwater and marine fish, and terrestrial, freshwater and marine invertebrates) and non-vascular plants (bryophytes and macroalgae) were assessed or reassessed for risk of extinction using the New Zealand Threat Classification System. Lists of threatened vascular plants were taken from a previous publication. Threatened fungi were not reassessed. The number of species listed as threatened increased from 2372 to 2788, and the number listed as Data Deficient increased from 2047 to 3031. These changes were mostly as a result of improved knowledge and the assessment of species that had not been considered previously, but a small number of species changed category as a result of genuinely improved or deteriorated status. Each of these changes is discussed, and complete lists of threatened and Data Deficient species are provided.

Keywords: Threat Classification System, threatened species, New Zealand

# Introduction

The New Zealand Threat Classification System developed by Molloy et al. (2002) was used to list nominated native New Zealand taxa according to threat of extinction. These lists are provided in Appendix 1.

## Interpreting these lists—the importance of qualifiers

The qualifiers listed for many species provide additional information that adds meaning to the threat classification. The qualifiers are an integral part of the classification of each taxon. When using these listings to inform management decisions, these factors should be taken into account.

For example, all taxa listed as Nationally Critical have a very high risk of extinction in New Zealand because of extremely small known populations or areas of occupancy, and/or very high rates of decline. However, some of the listings carry qualifiers that indicate that they are likely to be of lower priority for recovery action, e.g. SO (Secure Overseas), ST (Stable), or RC (Recovering).

**Example:** Among marine mammals, four taxa are listed as Nationally Critical: southern elephant seal (*Mirounga leonina*), orca/killer whale (*Orcinus orca*), Brydes whale (*Balaenoptera edeni/brydei*), and Maui's dolphin (*Cephalorhynchus hectori māui*). All are believed to have New Zealand populations of less than 250 breeding adults. However, they differ very significantly in the associated qualifiers:

- Maui's dolphin has the qualifiers CD (Conservation Dependent) and HI (Human Induced loss of range). The absence of an SO or TO (Threatened Overseas) qualifier means that it is endemic.
- The southern elephant seal has the qualifiers S?O (the questionmark indicating uncertainty about how secure the taxon is overseas) and HI. The Australasian population, of which the New Zealand animals are a small part, is in severe decline, but the overall world population now appears roughly stable.
- In contrast, the Brydes and killer whales both have the qualifiers SO and DP (Data Poor), indicating that they are not considered endangered on a world scale and the information on their population sizes could be improved.

# The process

1. Questionnaires (Appendix 2) requesting information on numbers, population trends and the geographic range of threatened and potentially threatened taxa (i.e. species, subspecies, and undescribed and tag-named taxa) were sent to many experts both within and outside the New Zealand Department of Conservation (DOC).
2. The taxa were divided into broad taxonomic groups (e.g. birds, terrestrial invertebrates and vascular plants). An expert panel for each taxonomic group met late in 2004 for one or more days, depending on the number of species to be discussed. (Members of the expert panels are listed in Appendix 3.) Exceptions were vascular plants, for which a review had occurred in association with the publication of a threatened plant list in the ‘New Zealand Journal of Botany’ by de Lange et al. (2004), and fungi, for which new information and expert time were not currently available to allow a review of the listings. Each taxon for which information was available in the form of either a completed questionnaire, a recent publication on its status or unpublished information that was known to panel members was assessed by the panel according to the criteria outlined in the New Zealand Threat Classification System (Molloy et al. 2002). Rod Hitchmough and Leigh Bull facilitated all meetings to ensure that, as far as possible, the criteria were applied consistently across all groups. Decisions on threat listings were recorded electronically as they were made and were projected on screen for checking by the panel as they were recorded. Notes were recorded about the information on which the listings were based. Records were also kept of all changes in formal threat status since the previous listing.
3. The listing process was either completed at the panel meetings or required various degrees of follow-up, according to the amount of information available at the meeting. Considerable additional information was collected after the meetings for some lesser known groups. In these instances, this information and draft listings that were inferred from it were then circulated to panel members for their confirmation.
4. A draft list was then circulated both to all panel members and to DOC conservancies for checking for accuracy and identification of significant gaps.

# Additional information available in electronic but not published form

Appendix 1 of this document lists all threatened and Data Deficient species currently listed under the New Zealand Threat Classification System, with the exception of Data Deficient fungi, for which the lists are unchanged from those published by Hitchmough (comp. 2002). However, many other taxa have been assessed and been listed as Not Threatened, Migrant, Coloniser, Vagrant or Introduced and Naturalised. These additional listings and more background information are available electronically (see below). This information includes additional notes on distribution and status, criteria triggering listing, a record of the date on which the entry for that taxon was last modified, previous listings under other systems (Molloy & Davis 1992; the International Union for Conservation of Nature and Natural Resources (IUCN); and (for plants) de Lange et al. 1999), and synonyms (e.g. tag names that have now been superseded by a published formal name) that have been used in recent documents with cross-references to the name used in this listing.

In addition to the information published in Appendix 1 of this document, a computer spreadsheet with additional background information is available next to the pdf for this document on the Science Publications page of the DOC website ([www.doc.govt.nz](http://www.doc.govt.nz) > Publications > Science & technical). The same listings and additional background information about the species will shortly be available on the DOC BIOWEB database. The electronic database will be ‘live’ and will be updated as any significant changes in the status of taxa occur. If there has been a change since the last hard-copy publication of these lists, the database should be regarded as containing the more authoritative information. If further information is required, contact your local DOC conservancy office.

## Coverage

The system has the capacity to classify the status of all species, subspecies, etc. of organisms occurring in the wild in New Zealand. Every effort has been made to include as many species as possible in this list, but inevitably the completeness of coverage varies between taxonomic groups. We hope that taxa not presently covered will be included in future revisions of this list.

For all native vertebrate groups except marine fish, the entire fauna has had its status assessed and all known native taxa including non-threatened taxa are listed in the electronic database.

For vascular plants, all native taxa have been assessed, but non-threatened taxa are listed in the electronic database only if they have been considered threatened

at some stage in the past. Any native vascular plant not listed can be assumed to be Not Threatened.

For all other groups (bryophytes, algae, fungi, freshwater invertebrates, marine fish, marine invertebrates and terrestrial invertebrates), only taxa nominated as likely to be threatened have been assessed. Many more taxa in these groups are candidates for assessment in the future. The status of fungi has not been reviewed since the publication of the 2002 lists. Threatened fungi from that listing are included here, but for the large number of fungi listed as Data Deficient, see Hitchmough (comp. 2002) or the electronic database.

Introduced and Naturalised taxa are listed in the electronic database only if they are IUCN-listed in their country of origin.

Formally described and generally accepted (taxonomically determinate) taxa are separated in this published list from taxonomically indeterminate taxa that lack published names or are of disputed validity. It is desirable to avoid confusion about the identity of tag-named, undescribed taxa by linking them to numbered voucher specimens. This was achieved as far as possible, but for some groups this was not possible; for example, it was not possible for many arthropods, for which many collections do not use individual identification numbers. The intention is that in future revisions of this list, voucher numbers or other evidence, such as reference to a published or filed photograph, will be included for undescribed taxa where possible.

## Listing at one taxonomic level only

Taxa are listed only once. If subspecies are recognised or informally proposed in a species, then it is the subspecies that are listed. For example, the following endemic vertebrate species will not be found listed as species because they are listed at the subspecific level:

Long-tailed bat—*Chalinolobus tuberculata* (undescribed subspecies)

Short-tailed bat—*Mystacina tuberculata*

Hector's dolphin—*Cephalorhynchus hectori*

Rifleman—*Acanthisitta chloris*

Brown teal—*Anas chlorotis* (undescribed subspecies)

Bellbird—*Anthonis melanura*

Pipit—*Anthus novaeseelandiae*

Fernbird, karoti, matata—*Bowdleria punctata*

Kokako—*Callaeas cinerea*

Banded dotterel—*Charadrius bicinctus*

New Zealand dotterel—*Charadrius obscurus*

New Zealand snipe—*Coenocorypha aucklandica*

Red-crowned kakariki—*Cyanorhamphus novaezelandiae*  
New Zealand falcon—*Falco novaeseelandiae* (undescribed subspecies)  
Weka—*Gallirallus australis*  
Kaka—*Nestor meridionalis*  
New Zealand robin—*Petroica australis*  
Tomtit—*Petroica macrocephala*  
Saddleback, tieke—*Philesturnus carunculatus*  
Tui—*Prosthemadera novaeseelandiae*  
White-fronted tern—*Sterna striata*  
Marbled skink—*Cyclodina oliveri* (undescribed subspecies)  
Striped gecko—*Hoplodactylus stephensi* (undescribed subspecies)  
Auckland and Wellington green geckos—*Naultinus elegans*  
Common and Chatham Island skinks—*Oligosoma nigriplantare*  
Cook Strait and northern tuatara—*Sphenodon punctatus* (undescribed  
subspecies)

## Distribution of listings

Table 1 shows a breakdown of the numbers of species in each threat category, separated by taxonomic group. In most groups there are large numbers of Nationally Critical and Nationally Endangered species, relatively few that are listed as Nationally Vulnerable or in Serious Decline, and large numbers again in the Gradual Decline, Sparse, and Range Restricted categories. There are fewer taxa in the Nationally Vulnerable and Serious Decline categories because a combination of high rates of decline and moderate to large population sizes is required to trigger classification in these categories. Unless facing a new threat, most New Zealand taxa that are declining at these high rates will already have very low population sizes, triggering a higher threat listing. Taxa that have suffered high rates of decline in the past but have now stabilised in refugia will no longer trigger the trend criteria for these categories and will be listed as Sparse or Range Restricted.

## Conservancy distributions

The 2002 lists included a breakdown of where the taxa were found by DOC conservancies. However, in 2004–05 there were no resources available for the very time-consuming task of collecting and checking information on conservancy distributions of the nearly 7000 taxa that are currently listed. This information should soon become available via the ‘casual observations’ application of the DOC BIOWEB database. The recording of observations of threatened species in this database is strongly encouraged.

TABLE 1. DISTRIBUTIONS OF THREAT RANKINGS BY TAXONOMIC GROUP.

All groups, except fungi and vascular plants, were reviewed in 2004–05. Fungi listings were not reviewed and remain the same as in the 2002 listing; vascular plant listings were reviewed by de Lange et al. (2004) and have been taken from that document.

THREAT CATEGORY	TAXONOMIC GROUP						
	BATS	BIRDS	BRYOPHYTES	FRESHWATER FISH	FRESHWATER INVERTEBRATES	FROGS	FUNGI
Nationally Critical	21	69	1	11	2	49	1
Nationally Endangered	3	29	18	2	2	1	
Nationally Vulnerable	1	12	1	3	1		
Serious Decline	4			1			1
Gradual Decline	21			13	3		10
Sparse	8		33	2	11	1	5
Range Restricted	58	54	4	86			37
<b>Total threatened</b>	<b>5</b>	<b>153</b>	<b>175</b>	<b>26</b>	<b>114</b>	<b>4</b>	<b>65</b>
Data Deficient	1	1	51	3	27		38
Extinct (since 1840)	16		1	1		1447*	23

\* Unchanged since 2002 except for the removal of eight duplicate listings; not listed in this document.

*Table 1—continued*

THREAT CATEGORY	TAXONOMIC GROUP						
	MARINE FISH	MARINE INVERTEBRATES	MARINE MAMMALS	REPTILES	TERRESTRIAL INVERTEBRATES	VASCULAR PLANTS	TOTAL
Nationally Critical	10	4	5	5	125	85	383
Nationally Endangered	3	2	4	4	101	67	232
Nationally Vulnerable			1	1	11	23	53
Serious Decline	5		1	14	14	29	55
Gradual Decline	14		22	38	79		202
Sparse	4		12	115	140		349
Range Restricted	32	234	22	539	445		1514
<b>Total threatened</b>	<b>52</b>	<b>270</b>	<b>8</b>	<b>67</b>	<b>943</b>	<b>868</b>	<b>2788</b>
Data Deficient	34	8	14	10	1316	96	3031
Extinct (since 1840)			1	1	9	6	33

# Changes in the threatened category status of indigenous species since the previous listing exercise

The total number of threatened species reported in the listing exercise carried out in 2004–05 was 2788 (up by 416). There were net increases of 23 species listed in the Nationally Critical category, 32 species in Nationally Endangered and 10 species in Nationally Vulnerable, a reduction of 8 in the total listed as in Serious Decline, and increases of 23 in Gradual Decline, 72 in Sparse and 264 in Range Restricted. These are net changes: in each category, some species were added and others were removed; these changes include both movements between categories and new additions to the list (see Table 2).

An additional 984 taxa were newly listed as Data Deficient—that is, likely to be threatened, but with too little information to fit them into a threatened category. This brings the total in this category to 3031.

The majority of the newly listed threatened and Data Deficient taxa were added as a result of new information about their status rather than a real sudden change in status. However, there were some genuine declines and recoveries. The status of five species is considered to have genuinely improved sufficiently over the past 3 years to have resulted in a changed threat classification. These are:

- Codfish Island fernbird (*Bowdleria punctata caudata*): Nationally Critical to Range Restricted. Originally found in only one location, its population has recovered from temporary reductions caused by the kiore eradication on Codfish Island; birds are now much more abundant than before the rat eradication took place. A second, healthy population has also been established on another pest-free island using translocation techniques.
- Crested grebe (*Podiceps cristatus australis*): Nationally Critical to Nationally Endangered. Survey results show that this species has increased in numbers. Predator control work has been undertaken at two important breeding sites (around Lake Pearson and on a nesting island on Lake Clearwater), along with the management of willows to improve nesting sites.
- Campbell mollymawk/New Zealand black-browed albatross (*Thalassarche impavida*): Nationally Vulnerable to Range Restricted. This species' recovery from earlier decline has been confirmed. This recovery shows a close correlation with the reduction in fishing intensity around New Zealand in the mid-1980s.
- Black petrel (*Procellaria parkinsoni*): Gradual Decline to Range Restricted. Although formerly in decline, there is no current evidence that this continues. Populations are found on Hauturu/Little Barrier Island and Great Barrier Island (Aotea Island). The eradication of cats on Hauturu/Little Barrier Island substantially improved the security of this sector of the population.

- *Therasia* sp. “chathamensis” (NMNZ M.52076): Gradual Decline to Range Restricted. Found only in dunes on the Chatham Islands, this snail’s decline has been reduced through management of the dunes’ ecosystem.

At the same time, 40 species are considered to have genuinely worsened in status sufficiently over the past 3 years to have resulted in a changed threat classification. These are:

- Bounty Island shag (*Leucocarbo ranfurlyi*): Range Restricted to Nationally Critical. This change in listing was due to increased concern about *Leucocarbo* shags in general. This species is a Bounty Island endemic and therefore is seldom surveyed. Furthermore, previous surveys used different methods, so may not be directly comparable. However, these surveys did indicate a substantial decline in numbers prior to 1997.
- Fiordland crested penguin (*Eudyptes pachyrhynchus*): Gradual Decline to Nationally Endangered. There has been a massive historical decline in this species since European settlement, and this is assumed to be continuing at some sites. A reconsideration of the total population size prompted revision of the listing. In the early 1990s, a population survey found 2260 nests, giving a conservative population estimate of less than 5000 mature individuals.
- Chatham Island shag (*Leucocarbo onslowi*): Range Restricted to Nationally Endangered. The low count in 2003 may reflect a poor breeding season rather than population decline, but the expert panel took a precautionary approach. The data need confirmation by repeated counts.
- Grey duck (*Anas superciliosa superciliosa*): Serious Decline to Nationally Endangered. Continued introgressive hybridisation with introduced mallard ducks is causing continued severe population decline.
- Black-fronted tern (*Sterna albostriata*): Serious Decline to Nationally Endangered. This change in listing was due to both improved knowledge and continuing decline. The species is very susceptible to predation and disturbance. The Ashburton River/Hakatere population decreased from 750 birds in 1981 to 200 birds in 1990.
- Pitt Island shag (*Stictocarbo featherstoni*): Range Restricted to Nationally Vulnerable. A survey in 2003 revealed a 25% decline over 6 years.
- Salvin’s mollymawk (*Thalassarche salvini*): Range Restricted to Nationally Vulnerable. This change in listing was due to a precautionary approach being taken by the expert panel and concern regarding high levels of bycatch. Population decline at the breeding site (Bounty Island) was recorded between 1978 and 1997, but the methods used in the counts may not be comparable.
- South Island rifleman (*Acanthisitta chloris chloris*): Not Threatened to Gradual Decline. This species has recently disappeared from Stewart Island/Rakiura, and anecdotal it also seems to have disappeared or be less conspicuous in some South Island lowland forest areas.
- North Island rifleman (*Acanthisitta chloris granti*): Not Threatened to Gradual Decline. This change was based on a general impression of declining conspicuousness.
- Red-billed gull (*Larus novaehollandiae scopulinus*): Not Threatened to Gradual Decline. Population decline documented at the three largest colonies (Three Kings, Mokohinau and Kaikoura) has been sufficient to meet the Gradual Decline criteria.

TABLE 2. SUMMARY OF STATUS CHANGES BETWEEN LISTINGS IN HITCHMOUGH (COMP. 2002) AND THOSE IN APPENDIX 1.

THREAT CATEGORY	TAXONOMIC GROUP														
	BATS			BIRDS			BRYOPHYTES			FRESHWATER FISH					
	2001	2004	IN	OUT	CHANGE	2001	2004	IN	OUT	CHANGE	2001	2004	IN	OUT	CHANGE
<b>THREAT DIVISION</b>															
Nationally Critical						25	21	1	5	-4	80	69	18	29	-11
Nationally Endangered	3	3	0	0	0	22	29	8	1	7	9	18	12	3	9
Nationally Vulnerable	1	1	0	0	0	9	12	4	1	3	1	1	3	1	2
Serious Decline						8	4	0	4	-4				2 <sup>a</sup>	1
Gradual Decline						20	21	3	2	1				10	13
Sparse						11	8	0	3	-3	5	33	30	2	28
Range Restricted	1	1	0	0	0	50	58	13	5	8	35	54	28	9	19
<b>Total threatened</b>	<b>5</b>	<b>5</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>145</b>	<b>153</b>	<b>29</b>	<b>21</b>	<b>8</b>	<b>129</b>	<b>175</b>	<b>89</b>	<b>43</b>	<b>46</b>
Data Deficient	1	1	0	0	0	0	1	1	0	1	14	51	39	2	37
<b>Total threatened &amp; Data Deficient</b>	<b>6</b>	<b>6</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>145</b>	<b>154</b>	<b>30</b>	<b>21</b>	<b>9</b>	<b>143</b>	<b>226</b>	<b>128</b>	<b>45</b>	<b>83</b>
Extinct (since 1840)						16	16	1	1	0				1	1
<b>THREAT LEVEL</b>															
Acutely Threatened	4	4	0	0	0	56	62	13	7	6	89	88	31	32	-1
Chronically Threatened	0	0	0	0	0	28	25	3	6	-3	0	0	0	0	12
At Risk	1	1	0	0	0	61	66	13	8	5	40	87	58	11	47

<sup>a</sup> Serious Decline freshwater fish included two species in Appendix 1 of Hitchmough (comp. 2002), but the summary in Table 1 listed only one species. This was because at taxonomic split was agreed during the publication process and the table was not updated.

*Continued on next page*

Table 2—continued

THREAT CATEGORY	TAXONOMIC GROUP											
	FRESHWATER INVERTEBRATES						MACROALGAE					
	2001	2004	IN	OUT	CHANGE		2001	2004	IN	OUT	CHANGE	
Nationally Critical	4	11	7	0	7		2	2	0	0	50	49
Nationally Endangered	2	2	0	2	1		1	0	0	0	-1 <sup>b</sup>	
Nationally Vulnerable	1	1	0	1							1	1
Serious Decline											0	0
Gradual Decline	3	3	0	3					10	10	0	0
Sparse	6	11	6	1	5		1	1	0	0	5	5
Range Restricted	38	86	54	6	48				0	0	0	0
<b>Total threatened</b>	<b>48</b>	<b>114</b>	<b>73</b>	<b>7</b>	<b>66</b>		<b>4</b>	<b>4</b>	<b>0</b>	<b>0</b>	<b>66</b>	<b>65</b>
Data Deficient	20	27	10	3	7						-1	<b>36</b>
<b>Total threatened &amp; Data Deficient</b>	<b>68</b>	<b>141</b>	<b>83</b>	<b>10</b>	<b>73</b>		<b>4</b>	<b>4</b>	<b>0</b>	<b>0</b>	<b>1455</b>	<b>1447</b>
Extinct (since 1840)											8 <sup>c</sup>	23
											0	0
											59	<b>61</b>
											2	0
											2	2
THREAT DIVISION												
Acutely Threatened	4	14	10	0	10		3	3	0	0	50	49
Chronically Threatened	0	3	3	0	3		0	0	0	0	11	11
At Risk	44	97	60	7	53		1	1	0	0	5	5
											36	37
											1	0

<sup>b</sup> Nationally Critical fungi have been reduced by one species because of the removal of a double listing that was included in Hitchmough (comp. 2002).

<sup>c</sup> Data Deficient fungi have been reduced by eight species because of the removal of double listings (these were edited out of the published list in Hitchmough (comp. 2002), but were included in the statistics in Table 1 of that publication).

Table 2—continued

THREAT CATEGORY	TAXONOMIC GROUP																			
	MARINE FISH						MARINE INVERTEBRATES													
	2001	2004	IN	OUT	CHANGE	2001	2004	IN	OUT	CHANGE	2001	2004	IN	OUT	CHANGE					
<b>THREAT DIVISION</b>																				
Nationally Critical			12	10	0	2	-2	4	4	0	0	4	5	3	2	1				
Nationally Endangered			3	3		0		1	2	1	0	1	4	4	2	2	0			
Nationally Vulnerable								1	0	1	-1	0	1	1	0	1				
Serious Decline			5	5		0						1	1	0	0	0				
Gradual Decline	2	2	0	0	0	5	14	9	0	9		15	22	7	0	7				
Sparse	17	18	1	0	1	4	4	0				16	12	1	5	-4				
Range Restricted	32	32	2	2	0	215	234	21	2	19	1	2	1	0	19	22	4	1	3	
<b>Total threatened</b>	<b>51</b>	<b>52</b>	<b>3</b>	<b>2</b>	<b>1</b>	<b>244</b>	<b>270</b>	<b>30</b>	<b>4</b>	<b>26</b>	<b>7</b>	<b>8</b>	<b>2</b>	<b>1</b>	<b>59</b>	<b>67</b>	<b>18</b>	<b>10</b>	<b>8</b>	
Data Deficient	36	34	7	9	-2	2	8	6	0	6	12	14	2	0	2	8	10	7	5	2
<b>Total threatened &amp; Data Deficient</b>	<b>87</b>	<b>86</b>	<b>10</b>	<b>11</b>	<b>-1</b>	<b>246</b>	<b>278</b>	<b>36</b>	<b>4</b>	<b>32</b>	<b>19</b>	<b>22</b>	<b>4</b>	<b>1</b>	<b>3</b>	<b>67</b>	<b>77</b>	<b>25</b>	<b>15</b>	<b>10</b>
Extinct (since 1840)												1	1	0	0					
<b>THREAT DIVISION</b>																				
Acutely Threatened	0	0	0	0	0	15	13	0	2	-2	6	6	1	1	0	8	10	6	4	2
Chronically Threatened	2	2	0	0	0	10	19	9	0	9	0	0	0	0	0	16	23	7	0	7
At Risk	49	50	3	2	1	219	238	21	2	19	1	2	1	0	1	35	34	5	6	-1

Continued on next page

Table 2—continued

THREAT CATEGORY	TAXONOMIC GROUP														
	TERRESTRIAL INVERTEBRATES						VASCULAR PLANTS								
	2001	2004	IN	OUT	CHANGE	2001	2004	IN	OUT	CHANGE	2001	2004	IN	OUT	CHANGE
<b>THREAT DIVISION</b>															
Nationally Critical	89	125	45	9	36	89	85	3	7	-4	360	383	78	55	23
Nationally Endangered	87	101	19	5	14	68	67	0	1	-1	200	232	44	12	32
Nationally Vulnerable	8	11	3	0	3	23	23	1	1	0	43	53	14	4	10
Serious Decline	16	14	2	4	-2	30	29	2	3	-1	63	55	4	12	-8
Gradual Decline	33 <sup>d</sup>	38	9	4	5	84	79	2	7	-5	179	202	38	15	23
Sparse	66	115	62	13	49	144	140	5	9	-4	277	349	106	34	72
Range Restricted	370 <sup>e</sup>	539	244	75	169	451 <sup>f</sup>	445	16	22	-6	1250	1514	386	122	264
<b>Total threatened</b>	<b>669</b>	<b>943</b>	<b>384</b>	<b>110</b>	<b>274</b>	<b>889</b>	<b>868</b>	<b>29</b>	<b>50</b>	<b>-21</b>	<b>2372</b>	<b>2788</b>	<b>670</b>	<b>254</b>	<b>416</b>
Data Deficient	371 <sup>f</sup>	1316	1001	56	945	100 <sup>g</sup>	96	5	9	-4	2047	3031	1079	95	984
<b>Total threatened &amp; Data Deficient</b>	<b>1040</b>	<b>2259</b>	<b>1385</b>	<b>166</b>	<b>1219</b>	<b>989</b>	<b>964</b>	<b>34</b>	<b>59</b>	<b>-25</b>	<b>4419</b>	<b>5819</b>	<b>1749</b>	<b>349</b>	<b>1400</b>
Extinct (since 1840)	4	9	6	1	5	6	6	0	0	0	28	33	7	2	5
<b>THREAT CATEGORY</b>															
Acutely Threatened	184	237	67	14	53	180	175	4	9	-5	603	668	136	71	65
Chronically Threatened	49	52	11	8	3	114	108	4	10	-6	242	257	42	27	15
At Risk	436	654	306	88	218	595	585	21	31	-10	1527	1863	492	156	336

<sup>d</sup> Gradual Decline terrestrial invertebrates included 33 species in Appendix 1 of Hitchmough (comp. 2002), but the summary in Table 1 listed 34 species. This was because of a simple counting error in the table.

<sup>e</sup> Range Restricted terrestrial invertebrates 2001 is as per Hitchmough (comp. 2002). Subsequent to this publication, 73 spiders were added to a working spreadsheet, and *Panoa isolata* was moved from Range Restricted to Data Deficient; these changes are included in the 2004 totals.

<sup>f</sup> Data Deficient terrestrial invertebrates 2001 is as per Hitchmough (comp. 2002). Subsequent to this publication, 766 spiders were added to a working spreadsheet; these additions are included in the 2004 totals.

<sup>g</sup> Range Restricted vascular plants included 451 taxa in Appendix 1 of Hitchmough (comp. 2002), but the summary in Table 1 listed 452 taxa. This was because of a simple counting error in the table.

<sup>h</sup> Data Deficient vascular plants included 100 taxa in Appendix 1 of Hitchmough (comp. 2002), but the summary in Table 1 listed 102 taxa. This was because of a simple counting error in the table.

- Moss (*Chorisodontium aciphyllum*): Range Restricted to Nationally Endangered. The Stockton population (one of four known populations) was extirpated by mining since the last listing (2001). Confined to Stockton and Denniston Plateaus.
- Moss (*Seligeria diminuta*): Range Restricted to Nationally Endangered. This Canterbury limestone endemic has been impacted by rock climbing at Castle Hill and by habitat modification associated with filming.
- Dune lakes galaxias (*Galaxias* sp.): Serious Decline to Nationally Vulnerable. There are two surviving populations at Kai Iwi lakes. Extirpation of the third population has now been confirmed; it has not been seen for more than a decade. The species is vulnerable to further exotic fish introductions and to water quality deterioration.
- Eldon's galaxias (*Galaxias eldoni*): Gradual Decline to Nationally Vulnerable. This change is based on both real decline and new knowledge. This species is known from 12 streams: five small populations are vulnerable to water quality changes resulting from farming, four to forestry impacts, and one to invasion of the habitat by predatory trout; two populations are secure. Restoration and habitat protection is occurring on some streams.
- Gollum galaxias (*Galaxias gollumoides*): Not Threatened to Gradual Decline. This change results from more information being gathered and from an increase in threats to water quality related to land use. The species is widespread (29 known populations), but few populations are secure (Catlins and Stewart Island/Rakiura). The species is vulnerable to predation by trout.
- Upland longjaw galaxias (*Galaxias prognathus*): Sparse to Gradual Decline. Decline has been observed when revisiting known sites and potential threats to water quality have increased. It is known from three or four catchments only. Local vulnerability/declines may get worse if land use intensifies and impacts water quality following tenure review. This species was referred to previously as the longjaw galaxias.
- “Southern” galaxias (*Galaxias* “Southern sp.”): Data Deficient to Gradual Decline. This change results from more information being gathered and from an increase in threats to water quality related to land use. Habitat degradation is causing decline in Southland. There are secure populations on Stewart Island/Rakiura. This southern taxon was previously thought to be part of the range of the flathead galaxias, but is now thought likely to be a distinct species.
- Grand skink (*Oligosoma grande*) and Otago skink (*Oligosoma otagense*): both Nationally Endangered to Nationally Critical. Population modelling based on a long-term mark-recapture study at the main stronghold indicated a severe threat of extinction within 10 years from 2003 for both species. Small peripheral populations are also disappearing.
- Whitakers skink (*Cyclodina whitakeri*): Range Restricted to Nationally Vulnerable. A very large historical decline can be inferred from the relict current distribution compared with the subfossil distribution. Monitoring confirms a continued decline at Pukerua Bay (the only mainland site). There are two secure and healthy populations on offshore islands. The total area occupied is only about 20 ha. There are also three translocated populations. The predicted decline on the mainland is likely to be offset by increases in

translocated populations in the longer term, and captive management is under way.

- *Hoplodactylus* “Matapia Island”: Sparse to Gradual Decline. Previously thought to be confined to Matapia Island, this small gecko is now known to also occur on the mainland (Te Paki area and Karikari Peninsula). Mainland decline has been inferred due to destruction of habitat during coastal development (particularly on Karikari Peninsula) and predator impacts.
- Goldstripe gecko (*Hoplodactylus chrysosireticus*): Sparse to Gradual Decline. Found on Mana Island and in flax near the coast in Taranaki. A reinterpretation of existing information led the expert panel to infer that the mainland population is declining, because the population recovery on Mana Island following mouse eradication demonstrates the species’ vulnerability to mammalian predation.
- Harlequin gecko (*Hoplodactylus rakiurae*): Sparse to Gradual Decline. Decline was inferred from recent observations of reduced rates of sightings at a few sites. There is extensive unsurveyed habitat, exact habitat preferences are unknown, and threats and trends are unknown. However, the species is likely to be extremely slow breeding and slow maturing, and is often found in the open on the ground both by day and night, making it vulnerable to predation by mammals (cats, rats and possums), which is severely impacting bird species at the same sites.
- Ornate skink (*Cyclodina ornata*): Not Threatened to Gradual Decline. Found in lowland areas throughout the North Island. Although there are numerous secure island populations, there is concern about the level of mainland decline. This species had been lost from known sites in Northland—one of its main strongholds. Decline on the mainland may be partly offset by population increases on islands following pest eradication. Taxonomy of the Three Kings population requires attention.
- Aphid *Paradoxaphis aristoteliae* Sunde, 1987: Data Deficient to Nationally Critical. This aphid is known from only two sites. It has been repeatedly found at Dolamore Park, Gore, and was previously found at Lake Rotoroa, Nelson Lakes; however, new information indicates that it is no longer found at the latter site.
- Snail *Rhytidida oconnori* Powell, 1946: Nationally Endangered to Nationally Critical. This large snail is confined to marble areas on Canaan around Harwoods Track and upper Gorge Creek. Recent surveys indicate approximately 80% decline in the last 4 years.
- Spider *Zealoctenus cardronaensis* Forster & Wilton, 1974: Data Deficient to Nationally Critical. This species is known from only one female specimen collected in 1969 in tussock at high altitudes at the southern end of Cardrona Valley. Recent search effort (1999 and 2000) in Cardrona Valley found no new specimens.
- Forest ringlet butterfly (*Dodonidia helmsii*): Gradual Decline to Serious Decline. Conceted survey effort indicates that this butterfly has disappeared from lowland sites, despite plentiful habitat. Very small remnant populations remain at high altitude. Vespid wasp predation is a possible cause of the decline.

- Moth *Notoreas* “Wellington”: Gradual Decline to Serious Decline. This species is endemic to the south Wellington coast (seven sites between Titahi Bay, Wellington, and White Rock, eastern Wairarapa). It feeds only on the prostrate coastal plant *Pimelea* cf. *urvilleana*. The habitat of the plant and moth has been partly destroyed by a range of incursions including four-wheel-drive vehicles, off-road motorbikes and cattle trampling.
- Moth *Gingidiobora nebulosa* (Philpott, 1917): Sparse to Gradual Decline. The sole host plant (*Gingidia montana*) is suffering browse reduction, which must be impacting on the moth population.
- Karikari tree weta (*Hemideina thoracica* chromosomes race  $2n=23,24$ ): Range Restricted to Gradual Decline. Habitat of this weta on Karikari Peninsula has been destroyed during coastal development, and this threat is continuing.
- Ground beetle *Mecodema howitti* Castelnau, 1867: Range Restricted to Gradual Decline. This eastern Banks Peninsula endemic lives in forest remnants and old logs in pasture. These are being lost (rotting away or being cleared and burnt), causing decline.
- Large land snail *Powelliphanta* “Haast”: Range Restricted to Gradual Decline. Good data indicate that high levels of thrush predation at Marks Range colonies are causing decline.
- *Cortaderia turbaria*: Nationally Endangered to Nationally Critical. The wild population now stands at 150 individuals. This reflects a drop of 60%–65% since 1999, and further losses are anticipated. The main threat comes from a root fungus—*Fusarium* wilt—which was first identified in the northern populations 1–2 years ago. It has now eliminated at least one natural population and is taking out another planted one. As yet, the fungus cannot be controlled once it has infected a plant, and it is probably the same fungus that has made cultivation of *C. turbaria* difficult. At present, it is unclear whether *Fusarium* is a recent arrival; it possibly arrived in cultivated *C. turbaria* stock and then spread from the nursery and/or in plantings. It is also unknown which wild populations are infected/clean. Imminent extinction is a strong possibility, but some conservation measures to halt the pathogen’s spread are now in place.
- *Hebe societatis*: Range Restricted to Nationally Critical. This status change reflects the very small area occupied and very high risk rather than current decline. In 2002, all available alpine habitat on the mountain was surveyed. However, this species was not found anywhere other than at the initial discovery site of 2 years previously. The estimated area of occupancy is 100 m × 200 m and the estimated number of plants is c. 1500. No loss of plants is apparent, although the slopes nearby are being seriously damaged by pigs. Pigs could easily turn over the whole 2 ha of *Hebe* habitat.
- *Puccinellia walkeri* subsp. *chathamica*: Range Restricted to Nationally Critical. This species is found on the Chatham and Auckland Islands. We know its situation well on the Chatham Islands, where c. 900 plants cover 0.1 ha. Threats come from the unrestricted access of browsing animals to the beaches it occupies, competition from weeds, and habitat loss via erosion, etc. The habitat size places it into the Nationally Critical category. However, we do not know its exact status on the Auckland Islands. A botanist who knows the plant well recently spent a month visiting most accessible coastal portions of the Auckland Islands and saw very little of it.

- *Leptinella rotundata*: Gradual Decline to Nationally Vulnerable. One of the three remaining populations (at South Head) had virtually disappeared in April 2002 through the collapse of a sea cliff and the spread of grasses.
- *Tetrachondra hamiltonii*: Gradual Decline to Serious Decline. This species is under threat from weeds spreading into turf habitats throughout its range. It is found at one site in the North Island, and at scattered sites in the South Island and on Stewart Island/Rakiura (mainly western). This species, which is endemic to New Zealand, is one of only two species in the family.
- *Leucogenes tarabaoa*: Range Restricted to Serious Decline. This species is endemic to the Mount Peel Range near Geraldine. Its situation has deteriorated through the spread of browsing animals and habitat deterioration partly caused by the impact of trampers.
- *Jovellana sinclairii*: Range Restricted to Gradual Decline. This species is endemic to the central North Island—Hicks Bay and Taumarunui south to the Ruahine Range. It is found on fertile river flats in forested areas and is threatened by browsing animals (particularly goats), competition from weeds and habitat fragmentation. Surveys in 2002 failed to find it at many of its former sites.

Other species may have substantially improved or deteriorated in status without triggering the criteria to change category.

## Extinction

Sixteen birds, one freshwater fish, one reptile, nine terrestrial invertebrates and six vascular plants are listed as having become extinct since 1840. This has changed since the last listing 3 years ago: one bird and six invertebrates have been added to the Extinct category, and one bird and one invertebrate have been removed. Each of the additions was of a species with no confirmed reports for many years; the additional 3 years of no records, despite searching, sufficiently increased the confidence that the taxon was extinct for the expert panel to list it as such—in each case, extinction probably actually occurred many years ago. The seven species newly listed as Extinct were:

- South Island kokako (*Callaeas cinerea cinerea*), for which there have been no confirmed sightings for 45 years.
- Snail (“Ellettia” “aucklandensis” (NMNZ M.127901)) and ground beetle (*Mecodema costellum* “spelles”). For both species, extinction probably predates 1840.
- Snail (*Zelandiscus elevata*). This species is very poorly known—the shell material in collections is possibly subfossil, but this has yet to be confirmed.
- Ground beetle (*Mecodema punctellum*). This species is known only from Stephens Island (Takapourewa) and has not been seen since 1931.
- Akaroa weevil (*Megacolabus sculpturatus*). This species was listed as Extinct by Molloy & Davis (1994); recent intensive pitfall trapping has failed to find any specimens.

- Flax snail (*Placostylus (Maoristylus) ambagiosus* “Kohuronaki”). This species has not been seen alive for several decades (although fresh shell fragments were found in the early 1990s) and there is no new evidence of survival.

One snail (*Zelandiscus worthyi*) was removed from the list of Extinct species, but may well in fact be extinct. However, the panel formed the opinion that it was too poorly known to state this with confidence.

The high-profile discovery of a population of New Zealand storm petrels (*Oceanites maorianus*) in the Hauraki Gulf, has resulted in this species being removed from the Extinct lists. Initially the panel was not certain that the new discovery was of the same species; therefore, the New Zealand storm petrel was still listed as Extinct and the recently discovered population was listed separately as Data Deficient. However, morphological data presented recently resulted in the panel merging these listings, and moving the New Zealand storm petrel from Extinct to Data Deficient.

The Canterbury knobbled weevil *Hadramphus tuberculatus* was a potential addition to the list of Extinct species, but instead changed status from Nationally Critical to Nationally Endangered at the last minute. It was not seen between 1922 and early 2005, when it was rediscovered at Burke’s Pass; it was previously considered extinct by many authorities (it is IUCN listed EX as *Karocolens tuberculatus*). There had been frequent unsuccessful survey efforts prior to its rediscovery. In 2002, although considered very likely to be extinct, some doubt remained; therefore, it was conservatively listed as Nationally Critical. A draft listing of Extinct had been agreed at the 2004 panel meeting because continued surveys had failed to locate specimens of this large, distinctive, conspicuous insect; however, this listing was changed to Nationally Endangered as a result of its rediscovery.

## Updating listings

The intention is that this list will be fully reviewed every 3 years and that any major changes in risk of extinction will be recorded in the electronic database as they occur. Suggestions for minor corrections will be stockpiled for consideration in the triennial review of the listings and should be sent to:

New Zealand Threat Classification System lists  
Terrestrial Conservation Unit  
Research Development and Improvement Division  
Department of Conservation  
PO Box 10420, The Terrace  
Wellington 6143  
New Zealand  
Fax (04) 471 3279

# Acknowledgements

Many people generously donated time and information to this project. In particular, the listing would have been impossible without the information provided in completed questionnaires on threatened species. The expert panel members listed in Appendix 3 made the panel meetings enjoyable, informative and productive.

# References

- de Lange, P.J.; Heenan, P.B.; Given, D.; Norton, D.A.; Ogle, C.C.; Johnson, P.N.; Cameron, E.K. 1999: Threatened and uncommon plants of New Zealand. *New Zealand Journal of Botany* 37: 603–628.
- de Lange, P.J.; Norton, D.A.; Heenan, P.B.; Courtney, S.P.; Molloy, B.P.J.; Ogle, C.C.; Rance, B.D.; Johnson, P.N.; Hitchmough, R. 2004: Threatened and uncommon plants of New Zealand. *New Zealand Journal of Botany* 42: 45–76.
- Hitchmough, R. (comp.) 2002: New Zealand Threat Classification System lists 2002. *Threatened Species Occasional Publication* 23. Department of Conservation, Wellington. 210 p. Electronic version available from the DOC website: [www.doc.govt.nz](http://www.doc.govt.nz).
- Molloy, J.; Davis, A. 1992: Setting priorities for the conservation of New Zealand's threatened plants and animals. Department of Conservation, Wellington. 44 p.
- Molloy, J.; Davis, A. 1994: Setting priorities for the conservation of New Zealand's threatened plants and animals. 2nd edition. Collated by C. Tisdall. Department of Conservation, Wellington. 64 p.
- Molloy, J.; Bell, B.; Clout, M.; de Lange, P.; Gibbs, G.; Given, D.; Norton, D.; Smith, N.; Stephens, T. 2002: Classifying species according to threat of extinction—a system for New Zealand. *Threatened Species Occasional Publication* 22. Biodiversity Recovery Unit, Department of Conservation, Wellington. 26 p. Electronic version available from the DOC website: [www.doc.govt.nz](http://www.doc.govt.nz).

# Appendix 1

## TAXONOMIC LISTS BY THREAT CATEGORY

Taxa (as classified in 2004) are listed by threat category. Qualifiers are given where appropriate. International Union for Conservation of Nature and Natural Resources (IUCN) listings are also shown where applicable.

### Extinct

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
<b>BIRDS</b>			
<b>Taxonomically determinate</b>			
Chatham Island bellbird	<i>Anthonis melanocephala</i>		
Chatham Island fernbird	<i>Bowdleria rufescens</i>		
Chatham rail	<i>Cabalus modestus</i>		
South Island kokako	<i>Callaeas cinerea cinerea</i>		
Little Barrier Island snipe	<i>Coenocorypha aucklandica barrierensis</i>		NT (for <i>C. aucklandica</i> s.l. <sup>c</sup> )
Stewart Island snipe	<i>Coenocorypha aucklandica tredalei</i>		NT (for <i>C. aucklandica</i> s.l. <sup>c</sup> )
New Zealand quail	<i>Coturnix novaezelandiae</i>		
Dieffenbach's rail	<i>Gallirallus dieffenbachii</i>		
Huia	<i>Heteralocha acutirostris</i>		
New Zealand little bittern	<i>Ixobrychus novaezelandiae</i>		
Auckland Island merganser	<i>Mergus australis</i>		
Laughing owl	<i>Sceloglaux albifacies</i>		
Stephens Island wren	<i>Traversia lyalli</i>		
South Island piopio	<i>Turnagra capensis</i>		
North Island piopio	<i>Turnagra tanagra</i>		
Bush wren	<i>Xenicus longipes</i>		
<b>FRESHWATER FISH</b>			
<b>Taxonomically determinate</b>			
Grayling	<i>Prototroctes oxyrhynchus</i>	HI	EX
<b>REPTILES</b>			
<b>Taxonomically determinate</b>			
Kawekaweau	<i>Hoplodactylus delcourti</i>		EX
<b>TERRESTRIAL INVERTEBRATES</b>			
<b>Taxonomically determinate</b>			
Ground beetle	<i>Mecodema punctellum</i> Broun, 1921		EX
Akaroa weevil	<i>Megacolabus sculpturatus</i> Broun, 1893	OL	
Weevil	<i>Thotmus balli</i> Broun, 1911		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

<sup>c</sup> Subspecies likely to be elevated to species level in the future.

*Continued on next page*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Snail	<i>Zelandiscus elevata</i> (Climo, 1978)		
<b>Taxonomically indeterminate</b>			
Snail	“Ellettia” “aucklandensis” (NMNZ M.127901)	OL	
Native bee, manuscript name “tautahi”	<i>Leioproctus</i> “otautahi”		
Ground beetle	<i>Mecodema costellum</i> “spelles”		
Flax snail	<i>Placostylus (Maoristylus) ambagiosus</i> “Herangi Hill”		VU B1+2abcde (for <i>P. ambagiosus</i> s.l.)
Flax snail (pupuharakeke)	<i>Placostylus (Maoristylus) ambagiosus</i> “Kohuronaki”	OL	VU B1+2abcde (for <i>P. ambagiosus</i> s.l.)
<b>VASCULAR PLANTS</b>			
<b>Taxonomically determinate</b>			
	<i>Lepidium obtusatum</i> Kirk		
	<i>Logania depressa</i> Hook.f.		
	<i>Myosotis traversii</i> var. <i>cinerascens</i> (Petrie) L.B. Moore (WELT 2585)	DP	
	<i>Stellaria elatinoides</i> Hook.f.		
	<i>Trilepidea adamstii</i> (Cheeseman) Tiegh.		
<b>Taxonomically indeterminate</b>			
	<i>Pseudognaphalium</i> (a) (CHR 365358; Zoo)	DP	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

## Nationally Critical

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
<b>BATS</b>			
<b>Taxonomically indeterminate</b>			
Southern North Island southern short-tailed bat	<i>Mystacina tuberculata tuberculata</i> (southern North Island)	HI	VU A2c, C2a (for <i>M. tuberculata</i> s.l.)
<b>BIRDS</b>			
<b>Taxonomically determinate</b>			
Campbell Island teal	<i>Anas nesiotis</i>	CD, HI, OL	CR D
Okarito brown kiwi	<i>Apteryx rowi</i>	CD, RF, OL	CR C1
Southern New Zealand dotterel	<i>Charadrius obscurus obscurus</i>	CD, HI, OL	EN C1
Orange-fronted parakeet	<i>Cyanoramphus malherbi</i>	HI, EF	CR C1, C2 a(i)(ii), C2 b
White heron	<i>Egretta alba modesta</i>	ST, SO, OL	
White tern	<i>Gygis alba royanus</i>	SO, OL	
Chatham Island oystercatcher	<i>Haematopus chathamensis</i>	CD, RC	EN D1
Chatham Island pigeon, parea	<i>Hemiphaga chathamensis</i>	CD, RC, HI, OL	EN D
Black stilt	<i>Himantopus novaezelandiae</i>	CD, ST, HI, OL	CR C2 a(i), D
Bounty Island shag	<i>Leucocarbo ranfurlyi</i>	DP	CR B2 a,c(i)(ii)(iii)(iv)
Kermadec white-faced storm petrel	<i>Pelagodroma marina albiclinus</i>	DP, HI, OL?	
Black robin	<i>Petroica traversi</i>	ST, HI	EN D
Takahe	<i>Porphyrio hochstetteri</i>	CD, ST, HI	EN D
Taiko	<i>Pterodroma magentae</i>	CD, OL	CR A1 e B2 a, b(v) C1 C2 a(i) D
New Zealand fairy tern	<i>Sterna nereis davisae</i>	OL, CD, HI	
Kakapo	<i>Strigops habroptilus</i>	CD, HI	CR D
New Zealand shore plover, tuturuatu	<i>Thinornis novaeseelandiae</i>	ST, HI, CD	EN D
<b>Taxonomically indeterminate</b>			
South Island brown teal	<i>Anas chlorotis</i> "South Island"	HI, OL	EN A2 a,c
Haast tokoeka	<i>Apteryx Haast</i> "	RF, OL, CD	
Campbell Island snipe	<i>Coenocorypha Campbell</i> "	DP, OL	
Codfish Island South Georgian diving petrel	<i>Pelecanoides georgicus</i> "Codfish Island"	HI, OL	
<b>BRYOPHYTES</b>			
<b>Taxonomically determinate</b>			
Liverwort	<i>Acromastigum brachiphyllum</i>		
Liverwort	<i>Acromastigum verticale</i>		
Liverwort	<i>Allisoniella recurva</i>	DP	
Liverwort	<i>Allisoniella scottii</i>	DP	
Liverwort	<i>Anastrophyllum papillosum</i>	DP	
Liverwort	<i>Andrewsianthus bodgsonae</i>		
Liverwort	<i>Austroscyphus nitidissimus</i>	OL	
Moss	<i>Bartramia alaris</i> Dixon & Sainsbury	TO, DP	
Liverwort	<i>Bragginsella anomala</i>	OL	
Moss	<i>Bryum tenuidens</i> (Dixon & Sainsbury)	DP	
Moss	<i>Calliergidium austro-stramineum</i> (Mull. Hal.) E.B. Bartram	SO, OL	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Nationally Critical—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Liverwort	<i>Cheirolejeunea tenella</i>	OL	
Liverwort	<i>Cololejeunea cardiocarpa</i>	DP, SO, OL	
Liverwort	<i>Cololejeunea ellipsoidea</i>	DP, OL	
Liverwort	<i>Cololejeunea falcidentata</i>		
Liverwort	<i>Cololejeunea pulchella</i> var. <i>stylifera</i>		
Moss	<i>Crossidium davidae</i> Catcheside	SO, OL	
Moss	<i>Crossidium geheebei</i> (Broth.) Broth.	SO, OL	
Moss	<i>Cyclodictyon blumeanum</i> (Mull. Hal.) O. Kuntze	ST, SO, OL	
Moss	<i>Dicranoweisia spenceri</i> Dixon & Sainsbury	DP	
Moss	<i>Didymodon calycinus</i> Dixon	DP	
Moss	<i>Ditrichum brachycarpum</i> Hampe	TO, OL	
Moss	<i>Ditrichum rufo-aureum</i> (Hampe) Willis	OL, DP, TO	
Liverwort	<i>Dumontiera birsuta</i>	SO	
Moss	<i>Epipterygium opararensense</i> Fife & A.J. Shaw	OL	
Moss	<i>Erpodium glaucum</i> (Wilson) I.G. Stone	S?O	
Moss	<i>Goniomitrium acuminatum</i> Hook. & Wilson	SO	
Moss	<i>Grimmia plagiopoda</i> Hedw.	TO	
Moss	<i>Hampeella pallens</i> (Sande Lac.) M. Fleisch.	DP, SO	
Liverwort	<i>Herzobryum atrocapillum</i>	SO, OL	
Liverwort	<i>Herzogobryum filiforme</i>	S?O	
Liverwort	<i>Herzogobryum vermiculare</i>	SO, OL	
Liverwort	<i>Isolembidium anomalum</i> var. <i>anomalum</i>	SO	
Liverwort	<i>Isotachis westlandica</i>		
Liverwort	<i>Leiomitra julacea</i>	SO	
Liverwort	<i>Lejeunea cyanophora</i>		
Moss	<i>Lindbergia maritima</i> Lewinsky	OL	
Liverwort	<i>Lopbozia autoica</i>	DP	
Liverwort	<i>Lopbozia nivicola</i>	DP	
Liverwort	<i>Lopbozia pumicola</i>	OL	
Liverwort	<i>Lopbozia subalpina</i>	DP	
Moss	<i>Macromitrium angulatum</i> Mitt.	S?O, OL	
Liverwort	<i>Nephelejeunea talinayi</i> (S. Arnell) Grolle	OL, T?O	
Liverwort	<i>Pachyschistochila papillifera</i>	OL	
Liverwort	<i>Paracromastigum fiordlandiae</i>		
Liverwort	<i>Petalophyllum bodgsoniae</i>	OL	
Liverwort	<i>Phaeoceros hirticalyx</i>	S?O	
Moss	<i>Physcomitrella patens</i> subsp. <i>readeri</i> (Mull. Hal.) B.C. Tan	S?O	
Moss	<i>Physcomitrium pusillum</i> Hook.f. & Wilson	TO	
Liverwort	<i>Plagiochila baylisii</i>		
Liverwort	<i>Plagiochila fragmentisima</i>		
Liverwort	<i>Plagiochila batcheri</i>	OL	
Liverwort	<i>Plagiochila kermadecensis</i>	OL	
Moss	<i>Plagiotropus oederiana</i> (Sw.) H.A. Crum & L.E. Anderson	SO, OL	
Liverwort	<i>Ptychanthus stephensoniana</i>	DP, OL	
Liverwort	<i>Riccardia intercellula</i>	OL	
Liverwort	<i>Riccardia pseudodendroceros</i>	OL	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Nationally Critical—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Liverwort	<i>Riccardia umida</i>	OL	
Liverwort	<i>Schistochila pellucida</i>	DP	
Liverwort	<i>Schistochila pluriciliata</i>		
Moss	<i>Scorpiurium cucullatum</i> (Mitt.) Hedenäs	SO	
Liverwort	<i>Seppeltia succuba</i>	SO, OL	
Liverwort	<i>Telaranea exigua</i>	OL	
Liverwort	<i>Telaranea plumulosa</i>	S?O, OL	
Liverwort	<i>Temnoma angustifolium</i>		
Liverwort	<i>Xenothallus vulcanicus</i>		
<b>Taxonomically indeterminate</b>			
Liverwort	<i>Brevianthus</i> sp.	OL, DP	
Liverwort	<i>Frullania</i> "Radar Bush"	OL	
Liverwort	<i>Riccardia</i> aff. <i>pallidevirens</i>	SO	
<b>FRESHWATER FISH</b>			
<b>Taxonomically determinate</b>			
Lowland longjaw galaxias	<i>Galaxias cobitinis</i>	EF, HI	
<b>FRESHWATER INVERTEBRATES</b>			
<b>Taxonomically determinate</b>			
Snail	<i>Lymnaea tomentosa</i> (Pfeiffer, 1855)		
Caddisfly	<i>Oeconesus angustus</i> Ward, 1997	DP, OL	
Caddisfly	<i>Pseudoeconesus haasti</i> Ward, 1997	DP, OL	
Caddisfly	<i>Psilochorema spinibarpax</i> Ward, 1996	DP, OL	
Diving beetle	<i>Rhantus plantaris</i> Sharp	DP	
Diving beetle	<i>Rhantus schauinslandi</i> Ordish, 1989	DP	
Caddisfly	<i>Tipphobiosis binewai</i> Ward, 1995	DP, OL	
Caddisfly	<i>Tipphobiosis kuscheli</i> Wise, 1972	OL	
Caddisfly	<i>Tipphobiosis quadrifurca</i> Ward, 1997	DP, OL	
Caddisfly	<i>Tipphobiosis schmidi</i> Ward, 1998	DP, OL	
Caddisfly	<i>Tipphobiosis trifurca</i> McFarlane, 1981		
<b>FROGS</b>			
<b>Taxonomically determinate</b>			
Archie's frog	<i>Leiopelma archeyi</i>		CR A2ae
Hamilton's frog	<i>Leiopelma hamiltoni</i>	CD, ST, HI, OL	EN D
<b>FUNGI</b>			
<b>Taxonomically determinate</b>			
	<i>Austrogaster novaezelandiae</i>	OL	
	<i>Berggrenia cyclospora</i>	OL	
	<i>Cantharellus elsaes</i>	OL	
	<i>Chalciporus aurantiacus</i>		
	<i>Chlorovibrissea bicolor</i>	OL	
	<i>Chlorovibrissea melanochlora</i>	OL	
	<i>Chlorovibrissea tasmanica</i>	OL	
Fischer's egg	<i>Claustula fischeri</i> K.M. Curtis, 1926	DP, ST, TO, OL	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
	<i>Colpoma notbogagi</i>		
	<i>Cordierites acanthophora</i>	OL	
	<i>Dichomitus newhookii</i>		
	<i>Gomphus dingleyae</i>	OL	
	<i>Gomphus novaezelandiae</i>	OL	
	<i>Gyroporus castaneus</i>	OL	
	<i>Hysterangium youngii</i>	OL	
	<i>Inonotus chondromyelus</i>		
	<i>Lactarius maruitaensis</i>	OL	
	<i>Phallobata alba</i>		
	<i>Phanerochaete citrina</i>	OL	
	<i>Phanerochaete corymbata</i>	OL	
	<i>Phanerochaete luteoaurantiaca</i>	OL	
Septate-spored polypore	<i>Polyporus septosporus</i> P.K. Buchanan & Ryvarden, 1998	DP, TO, OL	
Chatham Island sow thistle rust	<i>Puccinia embergeriae</i> McKenzie & P.R. Johnst. ined.	ST, OL	
	<i>Puccinia freycinetiae</i>	OL	
	<i>Ramaria aureorbiza</i>	OL	
	<i>Ramaria avellaneovertex</i>	OL	
	<i>Ramaria basirobusta</i>	OL	
	<i>Ramaria junquilleovertex</i>	OL	
	<i>Ramaria piedmontiana</i>	OL	
	<i>Ramariopsis avellanea</i>	OL	
	<i>Ramariopsis avellaneoinversa</i>	OL	
	<i>Ramariopsis tortuosa</i>	OL	
	<i>Russula inquinata</i>	DP, OL	
	<i>Russula littoralis</i>	OL	
	<i>Russula miniata</i>	DP, OL	
	<i>Russula papakaiensis</i>	DP, OL	
	<i>Russula pleurogena</i>	OL	
	<i>Russula solitaria</i>	DP, OL	
Russula	<i>Russula vivida</i> McNabb, 1973	DP	
	<i>Sarcosoma orientale</i>		
	<i>Squamianita squarrulosa</i>	OL	
	<i>Thaxterogaster cartilagineus</i>	OL	
	<i>Uredo chathamica</i>	OL	
	<i>Uredo salicorniae</i>		
	<i>Volvariella surrecta</i>	OL	
	<i>Xylaria wellingtonensis</i>	OL	
	<i>Xylaria zealandica</i>	OL	
<b>Taxonomically indeterminate</b>			
Pukatea bracket	<i>Ganoderma</i> sp. “Awaroa”	HI	
	Undescribed genus (Trichocomaceae)	DP	
<b>MACROALGAE</b>			
<b>Taxonomically determinate</b>			
Red seaweed—Bangiales	<i>Dione arcuata</i>	OL	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
<b>MARINE INVERTEBRATES</b>			
<b>Taxonomically determinate</b>			
Polychaete worm	<i>Boccardia magniovata</i>		
Barnacle	<i>Chionelasmus darwini</i> (Pilsbry, 1907)	DP	
Octopus	<i>Cirroctopus hochbergi</i> O'Shea, 2000	DP, HI	
Barnacle	<i>Ibla idiotica</i> Batham, 1945		
Octopus	<i>Opisthoteuthis chathamensis</i> O'Shea, 2000	DP, HI	
Octopus	<i>Opisthoteuthis mero</i> O'Shea, 2000	DP, HI	
Echinoid	<i>Porterpygus kieri</i> Baker, 1984	DP	
Polychaete worm	<i>Spio aequalis</i>		
Barnacle	<i>Volcanolepas osbeai</i> (Buckeridge, 2000)	OL	
Sea daisy	<i>Xyloplax medusiformis</i> Baker, Rowe & Clark, 1987	DP	
<b>Taxonomically indeterminate</b>			
Giant deep clam	<i>Calyptogena</i> sp. (NZOI )		
<b>MARINE MAMMALS</b>			
<b>Taxonomically determinate</b>			
Bryde's whale	<i>Balaenoptera edeni/brydei</i>	SO, DP	DD
Maui's dolphin	<i>Cephalorhynchus hectori māui</i>	CD, HI	EN A1d, C1 (for <i>C. hectori</i> s.l.)
Southern elephant seal	<i>Mirounga leonina</i>	S?O, HI	
Killer whale	<i>Orcinus orca</i>	SO, DP	LR/cd
<b>REPTILES</b>			
<b>Taxonomically determinate</b>			
Grand skink	<i>Oligosoma grande</i>	CD, HI	VU C2a
Otago skink	<i>Oligosoma otagense</i>	CD, HI	VU C2a
<b>Taxonomically indeterminate</b>			
Open Bay Islands gecko	<i>Hoplodactylus</i> sp. "Open Bay Islands gecko"	DP, OL	
Coromandel striped gecko	<i>Hoplodactylus stephensi</i> Coromandel populations		
Open Bay Island skink	<i>Oligosoma</i> "Open Bay Island skink"	DP, OL	
<b>TERRESTRIAL INVERTEBRATES</b>			
<b>Taxonomically determinate</b>			
Moth	<i>Aletia cyanopetra</i> (Meyrick, 1927)	OL, DP	
Snail	<i>Alsolemia cresswelli</i> (Climo, 1978)	DP	
Nemertine worm	<i>Antiponemertes allisonae</i> (Moore, 1973)	OL	VU D2
Aphid	<i>Aphis coprosmae</i> Laing ex Tilyard		
Bird louse	<i>Apterygon okarito</i> Palma & Price, 2004		
Moth	<i>Archyala culta</i> Philpott, 1931	OL	
Moth	<i>Archyala opulenta</i> Philpott, 1926	DP	
Moth	<i>Asaphodes imperfecta</i> (Philpott, 1905)	DP, HI	
Moth	<i>Asaphodes obarata</i> F & R, 1875	DP, HI	
Stick insect	<i>Asteliaphasma naomi</i> (Salmon)	OL, DP	
Moth	<i>Australothitis volatilis</i> Matthews & Patrick, 1998	HI	
Moth	<i>Bityla pallida</i> (Hudson, 1905)	OL, DP	
Snail	<i>Chaureopa roscoei</i> Climo, 1985	DP	
Moth	<i>Chersadaula ochrogaster</i> Meyrick, 1923	OL	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Nationally Critical—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Snail	<i>Climocella pukanui</i> Goulstone & Brook, 1999	OL	
Bird louse	<i>Coloceras harrisoni</i> (Tendeiro, 1972)		
Moth	<i>Coridomorpha stella</i> Meyrick, 1914	DP	
Snail	<i>Cyrtora birtsutissima</i> (Powell, 1951)	RC, OL	DD
Snail	<i>Delouagapia tasmani</i>	OL	
Moth	<i>Elachista eurychora</i> (Meyrick, 1919)	DP, OL	
Moth	<i>Erechthias lychnopa</i> Meyrick, 1927		
Moth	<i>Euxoa cerapachodes</i> Guenée, 1868	DP, OL	
Snail	<i>Fectola melchior</i> Goulstone & Brook, 1999	OL	
Snail	<i>Flammoconcha cumberi</i> (Powell, 1941)	DP, OL	
Mokohinau stag beetle	<i>Geodorcus ithaginis</i> (Broun, 1893)	OL	
Spider—Cyatholipidae	<i>Hanea paturau</i> Forster, 1988	OL, DP	
Moth	<i>Heterocrossa maculata</i> (Philpott, 1927)	DP	
Open Bay Island leech	<i>Hirudobdella antipodum</i> (Benham, 1904)	OL	
Ground beetle	<i>Holcaspis abdita</i> Johns, 2004	OL	
Ground beetle	<i>Holcaspis bidentella</i> Johns, 2004		
Ground beetle	<i>Holcaspis brevicula</i> Butcher, 1984	OL	
Moth	<i>Izatha psychra</i> (Meyrick, 1883)	DP	
Moth	<i>Izatha rigescens</i> Meyrick, 1929	DP	
Nematode	<i>Longidorus waikouaitii</i> Yeates, Boag & Brown, 1997	DP, OL	
Weevil	<i>Lyperobius nesidiotes</i> Kuschel	ST, OL	
Ida Valley carabid	<i>Mecodema laeviceps</i> Broun, 1904		
Darkling beetle	<i>Menimus sinuatus</i> Broun, 1886	OL, DP	
Mercury Islands tusked weta	<i>Motuweta isolata</i> Johns, 1997	EF, OL	
Moth	<i>Orocrambus fugitivellus</i> (Hudson, 1950)	DP, OL	
Moth	<i>Orthoclydon pseudostinaria</i> (Hudson, 1918)		
Aphid	<i>Paradoxaphis aristoteliae</i> Sunde, 1987	DP	
Snail	<i>Pbrixgnathus transitans</i> Suter, 1892	OL	
Pimelea bug	<i>Pimeleocoris viridis</i> Eyles & Schuh, 2003	DP	
King Island turrent snail	<i>Placostylus (Basileostylus) bollonsi caperatus</i> Powell, 1948	RC, HI, OL	VU D2 (for <i>P. bollonsi</i> s.l.)
Flax snail (pupuharakeke)	<i>Placostylus (Maoristylus) ambagiosus ambagiosus</i> Suter, 1906	RC?, HI, OL	VU B1+2abcde (for <i>P. ambagiosus</i> s.l.)
Flax snail (pupuharakeke)	<i>Placostylus (Maoristylus) ambagiosus consobrinus</i> Powell, 1938	CD, RF, OL	VU B1+2abcde (for <i>P. ambagiosus</i> s.l.)
Flax snail (pupuharakeke)	<i>Placostylus (Maoristylus) ambagiosus keenorum</i> Powell, 1947	CD, ST, RF, OL	VU B1+2abcde (for <i>P. ambagiosus</i> s.l.)
Flax snail (pupuharakeke)	<i>Placostylus (Maoristylus) ambagiosus pandora</i> Powell, 1951	CD, HI, RF, OL	VU B1+2abcde (for <i>P. ambagiosus</i> s.l.)
Flax snail (pupuharakeke)	<i>Placostylus (Maoristylus) ambagiosus wattii</i> Powell, 1947	CD, HI, RF, OL	VU B1+2abcde (for <i>P. ambagiosus</i> s.l.)
Flax snail (pupuharakeke)	<i>Placostylus (Maoristylus) ambagiosus whareana</i> Powell, 1951	CD, HI, RF, OL	VU B1+2abcde (for <i>P. ambagiosus</i> s.l.)
Large land snail	<i>Powelliphanta gagei</i> (Powell, 1938)	DP, OL	
Large land snail	<i>Powelliphanta gilliesi brunnea</i> (Powell, 1938)	CD, HI, OL	DD (for <i>P. gilliesi</i> s.l.)
Large land snail	<i>Powelliphanta traversi otakia</i> (Powell, 1946)	CD, HI	DD (for <i>P. traversi</i> s.l.)
Harvestman	<i>Prasmiola unica</i> Forster, 1954	OL	
Slug	<i>Pseudaneitea ramsayi</i> Climo, 1973	RC, OL	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Darkling beetle	<i>Pseudobelops antipodensis</i>	OL	
Bird louse	<i>Quadraceps dominella</i> Timmermann, 1953		
Bird louse	<i>Quadraceps novaezealandiae</i> Timmermann, 1953		
Nematode	<i>Radopholus cavenessi</i> Egunjobi, 1968	DP, OL	
Bird louse	<i>Rallicola (Apteriscola) rodericki</i> Palma, 1991		
Bird louse	<i>Rallicola (Rallicola) takabe</i> Holloway, 1956		
Snail	<i>Rhytidia oconnori</i> Powell, 1946	OL	CR B1+2d
Snail	<i>Rhytidarex buddlei</i> (Powell, 1948)	OL	
Bird louse	<i>Saemundssonia (Saemundssonia) bathamensis</i> Timmermann, 1977		
Paua slug	<i>Schizoglossa gigantea</i> Powell, 1930	OL	
Alpine grasshopper	<i>Sigaus homerensis</i> Morris, 2003		
Moth	<i>Stathmopoda campylocha</i> Meyrick, 1889	OL	
Moth	<i>Titanomis sisyrota</i> Meyrick, 1888	DP	
Moth	<i>Trachypepla roseata</i> Philpott, 1923	OL, DP	
Moth	<i>Xanthorhoe bulbulata</i> (Guenée, 1868)	HI	
Pitt Island longhorn	<i>Xylotoles costatus</i> Pascoe, 1875	OL	EN D
Spider—Miturgidae	<i>Zealoctenus cardronaensis</i> Forster & Wilton, 1974	OL, DP	
Back Beach beetle	<i>Zecillenus tillyardi</i> (Brookes, 1927)		
<b>Taxonomically indeterminate</b>			
Moth	<i>Arctesthes</i> sp. "Von"	DP	
Leaf-vein slug	<i>Athoracophorid</i> "Mt Hikurangi"	OL	
Slug	<i>Atboracophorus</i> sp. 3 (NMNZ M.151429) "Waiare"	DP, OL	
Snail	<i>Cavellia</i> "Kohaihai" (NMNZ M.36649)	DP, OL	
Snail	<i>Charopidae</i> sp. 46 (NMNZ M.87828)	DP	
Snail	<i>Charopidae</i> sp. 164 (NMNZ M.88458)	OL	
A pleasing fungus beetle	<i>Cryptodacne</i> sp. "Chathams"		
Snail	<i>Cytora</i> sp. 11 (NMNZ M.87893)	HI, OL	
Snail	<i>Delos</i> sp. 1 (NMNZ M.29346)	HI, OL	
Snail	<i>Delos</i> sp. 13 (NMNZ M.29345)	HI, OL	
Snail	<i>Egestula</i> "broomfieldi" (NMNZ M.78965)	OL	
Snail	<i>Flammoconcha</i> "marstoni" (NMNZ M.22464)	OL, DP	
Stag beetle	<i>Geodorcus</i> sp. "Sisters"	DP, OL	
Moth	<i>Gracillariidae</i> n. sp. "teucridium"	DP	
Ground weta	<i>Hemiandrus</i> "Cape Campbell"	DP	
Ground beetle	<i>Holcaspis</i> n. sp.		
Moth	<i>Kiwaia</i> sp. "Cloudy Bay"	HI, EF, OL	
Native bee	<i>Leioproctus</i> "nunui"	HI, OL	
Bird louse	<i>Longimenopon</i> sp.		
Ground beetle	<i>Mecodema</i> sp. "Te Paki"	DP, OL	
Ground beetle	<i>Megadromus</i> sp. 8 "Omeo Hut"	OL	
Moth	<i>Meterana</i> "Foveaux Strait"	HI	
Moth	<i>Notoreas</i> "Castlepoint"	OL	
Moth	<i>Notoreas</i> "Cape Turnagain"	OL	
Moth	<i>Notoreas</i> "Mason Bay"	HI, OL	
Moth	<i>Notoreas</i> "Rahu Saddle"	DP, OL	
Moth	<i>Notoreas</i> "South Shag River"	OL	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Nationally Critical—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Moth	<i>Notoreas</i> "Waiho Flats"	HI, OL	
Bird louse	<i>Penentirmus</i> sp.		
Snail	<i>Pbrixgnathus</i> "wallacei" (NMNZ M.88229)		
King Island turrent snail	<i>Placostylus (Basileostylus) bollonsi</i> "West"	OL	VU D2 (for <i>P. bollonsi</i> s.l.)
Flax snail (pupuharakeke)	<i>Placostylus (Maoristylus) ambagiosus</i> "Haupatoto"	CD, HI, RF, OL	VU B1+2abcde (for <i>P. ambagiosus</i> s.l.)
Flax snail (pupuharakeke)	<i>Placostylus (Maoristylus) ambagiosus</i> "Kauaetewhakapeke Stream"	DP, OL	VU B1+2abcde (for <i>P. ambagiosus</i> s.l.)
Flax snail (pupuharakeke)	<i>Placostylus (Maoristylus) ambagiosus</i> "Poroiki"	CD, HI, RF, OL	VU B1+2abcde (for <i>P. ambagiosus</i> s.l.)
Flax snail (pupuharakeke)	<i>Placostylus (Maoristylus) ambagiosus</i> "Rangiora"	DP, HI, OL	VU B1+2abcde (for <i>P. ambagiosus</i> s.l.)
Flax snail (pupuharakeke)	<i>Placostylus (Maoristylus) ambagiosus</i> "Tapotupotu"	OL	VU B1+2abcde (for <i>P. ambagiosus</i> s.l.)
Flax snail (pupuharakeke)	<i>Placostylus (Maoristylus) ambagiosus</i> "Te Paki"	CD, ST, OL	VU B1+2abcde (for <i>P. ambagiosus</i> s.l.)
Flax snail (pupuharakeke)	<i>Placostylus (Maoristylus) ambagiosus</i> "Tirikawa Coast"	HI, OL	VU B1+2abcde (for <i>P. ambagiosus</i> s.l.)
Flax snail (pupuharakeke)	<i>Placostylus (Maoristylus) ambagiosus</i> "Tirikawa Trig"	DP, HI, OL	VU B1+2abcde (for <i>P. ambagiosus</i> s.l.)
Large land snail	<i>Powelliphanta</i> "Anatoki Range"	DP, OL	
Large land snail	<i>Powelliphanta</i> "Baton"	DP, OL	
Large land snail	<i>Powelliphanta</i> "Mt Augustus"	DP, OL	
Large land snail	<i>Powelliphanta gilliesi</i> "Haidinger"	ST, HI, OL	DD (for <i>P. gilliesi</i> s.l.)
Large land snail	<i>Powelliphanta lignaria</i> "millertoni"	DP, OL	DD (for <i>P. lignaria</i> s.l.)
Snail	Punctidae sp. 6 (NMNZ M.151458)		
Snail	Punctidae sp. 27 (NMNZ M.79798)	DP	
Snail	Punctidae sp. 226 (NMNZ M.154908)	OL	
Moth	<i>Sabatinca</i> sp. "Secretary Island"	DP, OL	
Bird louse	<i>Saemundssonia (Puffinoecus)</i> sp.		
Moth	<i>Scythbris</i> sp. "stripe"	DP, OL	
Moth	<i>Stigmella</i> sp. "olearia"	DP	
Snail	<i>Wainutia</i> "Mount Tuhua"	DP, OL	
<b>VASCULAR PLANTS</b>			
<b>Taxonomically determinate</b>			
	<i>Acaena rorida</i> B.H. Macmill.	CD, OL	
	<i>Alectryon excelsus</i> subsp. <i>grandis</i> (Cheeseman) de Lange et E.K. Cameron	RC, OL	
	<i>Anzybas carsei</i> (Cheeseman) D.L. Jones et M.A. Clem.	CD, HI, RF, EF, OL	
	<i>Atriplex hollowayi</i> de Lange et D.A. Norton	CD, EF	
	<i>Brachyscome pinnata</i> Hook.f.	CD, OL	
	<i>Carex dolomitica</i> Heenan et de Lange	CD, OL	
	<i>Carmichaelia hollowayi</i> G. Simpson	CD, RF	
	<i>Carmichaelia muritai</i> (A.W. Purdie) Heenan	CD	CR D
	<i>Celmista macmabontii</i> Kirk var. <i>macmabontii</i>	ST	
	<i>Centipeda minima</i> (L.) A. Braun et Asch. subsp. <i>minima</i>	SO, EF	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
	<i>Ceratocephala pungens</i> Garn.-Jones	HI, EF	
	<i>Christella dentata</i> (Forssk.) Brownsey et Jermy s.s.	CD, SO, RF, OL, HI	
	<i>Chionthus puniceus</i> (G. Don) Sol. ex Lindl.	CD, HI, OL	EN D
	<i>Coprosma spathulata</i> subsp. <i>bikuruana</i> de Lange et Heenan	CD, HI, OL	
	<i>Cortaderia turbaria</i> Connor	CD, RF, OL	
	<i>Crassula bunua</i> A.P. Druce	HI	
	<i>Davallia tasmani</i> subsp. <i>cristata</i> von Konrat, Briggins et de Lange	CD, RF, OL, ST	
	<i>Gunnera hamiltonii</i> Kirk	CD, ST, RF	
	<i>Hebe breviracemosa</i> (W.R.B. Oliv.) Cockayne et Allan	CD, HI, OL	
	<i>Hebe societatis</i> Bayly et Kellow	OL, DP	
	<i>Lepidium banksii</i> Kirk	CD, HI	
	<i>Lepidium sisymbrioides</i> subsp. <i>matau</i> (Petrie) Thell.	CD, HI, EF	
	<i>Leptinella filiformis</i> (Hook.f.) D.G. Lloyd et C. Webb	EW, CD, HI	
	<i>Linguella puberula</i> Hook.f.	HI, EF	
	<i>Linum monogynum</i> var. <i>chathamicum</i> Cockayne (CHR 417633)	CD, EF	
	<i>Mazus novaezeelandiae</i> subsp. <i>impolitus</i> f. <i>birta</i> Heenan	CD, HI	
	<i>Metrosideros bartlettii</i> J.W. Dawson	CD, HI	EN D
	<i>Myosotis albosericea</i> Hook.f.	ST, OL	
	<i>Myosotis angustata</i> Cheeseman	ST, OL	
	<i>Myosotis australis</i> var. <i>lytteltonensis</i> Laing et A. Wall	CD, HI, EF	
	<i>Myosotis cheesemani</i> Petrie	DP, EF	
	<i>Myosotis petiolata</i> Hook.f. var. <i>petiolata</i>	DP, HI, OL	
	<i>Neopaxia drucei</i> Heenan	ST, OL	
	<i>Olearia gardneri</i> Heads	CD, HI, RF	
	<i>Pachycladon exilis</i> (Heenan) Heenan et A. Mitch.	CD, HI, OL	
	<i>Pennantia baylisiana</i> (W.R.B. Oliv.) G.T.S. Baylis	CD, RF, OL, ST	CR D
	<i>Pittosporum turneri</i> Petrie	CD, RC	LR/cd
	<i>Poa spanta</i> Edgar et Molloy	CD, OL	
	<i>Poa sudicola</i> Edgar	DP, ST	
	<i>Pomaderris apetala</i> subsp. <i>maritima</i> N.G. Walsh et F. Coates	CD, SO, RF	
	<i>Pterostylis micromega</i> Hook.f.	CD, HI, EF	
	<i>Puccinellia raroflorens</i> Edgar	CD, DP	
	<i>Puccinellia walkeri</i> subsp. <i>chathamica</i> (Cheeseman) Edgar	DP	
	<i>Sebaea ovata</i> (Labill.) R. Br.	CD, SO, HI, EF	
	<i>Sicyos australis</i> Endl.	CD, TO	
	<i>Tecomanthe speciosa</i> W.R.B. Oliv.	CD, RF, OL	
	<i>Thelymitra sansicia</i> Hatch	DP, EF	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
	<i>Wahlenbergia pygmaea</i> subsp. <i>tararua</i> J.A. Petterson	DP, ST	
<b>Taxonomically indeterminate</b>			
	<i>Botrychium</i> aff. <i>lunaria</i> (CHR 289336; NW Nelson)	TO	
	<i>Calochilus</i> aff. <i>herbaceus</i> (CHR 65825; Kaimaumau)	SO, EF	
	<i>Cardamine</i> (a) (CHR 500569; Awahokomo)	CD, EF, OL	
	<i>Cardamine</i> (c) (CHR 65058; Reporoa Bog)	DP	
	<i>Celmisia</i> aff. <i>gracilenta</i> (b) (CHR 469722; Mangaweka)	CD, OL	
	<i>Craspedia</i> (a) (CHR 511522; Clutha River)	HI, OL	
	<i>Craspedia</i> (b) (CHR 516324; Leatham)	CD, OL	
	<i>Craspedia</i> (i) (CHR 395643; Fyfe River)	OL	
	<i>Craspedia</i> (j) (CHR 516302; Lake Heron)	HI, OL	
	<i>Euphrasia</i> (a) (CHR 471903; "white")	EF, OL	
	<i>Festuca</i> aff. <i>novae-zelandiae</i> (AK 252541; Awahokomo)	CD, EF, OL	
	<i>Gentiana</i> aff. <i>astonii</i> (a) (CHR 529112; Mt Brown)		
	<i>Gentiana</i> aff. <i>astonii</i> (b) (CHR 529111; Pareora River)	OL	
	<i>Gentiana</i> aff. <i>astonii</i> (e) (CHR 542276; Manahune)	OL	
	<i>Gentiana</i> aff. <i>saxosa</i> (AK 7316; Charleston)	OL	
	<i>Hebe</i> aff. <i>bishopiana</i> (AK 202263; Hikurangi Swamp)	CD, HI, OL	
	<i>Hypsela</i> aff. <i>rivalis</i> (CHR 369981; Burgoo Stream)	EF, OL	
	<i>Isoetes</i> aff. <i>kirkii</i> (CHR 247118A; Lake Omapere)	OL	
	<i>Koeleria</i> aff. <i>novozelandica</i> (AK 252546; Awahokomo)	CD, OL	
	<i>Lepidium</i> aff. <i>oleraceum</i> (a) (AK 230459; Chatham Islands)	CD, OL	
	<i>Lepidium</i> aff. <i>oleraceum</i> (d) (AK 255607; Mangere)	EF, OL	
	<i>Leptinella</i> (a) (CHR 515297; Clutha River)		
	<i>Limosella</i> (b) (CHR 515038; Manutahi)	ST	
	<i>Melicytus</i> aff. <i>obovatus</i> (b) (AK 235617; Mt Burnett)	CD, OL	
	<i>Microtis</i> aff. <i>unifolia</i> (CHR 532775; Fox)	DP, OL	
	<i>Myosotis</i> (b) (CHR 386966; Mt Tapuae-O-Uenuku)	OL	
	<i>Olearia</i> aff. <i>odorata</i> (CHR 386084; Canterbury Plains)	RF	
	<i>Pachycladon</i> aff. <i>fastigiata</i> (CHR 279206; Chalk Range)		
	<i>Pimelea</i> (a) (CHR 495025; Turakina)	CD, HI	
	<i>Pimelea</i> aff. <i>aridula</i> (a) (CHR 282959; Te Mata Peak)	OL	
	<i>Pratia</i> aff. <i>angulata</i> (AK 212143; Woodhill)		
	<i>Ranunculus</i> (a) (AKU 19876; Hope)		
	<i>Ranunculus</i> aff. <i>royi</i> (CHR 513327; Waihao)	HI, OL	
	<i>Thelymitra</i> (a) (WELT 79140; Ahipara)	CD, DP, HI, EF	
	<i>Tmesipteris</i> aff. <i>tannensis</i> (CHR 496779; Banks Peninsula)	DP	
	<i>Trichomanes</i> (AK 252983; Kerikeri)	DP, OL	
	<i>Trisetum</i> aff. <i>lepidum</i> (AK 251835; Awahokomo)	CD, EF, OL	
	<i>Uncinia perplexa</i> Heenan et de Lange	CD, HI, OL	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

## Nationally Endangered

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
<b>BATS</b>			
<b>Taxonomically determinate</b>			
Northern short-tailed bat	<i>Mystacina tuberculata aupourica</i>	DP, HI	VU A2c, C2a (for <i>M. tuberculata</i> s.l.)
<b>Taxonomically indeterminate</b>			
Long-tailed bat (South Island)	<i>Chalinolobus tuberculata</i> (South Island)	DP, HI	VU A2bc (for <i>C. tuberculata</i> s.l.)
South Island southern short-tailed bat	<i>Mystacina tuberculata tuberculata</i> (South Island)	HI	VU A2c, C2a (for <i>M. tuberculata</i> s.l.)
<b>BIRDS</b>			
<b>Taxonomically determinate</b>			
Brown teal	<i>Anas chlorotis</i> "North Island"	HI, CD	EN A2 a,c
Grey duck	<i>Anas superciliosa superciliosa</i>	SO	
Australasian bittern	<i>Botaurus poiciloptilus</i>	TO, HI	
Stewart Island fernbird, matata	<i>Bowdleria punctata stewartiana</i>	HI	NT (for <i>B. punctata</i> s.l.)
North Island kokako	<i>Callaeas cinerea wilsoni</i>	CD, HI, RF	EN A1 a b c e B1 a b(i)(ii)(iii)(iv)(v) B2 a b(i)(ii)(iii)(iv)(v) C1
Forbes' parakeet	<i>Cyanoramphus forbesi</i>	CD, RC, HI, OL	EN C2 a(ii)
Eastern rockhopper penguin	<i>Eudyptes chrysocome filholi</i>	TO	
Fiordland crested penguin	<i>Eudyptes pachyrhynchus</i>	HI	VU C1 2,a(i)
Erect-crested penguin	<i>Eudyptes sclateri</i>		
North Island weka	<i>Gallirallus australis greyi</i>	HI, EF	NT (for <i>G. australis</i> s.l.)
Stewart Island weka	<i>Gallirallus australis scotti</i>	ST, HI	NT (for <i>G. australis</i> s.l.)
Blue duck, whio	<i>Hymenolaimus malachorhynchos</i>	HI	EN C1, 2A(i)
Chatham Island shag	<i>Leucocarbo onslowi</i>		CR B2 a,c(i)(ii)(iii)(iv)
Yellowhead, mohua	<i>Moboua ochrocephala</i>	HI, EF	EN A 2a,e
South Island kaka	<i>Nestor meridionalis meridionalis</i>	HI	EN A 2a,e
North Island kaka	<i>Nestor meridionalis septentrionalis</i>	HI	EN A 2a,e
Kea	<i>Nestor notabilis</i>		VU C 1, 2a(ii)
Stitchbird, hihi	<i>Notiomystis cincta</i>	HI, OL	VU D2
Stewart Island robin	<i>Petroica australis rakiura</i>	HI	
Chatham Island tit	<i>Petroica macrocephala chathamensis</i>	RC, HI	
South Island saddleback, tieke	<i>Philesturnus carunculatus carunculatus</i>	HI	LC
Crested grebe	<i>Podiceps cristatus australis</i>	SO, HI, RC	
Chatham Island tui	<i>Prosthemadera novaeseelandiae chathamensis</i>	ST, HI, OL	
Chatham petrel	<i>Pterodroma axillaris</i>	CD, OL, RC	CR B2 a,b(v)
Kermadec petrel	<i>Pterodroma neglecta</i>	SO, HI, ST	
Hutton's shearwater	<i>Puffinus buttoni</i>	HI, OL	EB B2 a,b(v)
Black-fronted tern	<i>Sterna albostriata</i>	HI	EN A2 e
Grey-headed mollymawk	<i>Thalassarche chrysostoma</i>	TO, OL	
<b>Taxonomically indeterminate</b>			
Southern falcon	<i>Falco novaeseelandiae</i> "southern"		NT

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Nationally Endangered—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
<b>BRYOPHYTES</b>			
<b>Taxonomically determinate</b>			
Moss	<i>Archidium elatum</i> Dixon & Sainsbury	DP	EN B1+2c
Liverwort	<i>Brevianthus flavus</i>	TO	
Liverwort	<i>Cephalolobus squarrosus</i>	DP	
Liverwort	<i>Chaetophyllopsis whitelockei</i>	S?O	
Moss	<i>Chorisodontium aciphyllum</i> (Hook.f. & Wilson) Broth.	SO, HI	
Moss	<i>Fissidens berteroii</i> (Mont.) Mull. Hal.	S?O	
Moss	<i>Fissidens integerrimus</i> Mitt.	S?O	
Moss	<i>Fissidens strictus</i> Hook.f. & Wilson	S?O	
Liverwort	<i>Goebelobryum unguiculatum</i>	HI, SO	
Liverwort	<i>Neogrollea notabilis</i>	S?O	
Liverwort	<i>Petalophyllum australe</i>		
Liverwort	<i>Petalophyllum preissei</i>	S?O	
Liverwort	<i>Radula papulosa</i>		
Liverwort	<i>Ricciocarpus natans</i>	SO, HI	
Moss	<i>Seligeria diminuta</i> (R. Br. bis) Dixon		
Liverwort	<i>Stenolejeunea acuminata</i>		
<b>Taxonomically indeterminate</b>			
Liverwort	<i>Cololejeunea</i> sp. 1		
Liverwort	<i>Siphonolejeunea</i> "rock"	HI	
<b>FRESHWATER FISH</b>			
<b>Taxonomically determinate</b>			
Canterbury mudfish	<i>Neochanna burrowsius</i>	HI	VU B1+2bcd
Northland mudfish	<i>Neochanna heleios</i>	DP, HI	
<b>FRESHWATER INVERTEBRATES</b>			
<b>Taxonomically determinate</b>			
Caddisfly	<i>Edpercivalia banksiensis</i> McFarlane, 1939	OL	
Caddisfly	<i>Hydrobiosis styx</i> McFarlane, 1951		
<b>FROGS</b>			
<b>Taxonomically indeterminate</b>			
Maud Island frog	<i>Leiopelma pakeka</i>	CD, RC, HI, OL	VU D2
<b>MARINE INVERTEBRATES</b>			
<b>Taxonomically determinate</b>			
Squid	<i>Idiotethis cordiformis</i> (Chun, 1908)	SO, HI	
Lamp shell	<i>Pumilus antiquatus</i>		
<b>Taxonomically indeterminate</b>			
Barnacle	<i>Ibla</i> sp.	DP, HI, OL	
<b>MARINE MAMMALS</b>			
<b>Taxonomically determinate</b>			
Hector's dolphin	<i>Cephalorhynchus hectori hectori</i>	CD	EN A1d, C1 (for <i>C. hectori</i> s.l.)
Southern right whale	<i>Eubalaena australis</i>	RC, SO, HI	LR/cd

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
<b>REPTILES</b>			
<b>Taxonomically determinate</b>			
Chevron skink	<i>Oligosoma bomalonotum</i>	HI	VU D2
<b>Taxonomically indeterminate</b>			
Southern North Island speckled skink	<i>Oligosoma</i> aff. <i>infrapunctatum</i> “Southern North Island”	HI, DP	
Spotted skink “Central Canterbury”	<i>Oligosoma</i> aff. <i>lineoocellatum</i> “Central Canterbury”	HI	
Brothers Island tuatara	<i>Sphenodon guntheri</i>	ST	VU D1+2
<b>TERRESTRIAL INVERTEBRATES</b>			
<b>Taxonomically determinate</b>			
Bird louse	<i>Acidoproctus gottwaldbirschi</i> (Eichler, 1958)		
Snail	<i>Allodiscus fallax</i> Powell, 1952	OL	
Cook Strait click beetle	<i>Amychus granulatus</i> (Broun, 1886)	HI	
Stephens Island weevil	<i>Anagotus stephenensis</i> Kuschel, 1982	HI, OL	
Aphid	<i>Aphis bealyi</i> Cottier, 1953		
Moth	<i>Asaphodes frivola</i> (Meyrick, 1913)	DP	
Moth	<i>Asaphodes stinaria</i> (Guenée, 1868)	DP	
Bird louse	<i>Austrogoniodes struthbeus</i> Harrison, 1915		
Robust grasshopper	<i>Brachaspis robustus</i> Bigelow, 1967 s.s.	HI	
Moth	<i>Cephalissa siria</i> Meyrick, 1884	HI	
Bird louse	<i>Colpocephalum piligrimi</i> Price, 1967		
Herekopare weta	<i>Deinacrida carinata</i> Salmon, 1950	DP	
Little Barrier giant weta	<i>Deinacrida heteracantha</i> White, 1842	HI, OL	VU D2
Mahoenui giant weta	<i>Deinacrida mahoenui</i> Gibbs, 1999	CD, ST, HI	
Spider—Pisauridae	<i>Dolomedes schauinslandi</i> Simon, 1899	OL	
Moth	<i>Ericodesma aerodana</i> (Meyrick, 1881)	DP	
Snail	<i>Flammulina tepakiensis</i> Gardner, 1977	DP	
Bird louse	<i>Forficuloecus meinertzhageni</i> Guimarães, 1974		
Bird louse	<i>Forficuloecus piligrimi</i> Guimarães, 1985		
Canterbury knobbled weevil	<i>Hadramphus tuberculatus</i> (Pascoe, 1877)	DP	EX
Bird louse	<i>Heteromenopon</i> ( <i>Keamenopon</i> ) <i>kea</i> (Kellogg, 1907)		
Moth	<i>Kiwaia jeanae</i> Philpott, 1930	HI, OL	
Moth	<i>Kupea electilis</i> Philpott, 1930	OL	
Stephens Island ground beetle	<i>Mecodema costellum</i> <i>costellum</i> Broun, 1903	CD, HI, OL	
Darkling beetle	<i>Mimopeus parallelus</i> Watt, 1988	HI	
Bird louse	<i>Neopsittaconirmus kea</i> (Kellogg, 1907)		
Moth	<i>Orocrambus jansoni</i> Gaskin, 1975		
Moth	<i>Orocrambus sophistes</i> (Meyrick, 1905)	HI	
Bird louse	<i>Philopterus novaezealandiae</i> Palma & Price, 2000		
Snail	<i>Pbrixgnathus murdochii</i> Suter, 1894	DP	
King Island turrett snail	<i>Placostylus</i> ( <i>Basileostylus</i> ) <i>bollonsi arbutus</i> Powell, 1948	RC, HI, OL	VU D2 (for <i>P. bollonsi</i> s.l.)
King Island turrett snail	<i>Placostylus</i> ( <i>Basileostylus</i> ) <i>bollonsi bollonsi</i> Suter, 1908	RC, HI, OL	VU D2 (for <i>P. bollonsi</i> s.l.)

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Nationally Endangered—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Flax snail (pupuharakeke)	<i>Placostylus (Maoristylus) ambagiosus annectens</i> Powell, 1938	CD, RF	VU B1+2abcde (for <i>P. ambagiosus</i> s.l.)
Flax snail (pupuharakeke)	<i>Placostylus (Maoristylus) ambagiosus michiei</i> Powell, 1951	CD, RF, OL	VU B1+2abcde (for <i>P. ambagiosus</i> s.l.)
Flax snail (pupuharakeke)	<i>Placostylus (Maoristylus) ambagiosus paraspiritus</i> Powell, 1951	CD, HI, RF	VU B1+2abcde (for <i>P. ambagiosus</i> s.l.)
Large land snail	<i>Powelliphanta gilliesi aurea</i> (Powell, 1946)	HI, OL	DD (for <i>P. gilliesi</i> s.l.)
Large land snail	<i>Powelliphanta gilliesi gilliesi</i> (Smith, 1880)	CD, OL	DD (for <i>P. gilliesi</i> s.l.)
Large land snail	<i>Powelliphanta gilliesi jamesoni</i> (Powell, 1936)	CD, RF, OL	DD (for <i>P. gilliesi</i> s.l.)
Large land snail	<i>Powelliphanta gilliesi kaburangica</i> (Powell, 1936)	CD, OL	DD (for <i>P. gilliesi</i> s.l.)
Large land snail	<i>Powelliphanta gilliesi montana</i> (Powell, 1936)	CD, RF, OL	DD (for <i>P. gilliesi</i> s.l.)
Large land snail	<i>Powelliphanta hochstetteri anatokiensis</i> (Powell, 1938) red form	CD, RF, OL	DD (for <i>P. hochstetteri</i> s.l.)
Large land snail	<i>Powelliphanta hochstetteri anatokiensis</i> (Powell, 1938) yellow form	CD, RF	DD (for <i>P. hochstetteri</i> s.l.)
Large land snail	<i>Powelliphanta lignaria johnstoni</i> (Powell, 1946)	CD, RF	DD (for <i>P. lignaria</i> s.l.)
Large land snail	<i>Powelliphanta lignaria rotella</i> (Powell, 1938)	CD, HI, RF	DD (for <i>P. lignaria</i> s.l.)
Large land snail	<i>Powelliphanta lignaria ruforadiata</i> (Powell, 1949)	CD, OL	DD (for <i>P. lignaria</i> s.l.)
Large land snail	<i>Powelliphanta rossiana rossiana</i> (Powell, 1930)	DP	DD (for <i>P. rossiana</i> s.l.)
Large land snail	<i>Powelliphanta superba harveyi</i> (Powell, 1946)	CD, RF, OL	
Large land snail	<i>Powelliphanta superba mouatae</i> (Powell, 1946)	CD, RF, OL	
Large land snail	<i>Powelliphanta superba prouseorum</i> (Powell, 1946)	DP, RF, OL	
Large land snail	<i>Powelliphanta superba richardsoni</i> (Powell, 1946)	CD, RF, OL	
Large land snail	<i>Powelliphanta traversi florida</i> (Powell, 1946)	CD, HI	DD (for <i>P. traversi</i> s.l.)
Large land snail	<i>Powelliphanta traversi koputaroa</i> (Powell, 1946)	CD, HI	DD (for <i>P. traversi</i> s.l.)
Large land snail	<i>Powelliphanta traversi latizona</i> (Powell, 1949)	HI	DD (for <i>P. traversi</i> s.l.)
Large land snail	<i>Powelliphanta traversi tararuaensis</i> (Powell, 1938)	HI	DD (for <i>P. traversi</i> s.l.)
Large land snail	<i>Powelliphanta traversi traversi</i> (Powell, 1930)	CD, HI	DD (for <i>P. traversi</i> s.l.)
Scarab/chafer beetle	<i>Prodontria bicolorata</i> Given, 1964	HI	
Cromwell chafer beetle	<i>Prodontria lewisi</i> Broun, 1904	CD, OL	CR B1+2c
Darkling beetle	<i>Pseudobelops clandestinus</i> Watt, 1971		
Moth	<i>Schiffermuelleria orthophanes</i> (Meyrick, 1905)	OL	
Alpine grasshopper	<i>Sigaus childi</i> Jamieson, 1999		
Moth	<i>Statimopoda albimaculata</i> Philpott, 1931	DP	
Weevil	<i>Stephanorhynchus insolitus</i> Broun, 1893		
Moth	<i>Theoxena scissaria</i> (Guenée, 1868)	DP, EF?	
Bird louse	<i>Trabeculus flemingi</i> Timmermann, 1959		
Snail	<i>Wainuia clarkei</i> Powell, 1936	DP	CR B1+2ab
Moth	<i>Xanthorhoe frigida</i> Howes, 1946	DP	
<b>Taxonomically indeterminate</b>			
Bird louse	<i>Anaticola</i> sp.		
Grasshopper	<i>Brachaspis</i> “lowland”	DP	
Bird louse	<i>Brueelia</i> sp. (kokako)		
Bird louse	<i>Brueelia</i> sp. (saddleback)		
Snail	Charopidae sp. 27 (NMNZ M.58110)	DP, OL	
Snail	Charopidae sp. 165 (NMNZ M.99147)		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Nationally Endangered—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Snail	Charopidae sp. 166 (NMNZ M.79360)	DP, OL	
Snail	<i>Cyrtora</i> sp. 14 (NMNZ M.151437)	DP, HI, OL	
Moth	<i>Declana</i> sp. “grey toteuta”		
Moth	<i>Dichromodes</i> “Cloudy Bay”	DP, ST?, HI, OL	
Moth	<i>Gingidiobora subobscurata</i> (Walker, 1862) species complex “eastern Otago”	HI	
Moth	<i>Graphania</i> cf. <i>tetrachroa</i> “olearia”		
Moth	<i>Kiwatia</i> “plains jumper”	DP, HI	
Carabid	<i>Loxomerus</i> sp. “Bollons Island”	DP, OL	
Moth	<i>Maoritenes</i> sp. “olearia”		
Ground beetle	<i>Megadromus</i> “Omarama”		
Metallic green ground beetle	<i>Megadromus antarcticus</i> subsp. 1		
Moth	<i>Notoreas</i> “Cape Campbell”	HI	
Moth	<i>Notoreas</i> “northern”	HI	
Moth	<i>Orocrambus</i> “Mackenzie Basin”	OL	
Moth	<i>Pasiphila</i> sp. “olearia”		
Flax snail (pupuharakeke)	<i>Placostylus (Maoristylus) ambagiosus</i> “Ngaupoko”	HI, RF, OL	VU B1+2abcde (for <i>P. ambagiosus</i> s.l.)
Large land snail	<i>Powelliphanta</i> “Buller River”		
Large land snail	<i>Powelliphanta</i> “Gunner River”	CD, DP, RF	
Large land snail	<i>Powelliphanta</i> “Maungaharuru”	HI	
Large land snail	<i>Powelliphanta</i> “Owen”	DP	
Large land snail	<i>Powelliphanta</i> “Parapara”	CD, RF, OL	
Large land snail	<i>Powelliphanta</i> “patrickensis” (sensu Powell, 1949)	HI, OL	
Large land snail	<i>Powelliphanta gilliesi</i> “Heaphy”	CD, RF, OL	DD (for <i>P. gilliesi</i> s.l.)
Large land snail	<i>Powelliphanta superba</i> “Gouland Range”	DP, OL	
Moth	<i>Protosynaema</i> sp. “olearia”	HI	
Moth	<i>Pseudocoremia</i> sp. “knobby”	HI	
Snail	Punctidae sp. 12 (NMNZ M.87990)	OL	
Snail	Punctidae sp. 20 (NMNZ M.116650, “Microloma” “unicolorata”)	DP	
Snail	Punctidae sp. 64 (NMNZ M.68410)	DP	
<b>VASCULAR PLANTS</b>			
<b>Taxonomically determinate</b>			
	<i>Aciphylla traversii</i> (F. Muell.) Hook.f.	CD, EF, OL	
	<i>Ackama nubicola</i> de Lange	CD, RF, OL	
	<i>Amphibromus fluitans</i> Kirk	EF, TO	
	<i>Asplenium pauperequitum</i> Brownsey et P. Jackson	CD, HI, EF, OL	
	<i>Astelia chathamica</i> (Skottsb.) L.B. Moore	CD, RF, OL	
	<i>Australopyrum calcis</i> Connor et Molloy subsp. <i>calcis</i>	CD, ST, OL	
	<i>Boehmeria australis</i> var. <i>dealbata</i> (Cheeseman) Sykes	OL	
	<i>Brachyglossis buntii</i> (F. Muell.) B. Nord.	CD, RC, HI, OL	VU B1+2c
	<i>Carex inopinata</i> Cook	CD	
	<i>Carmichaelia curta</i> Petrie	HI	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Nationally Endangered—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
	<i>Carmichaelia juncea</i> Hook.f.	HI, EF, CD	
	<i>Carmichaelia kirkii</i> Hook.f.	DP	
	<i>Carmichaelia williamsii</i> Kirk	HI	
	<i>Chionthus maximus</i> Colenso		
	<i>Coprosma watima</i> A.P. Druce	CD, RC	
	<i>Crassula peduncularis</i> (Smith) F. Meigen	SO, EF	
	<i>Embergeria grandifolia</i> (Kirk) Boulos	OL, CD	
	<i>Epacris sinclairii</i> Hook.f.	RC	
	<i>Hebe armstrongii</i> (J.B. Armstr.) Cockayne et Allan	CD, HI	
	<i>Hebe salicornioides</i> (Hook.f.) Cockayne et Allan	RF	
	<i>Hebe speciosa</i> (A. Cunn.) Cockayne et Allan	CD, RF, HI	
	<i>Helichrysum dimorphum</i> Cockayne	CD, HI, RF	
	<i>Heliohebe raoulii</i> subsp. <i>maccaskillii</i> (Allan) Garn.-Jones		
	<i>Juncus holoschoenus</i> R. Br. var. <i>holoschoenus</i>	DP, SO	
	<i>Lepidium kirkii</i> Petrie	CD, HI, EF	
	<i>Lepidium oleraceum</i> Sparman s.s.	CD, HI, EF	
	<i>Lepidium sisymbrioides</i> subsp. <i>kawarau</i> (Petrie) Thell.	CD, HI	
	<i>Leptinella nana</i> (D.G. Lloyd) D.G. Lloyd et C. Webb	CD, EF	
	<i>Myosotidium hortensia</i> (Decne.) Baill.	CD, OL	
	<i>Myosotis colensoi</i> (Kirk) Macbride	CD, EF	
	<i>Myosotis petiolata</i> var. <i>pansa</i> L.B. Moore	EF	
	<i>Myosotis pygmaea</i> var. <i>glauca</i> G. Simpson et J.S. Thomson	CD, EF	
	<i>Myosurus minimus</i> subsp. <i>novae-zelandiae</i> (W.R.B. Oliv.) Garn.-Jones	HI, EF	
	<i>Myrsine argentea</i> Heenan et de Lange	OL	
	<i>Olearia crebra</i> E.K. Cameron et Heenan	CD, DP, RC	
	<i>Olearia pachyphylla</i> Cheeseman	HI	
	<i>Olearia polita</i> H.D. Wilson et Garn.-Jones	CD	EN C2b
	<i>Opbioglossum petiolatum</i> Hook.	CD, SO, HI	
	<i>Oreomyrrhis colensoi</i> var. <i>delicatula</i> Allan	CD, DP	
	<i>Phylloglossum drummondii</i> Kunze	SO, HI, EF	
	<i>Picris burbridgei</i> S. Holzapfel	SO	
	<i>Pittosporum obcordatum</i> Raoul	CD, RF	LR/nt
	<i>Pittosporum patulum</i> Hook.f.	CD, RF	EN D
	<i>Pittosporum serpentinum</i> (de Lange) de Lange	CD, RF, OL	
	<i>Pomaderris phyllicifolia</i> Lodd.	SO, HI	
	<i>Rorippa divaricata</i> (Hook.f.) Garn.-Jones et Jonsell	CD, EF	
	<i>Senecio kermadecensis</i> Belcher	EF	
	<i>Senecio scaberulus</i> (Hook.f.) D.G. Drury	HI, EF	
	<i>Simplicia laxa</i> Kirk	CD, HI	
	<i>Todea barbara</i> (L.) Moore	SO	
	<i>Triglochin palustris</i> L.	SO	
	<i>Uncinia strictissima</i> Petrie	DP, HI	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

*Nationally Endangered—continued*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
<b>Taxonomically indeterminate</b>			
	<i>Brachyscome</i> (a) (WELT 10278; Ward)		
	<i>Bulbinella modesta</i> L.B. Moore	DP	
	<i>Cardamine</i> (b) (CHR 312947; “tarn”)	DP, EF	
	<i>Cardamine</i> (d) (CHR 511706; Pisa Range)	DP	
	<i>Craspedia</i> (e) (CHR 514391; “tarn”)	CD, OL	
	<i>Gingidia</i> aff. <i>montana</i> (a) (CHR 510570; Mt Burnett)	HI	
	<i>Hebe</i> aff. <i>albicans</i> (AK 252966; Mt Burnett)	OL	
	<i>Hibiscus</i> aff. <i>trionum</i> (AK 218967; North Island)		
	<i>Lepidium</i> aff. <i>oleraceum</i> (b) (AK 208579; Antipodes)	CD, EF, DP	
	<i>Lepidium</i> aff. <i>oleraceum</i> (c) (CANU 5995; Snares)	DP	
	<i>Melicytus</i> (a) (CHR 355077; Matiri Range)		
	<i>Myosotis</i> aff. <i>pygmaea</i> (CHR 244566; Volcanic Plateau)	CD	
	<i>Pimelea</i> aff. <i>aridula</i> (b) (AK 230900; Cook Strait)	OL	
	<i>Ranunculus</i> aff. <i>stylosus</i> (CHR 515131; Manuhune)	HI, OL	
	<i>Rhopalostylis</i> aff. <i>sapida</i> (AK 227148; Chatham Islands)	CD	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

*Continued on next page*

## Nationally Vulnerable

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
<b>BATS</b>			
<b>Taxonomically indeterminate</b>			
Long-tailed bat (North Island)	<i>Chalinolobus tuberculata</i> (North Island)	DP, HI	VU A2bc (for <i>C. tuberculata</i> s.l.)
<b>BIRDS</b>			
<b>Taxonomically determinate</b>			
Wrybill, ngutu-pare	<i>Anarhynchus frontalis</i>		VU C2 a(i)
Northern New Zealand dotterel	<i>Charadrius obscurus aquilonius</i>	CD, ST	EN C1
Northern royal albatross, toroa	<i>Diomedea sanfordi</i>	RF	
Reef heron	<i>Egretta sacra sacra</i>	DP, SO	
White-flippered penguin	<i>Eudyptula minor albosignata</i>	HI	
Stewart Island shag	<i>Leucocarbo chalconotus</i>	HI	VU B1 a,c(iii)(iv) B2 a,c(iii)(iv) D2
Yellow-eyed penguin	<i>Megadyptes antipodes</i>	ST, HI	VU C1
Caspian tern	<i>Sterna caspia</i>	SO	
Pitt Island shag	<i>Stictocarbo featherstoni</i>		EN C1
Salvin's mollymawk	<i>Thalassarche salvini</i>	DP, TO	
Rock wren	<i>Xenicus gilviventris</i>	DP	VU C 1, 2a(i)
<b>Taxonomically indeterminate</b>			
Bush falcon	<i>Falco novaeseelandiae</i> "bush"	DP, HI	NT
<b>BRYOPHYTES</b>			
<b>Taxonomically determinate</b>			
Liverwort	<i>Zoopsis nitida</i>		
<b>FRESHWATER FISH</b>			
<b>Taxonomically determinate</b>			
Eldon's galaxias	<i>Galaxias eldoni</i>	CD, HI	
<b>Taxonomically indeterminate</b>			
Possible new non-diadromous galaxias	<i>Galaxias</i> "Teviot"	HI	
Dune lakes galaxias	<i>Galaxias</i> sp.	CD, HI	
<b>FRESHWATER INVERTEBRATES</b>			
<b>Taxonomically determinate</b>			
Caddisfly	<i>Kokiria milbaro</i> McFarlane, 1964		
<b>REPTILES</b>			
<b>Taxonomically determinate</b>			
Whitaker's skink	<i>Cyclodina whitakeri</i>	CD, HI	VU D1+2
<b>TERRESTRIAL INVERTEBRATES</b>			
<b>Taxonomically determinate</b>			
Snail	<i>Amborhytida tarangensis</i> (Powell, 1930)	OL	
Bird louse	<i>Austrogonioides vanalphenae</i> Banks & Palma, 2003		
Coxella weevil	<i>Hadramphus spinipennis</i> Broun, 1911	HI, EF	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Nationally Vulnerable—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Large land snail	<i>Powelliphanta fiordlandica</i> (Climo, 1971)	DP	
Large land snail	<i>Powelliphanta gilliesi compta</i> (Powell, 1930)	CD, OL	DD (for <i>P. gilliesi</i> s.l.)
Large land snail	<i>Powelliphanta lignaria lignaria</i> (Hutton, 1888)	CD, OL	DD (for <i>P. lignaria</i> s.l.)
Large land snail	<i>Powelliphanta lignaria lusca</i> (Powell, 1949)	CD, OL	DD (for <i>P. lignaria</i> s.l.)
Large land snail	<i>Powelliphanta lignaria oconnori</i> (Powell, 1938)	CD	DD (for <i>P. lignaria</i> s.l.)
Large land snail	<i>Powelliphanta lignaria unicolorata</i> (Powell, 1930)		DD (for <i>P. lignaria</i> s.l.)
Snail	<i>Rhytidida greenwoodi webbi</i> Powell, 1949		
<b>Taxonomically indeterminate</b>			
Moth	<i>Pyrgotis</i> sp. “olearia”		
<b>VASCULAR PLANTS</b>			
<b>Taxonomically determinate</b>			
	<i>Aciphylla dieffenbachii</i> Kirk	CD, EF	
	<i>Australopyrum calcis</i> subsp. <i>optatum</i>	CD	
	Connor et Molloy		
	<i>Carmichaelia carmichaeliae</i> (Hook.f.) Heenan		LR/nt
	<i>Dracophyllum longifolium</i> var. <i>septentrionale</i> W.R.B. Oliv.	RF, OL	
	<i>Dracophyllum urvilleanum</i> A. Rich.	DP	
	<i>Hebe barkeri</i> (Cockayne) Cockayne	CD, OL	VU D2
	<i>Hebe bishopiana</i> (Petrie) Hatch	CD	
	<i>Hebe cupressoides</i> (Hook.f.) Cockayne et Allan	CD, HI	
	<i>Hebe perbellia</i> de Lange		
	<i>Hebe scopulorum</i> Bayly, de Lange et Garn.Jones	CD	
	<i>Hibiscus diversifolius</i> Jacq.	SO	
	<i>Lepidium flexicaule</i> Kirk	CD, TO, EF	
	<i>Leptinella featherstonii</i> F. Muell.	CD	
	<i>Leptinella rotundata</i> (Cheeseman) D.G. Lloyd et C. Webb		
	<i>Lycopodiella serpentina</i> (Kunze) B. Øllg.	TO	
	<i>Muehlenbeckia astonii</i> Petrie	CD, RF	
	<i>Myosotis pygmaea</i> var. <i>minutiflora</i> G. Simpson et J.S. Thomson	CD, EF	
	<i>Olearia Hectorii</i> Hook.f.	CD, RF	EN C2a
	<i>Pittosporum dallii</i> Cheeseman	CD, RF	VU B1+2c
	<i>Ranunculus ternatifolius</i> Kirk		
	<i>Scutellaria novae-zelandiae</i> Hook.f.	CD	
<b>Taxonomically indeterminate</b>			
	<i>Kunzea</i> aff. <i>ericoides</i> (a) (AK 255350; Thornton)	CD, RC, OL	
	<i>Prasophyllum</i> aff. <i>patens</i> (AK 236408; New Zealand)	CD, DP	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

## Serious Decline

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
<b>BIRDS</b>			
<b>Taxonomically determinate</b>			
North Island brown kiwi	<i>Apteryx mantelli</i>	HI, RF, CD	EN A2a,e 3a,e 4a,e
Western weka	<i>Gallirallus australis australis</i>	HI, EF	NT (for <i>G. australis</i> s.l.)
Black-billed gull	<i>Larus bulleri</i>		EN A2,3,4 a,c,e
Chatham Island mollymawk	<i>Thalassarche eremita</i>	OL	CR B2 a,b(v)
<b>FRESHWATER FISH</b>			
<b>Taxonomically determinate</b>			
Dwarf inanga	<i>Galaxias gracilis</i>	CD	VU A1ce, B1+2cd, D2
<b>FUNGI</b>			
<b>Taxonomically determinate</b>			
	<i>Melampsora novaezelandiae</i>		
<b>MARINE INVERTEBRATES</b>			
<b>Taxonomically determinate</b>			
Coral	<i>Enallopsammia maranzelleri</i>	DP, S?O, HI	
Coral	<i>Madrepora oculata</i> Linnaeus, 1758	DP, S?O, HI	
Octopus	<i>Octopus kabaroa</i> O'Shea, 2000	DP	
<b>Taxonomically indeterminate</b>			
Gorgonian	<i>Iridogorgia</i> sp.	DP, SO?, HI	
Gorgonian	<i>Metallogorgia</i> cf. <i>melanotrichos</i>	DP, SO?, HI	
<b>REPTILES</b>			
<b>Taxonomically determinate</b>			
Small-scaled skink	<i>Oligosoma microlepis</i>	DP	VU B1+2bcd
<b>TERRESTRIAL INVERTEBRATES</b>			
<b>Taxonomically determinate</b>			
Bird louse	<i>Apterygon mirum</i> Clay, 1961		
Forest ringlet	<i>Dodonidia helmsii</i>	DP	
Spider—Theridiidae—black katipo spider	<i>Latrodectus atritus</i> Urquhart, 1890	HI	
Spider—Theridiidae—red katipo	<i>Latrodectus katipo</i> Powell, 1871	HI	
Ground beetle	<i>Mecodema costellum obesum</i> Townsend, 1965	DP	
Large land snail	<i>Powelliphanta annectens</i> (Powell, 1936)	CD, RF	
Large land snail	<i>Powelliphanta marchanti</i> (Powell, 1932)	CD	LR/nt
Large land snail	<i>Powelliphanta spedeni spedeni</i> (Powell, 1932)	HI	
Large land snail	<i>Powelliphanta superba superba</i> (Powell, 1930)	CD, RF	
Snail	<i>Succinea archeyi</i> Powell, 1933		
<b>Taxonomically indeterminate</b>			
Moth	<i>Notoreas</i> "Taranaki Coast"		
Moth	<i>Notoreas</i> "Wellington"		
Large land snail	<i>Powelliphanta</i> "Urewera"		
Alpine grasshopper	<i>Sigaus</i> sp. A	DP	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
<b>VASCULAR PLANTS</b>			
<b>Taxonomically determinate</b>			
	<i>Brachyglottis kirkii</i> (Kirk) C. Webb var. <i>kirkii</i>	CD, DP	
	<i>Carex litorosa</i> Bailey	DP, HI	
	<i>Dactylanthus taylorii</i> Hook.f.	CD, RF	
	<i>Daucus glochidiatus</i> (Labill.) Fisch., C.A. Mey. & Avé-Lall.	DP, SO	
	<i>Drymoanthus flavus</i> St George et Molloy		
	<i>Euphorbia glauca</i> G. Forst.	EF	
	<i>Heliohebe acuta</i> Garn.-Jones	CD, DP	
	<i>Heliohebe lavaudiana</i> (Raoul) Garn.-Jones	CD	
	<i>Hydatella inconspicua</i> (Cheeseman) Cheeseman	EF	
	<i>Isolepis basilaris</i> Hook.f.	EF, HI	
	<i>Kunzea ericoides</i> var. <i>linearis</i> (Kirk) W. Harris	HI	
	<i>Leucogenes tarabaoa</i> Molloy	OL	
	<i>Luzula celata</i> Edgar		
	<i>Marattia salicina</i> Smith	CD, SO	
	<i>Mazus novaezeelandiae</i> subsp. <i>impolitus</i> Heenan f. <i>impolitus</i>	CD, HI	
	<i>Mazus novaezeelandiae</i> W.R. Barker subsp. <i>novaezeelandiae</i>	CD, HI	
	<i>Myosotis pygmaea</i> Colenso var. <i>pygmaea</i>	DP, CD	
	<i>Olearia fimbriata</i> Heads		
	<i>Pachycladon stellata</i> (Allan) Heenan et A. Mitch.	DP	
	<i>Pimelea tomentosa</i> (J.R. Forst. et G. Forst.) Druce s.s.	EF	
	<i>Pittosporum kirkii</i> Hook.f.	CD, DP	
	<i>Plagianthus chathamicus</i> Cockayne	CD, RC	
	<i>Plumatochilus tasmanicum</i> (D.L. Jones) D.L. Szlachetko	SO, EF, HI	
	<i>Pterostylis paludosa</i> D.L. Jones, Molloy et M.A. Clem.	EF	
	<i>Tetrachondra hamiltonii</i> D. Oliver	DP	
<b>Taxonomically indeterminate</b>			
	<i>Hypericum aff. japonicum</i> (a) (CHR 165889; Volcanic Plateau)	DP	
	<i>Pimelea aff. arenaria</i> (AK 216133; southern New Zealand)	DP, RF	
	<i>Pittosporum aff. crassifolium</i> (AK 253259; Raoul Island)	RF, OL	
	<i>Sicyos aff. australis</i> (AK 252822; New Zealand)		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

## Gradual Decline

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
<b>BIRDS</b>			
<b>Taxonomically determinate</b>			
South Island rifleman	<i>Acanthisitta chloris chloris</i>	DP, HI	
North Island rifleman	<i>Acanthisitta chloris granti</i>	DP, HI	
Great spotted kiwi	<i>Apteryx haastii</i>	RF	
Southern tokoeka	<i>Apteryx australis</i>	HI, RF, DP	VU A1 c,e
Banded dotterel	<i>Charadrius bicinctus bicinctus</i>		
Yellow-crowned kakariki	<i>Cyanoramphus auriceps</i>	HI, EF	NT C1, 2b
Long-tailed cuckoo	<i>Eudynamys taitensis</i>	DP, HI	
Northern little blue penguin	<i>Eudyptula minor tredalei</i>	HI, EF	
Southern little blue penguin	<i>Eudyptula minor minor</i>	HI	
New Zealand pigeon, kereru, kukupa	<i>Hemiphaga novaeseelandiae</i>	RF	LC
Red-billed gull	<i>Larus novaehollandiae scopulinus</i>		
Antarctic prion	<i>Pachyptila desolata</i>	SO	
Light-mantled sooty albatross	<i>Phoebetria palpebrata</i>	DP, TO	
Grey petrel	<i>Procellaria cinerea</i>	TO, HI	
Cook's petrel, titi	<i>Pterodroma cookii</i>	RF	EN B2 a, b(v)
Flesh-footed shearwater	<i>Puffinus carneipes</i>	SO	
Sooty shearwater	<i>Puffinus griseus</i>	SO, HI	
New Zealand sooty tern	<i>Sterna fuscata kermadecii</i>	TO	
Southern white-fronted tern	<i>Sterna striata aucklandica</i>	HI	
White-fronted tern	<i>Sterna striata striata</i>		
<b>Taxonomically indeterminate</b>			
Eastern falcon	<i>Falco novaeseelandiae</i> "eastern"	HI, DP	NT
<b>FRESHWATER FISH</b>			
<b>Taxonomically determinate</b>			
Longfin eel	<i>Anguilla dieffenbachii</i>	HI	
Roundhead galaxias	<i>Galaxias anomalus</i>	EF, HI	
Giant kokopu	<i>Galaxias argenteus</i>	DP, HI	VU C2a
Flathead galaxias	<i>Galaxias depressiceps</i>	HI	
Dwarf galaxias	<i>Galaxias divergens</i>	DP, HI	
Gollum galaxias	<i>Galaxias gollumoides</i>	HI	
Bignose galaxias	<i>Galaxias macronasus</i>	DP	
Upland longjaw galaxias	<i>Galaxias prognathus</i>	DP	
Dusky galaxias	<i>Galaxias pullus</i>	CD, HI	
Brown mudfish	<i>Neochanna apoda</i>	HI	LR/nt
Black mudfish	<i>Neochanna diversus</i>	HI	DD
<b>Taxonomically indeterminate</b>			
Possible new non-diadromous galaxias	<i>Galaxias</i> "Southern sp."	HI	
Possible new non-diadromous galaxias	<i>Galaxias</i> sp. D	HI	
<b>FRESHWATER INVERTEBRATES</b>			
<b>Taxonomically determinate</b>			
Freshwater mussel	<i>Hyridella menziesi</i> (Gray, 1843)		
Koura	<i>Paranephrops planifrons</i> White		
Koura	<i>Paranephrops zealandicus</i> White		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
<b>FUNGI</b>			
<b>Taxonomically determinate</b>			
	<i>Propolis desmoschoenii</i>	CD, EF	
<b>Taxonomically indeterminate</b>			
	<i>Diaporthe</i> sp. 1	CD, HI	
	<i>Diaporthe</i> sp. 2	CD, HI	
	<i>Glonium</i> sp.	CD, EF	
	<i>Leucostoma</i> sp. 1	CD, HI	
	<i>Leucostoma</i> sp. 2	CD, HI	
	<i>Pestalotiopsis</i> sp.	CD, EF	
	<i>Phomopsis</i> sp.	CD, OL	
	<i>Seimatosprium</i> sp.	CD, EF	
	<i>Truncatella</i> sp.	CD, EF	
<b>MARINE FISH</b>			
<b>Taxonomically determinate</b>			
Great white shark/white pointer	<i>Carcharodon carcharias</i>	TO	
Basking shark	<i>Cetorhinus maximus</i>	TO	
<b>MARINE INVERTEBRATES</b>			
<b>Taxonomically determinate</b>			
Volute	<i>Alcithoe benthicola</i>	DP	
Volute	<i>Alcithoe fissurata</i>	DP	
Volute	<i>Alcithoe flemingi</i>	DP	
Volute	<i>Alcithoe larocheti</i>	DP	
Volute	<i>Alcithoe lutea</i>	DP	
Giant deep mussel	<i>Bathymodiolus tangaroa</i> Cosel & Marshall, 2003		
Snail	<i>Calliostoma turnerarum</i>		
Cyclostome bryozoan	<i>Calvetia osbeai</i> Taylor & Gordon, 2003		
Hermit crab	<i>Cancellus laticoxa</i> Forest & McLaughlan, 2000	OL	
Limpet	<i>Cellana flava</i> (Hutton, 1873)		
Basket starfish	<i>Gorgonocephalus dolichodactylus</i> Döderlein, 1911		
Golden volute	<i>Provocator mirabilis</i>		
Cyclostome bryozoan	<i>Spiritonopora perplexa</i> Taylor & Gordon, 2003	DP	
Bryozoan	<i>Steginoporella perplexa</i>		
<b>REPTILES</b>			
<b>Taxonomically determinate</b>			
Ornate skink	<i>Cyclodina ornata</i>	DP	
Goldstripe gecko	<i>Hoplodactylus chrysostrelicus</i>	HI	
Pacific gecko	<i>Hoplodactylus pacificus</i>	HI	
Harlequin gecko	<i>Hoplodactylus rakturiae</i>	DP	LR/nt
Auckland green gecko	<i>Naultinus elegans elegans</i>	HI	
Wellington green gecko	<i>Naultinus elegans punctatus</i>	HI	
Jewelled gecko	<i>Naultinus gemmeus</i>	HI	
Northland green gecko	<i>Naultinus grayii</i>	HI	
Rough gecko	<i>Naultinus rufus</i>	DP, HI	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

*Gradual Decline—continued*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Nelson green gecko	<i>Naultinus stellatus</i>	DP, HI	
Green skink	<i>Oligosoma chloronoton</i>	HI	
Cryptic skink	<i>Oligosoma inconspicuum</i>	DP, HI	
Speckled skink	<i>Oligosoma infrapunctatum</i>	HI	LR/nt
Spotted skink	<i>Oligosoma lineoocellatum</i>	HI	
Scree skink	<i>Oligosoma waimatense</i>	HI	VU B1+2bd
<b>Taxonomically indeterminate</b>			
Canterbury gecko	<i>Hoplodactylus</i> “Canterbury”	HI	
Matapia gecko	<i>Hoplodactylus</i> “Matapia Island”	HI	
Large Otago gecko	<i>Hoplodactylus</i> “Otago large”		
Southern forest gecko	<i>Hoplodactylus</i> “Southern forest gecko”	DP, HI	
Green skink “West Otago”	<i>Oligosoma</i> aff. <i>chloronoton</i> “West Otago”	DP	
Spotted skink “Mackenzie Basin”	<i>Oligosoma</i> aff. <i>lineoocellatum</i> “Mackenzie Basin”	DP	
Spotted skink “South Marlborough”	<i>Oligosoma</i> aff. <i>lineoocellatum</i> “South Marlborough”		
<b>TERRESTRIAL INVERTEBRATES</b>			
<b>Taxonomically determinate</b>			
Snail	<i>Amborhytida dunniae</i> (Gray, 1840)		
Snail	<i>Amborhytida duplicita</i> (Suter, 1904)		
Snail	<i>Amborhytida forsythi</i> (Powell, 1952)		
Bird louse	<i>Apterygon dumosum</i> Tandan, 1972		
Bird louse	<i>Apterygon hintoni</i> Clay, 1966		
Kaikoura giant weta	<i>Deinacrida parva</i> Buller, 1895	HI, EF	DD
Moth	<i>Gingidiobora nebulosa</i> (Philpott, 1917)		
Sphagnum porina	<i>Heloxycanus patricki</i> Dugdale, 1994		
Moth	<i>Hydriomena canescens</i> Philpott, 1918		
Moth	<i>Hydriomena clarkei</i> (Howes, 1917)		
Ground beetle	<i>Mecodema bowitti</i> Castelnau, 1867		
Moth	<i>Meterana exquisita</i> (Philpott, 1903)	HI	
Moth	<i>Meterana grandiosa</i> (Philpott, 1903)	HI	
Moth	<i>Paranotoreas fulva</i> (Hudson, 1905)		
Kauri snail, pupurangi	<i>Paryphanta busbyi busbyi</i> (Gray, 1840)		LR/nt (for <i>P. busbyi</i> s.l.)
Kauri snail, pupurangi	<i>Paryphanta busbyi wattii</i> Powell, 1946	CD, HI	LR/nt (for <i>P. busbyi</i> s.l.)
Large land snail	<i>Powelliphanta gilliesi fallax</i> (Powell, 1930)		DD (for <i>P. gilliesi</i> s.l.)
Large land snail	<i>Powelliphanta gilliesi subfusca</i> (Powell, 1930)	HI	DD (for <i>P. gilliesi</i> s.l.)
Large land snail	<i>Powelliphanta hochstetteri bicolor</i> (Powell, 1930)	CD, RF	DD (for <i>P. hochstetteri</i> s.l.)
Large land snail	<i>Powelliphanta hochstetteri consobrina</i> (Powell, 1936)	RF	DD (for <i>P. hochstetteri</i> s.l.)
Large land snail	<i>Powelliphanta hochstetteri hochstetteri</i> (Pfeiffer, 1862) brown based	CD	DD (for <i>P. hochstetteri</i> s.l.)
Large land snail	<i>Powelliphanta hochstetteri hochstetteri</i> (Pfeiffer, 1862) yellow based	CD	DD (for <i>P. hochstetteri</i> s.l.)
Large land snail	<i>Powelliphanta hochstetteri obscura</i> (Beutler, 1901)	CD, RF	DD (for <i>P. hochstetteri</i> s.l.)
Moth	“ <i>Pseudocoremia</i> ” <i>cineracia</i> (Howes, 1942)	HI	
Bird louse	<i>Rallicola</i> ( <i>Apteriscola</i> ) <i>gadowi</i> Harrison, 1915		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

*Continued on next page*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Bird louse	<i>Rallicola (Aptericola) gracilentus</i> Clay, 1953		
Snail	<i>Rhytidia stephenensis</i> Powell, 1930		
Alpine grasshopper	<i>Sigaus minutus</i> Bigelow, 1967		
Snail	<i>Wainutia edwardi</i> (Suter, 1899)		
Snail	<i>Wainutia urnula nasuta</i> Powell, 1946		
<b>Taxonomically indeterminate</b>			
Snail	<i>Ambarhytida</i> aff. <i>forsythi</i> (NMNZ M.173834)		
Slug	<i>Athoracophorus</i> sp. 4 (NMNZ M.151430) “northern NZ”	HI	
Moth	<i>Gingidiobora subobscurata</i> (Walker, 1862) species complex	DP	
Karikari tree weta	<i>Hemideina thoracica</i> 2n=23,24		
Moth	<i>Loxostege</i> sp. “salt pan”		
Large land snail	<i>Powelliphanta</i> “Haast”		DP
Moth	<i>Stathmopoda</i> sp. “olearia”		
Snail	<i>Wainutia</i> “Mount Oxford”		
<b>VASCULAR PLANTS</b>			
<b>Taxonomically determinate</b>			
Pingao	<i>Acaena buchananii</i> Hook.f.	DP	
	<i>Alepis flavidia</i> (Hook.f.) Tiegh.	CD, HI	
	<i>Anogramma leptophylla</i> (L.) Link	SO, EF	
	<i>Austrostipa littoralis</i> (Labill.) E.B. Alexev.	CD, SO, HI	
	<i>Brachyglottis perdicoides</i> (Hook.f.) B. Nord.	CD	LR/nt
	<i>Brachyglottis sciadophila</i> (Raoul) B. Nord.		
	<i>Carex astonii</i> Hamlin	DP	
	<i>Carex cirrhosa</i> Bergg.	DP, HI	
	<i>Carmichaelia crassicaule</i> Hook.f.	RF	
	<i>Carmichaelia stevensonii</i> (Cheeseman) Heenan		VU D2
	<i>Carmichaelia vexillata</i> Heenan	RF	
	<i>Celmisia major</i> Cheeseman var. <i>major</i>	HI	
	<i>Clematis marmoraria</i> Sneddon	CD, RF	
	<i>Colensoa physaloides</i> (A. Cunn.) Hook.f.		
	<i>Coprosma obconica</i> Kirk	CD, RF	
	<i>Coprosma pedicellata</i> Molloy, de Lange et B.D. Clarkson	CD	
	<i>Coprosma wallii</i> Petrie	CD, RF	VU B1+2c, D2
	<i>Crassula kirkii</i> (Allan) A.P. Druce et D.R. Given	DP	
	<i>Crassula manaia</i> A.P. Druce et Sykes	CD	
	<i>Cyclotoma interrupta</i> (Willd.) H. Itô	SO	
	<i>Deschampsia cespitosa</i> (L.) Beauv.	SO, HI	
	<i>Desmoschoenus spiralis</i> (A. Rich.) Hook.f.	CD, EF	
	<i>Doodia squarrosa</i> Colenso		
	<i>Drosera pygmaea</i> DC.	SO	
	<i>Eleocharis neozelandica</i> Kirk	EF	
	<i>Epilobium chionanthum</i> Hauss.	DP	
	<i>Eryngium</i> aff. <i>vesiculosum</i> (AK 232583; New Zealand)		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

*Gradual Decline—continued*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
	<i>Gratiola nana</i> Benth.	SO, HI	
	<i>Gunnera arenaria</i> Cheeseman		
	<i>Hebe pimeleoides</i> subsp. <i>faucicola</i> Kellow et Bayly		
	<i>Iphigenia novae-zelandiae</i> (Hook.f.) Baker	DP, HI	
	<i>Isolepis fluitans</i> (L.) R. Br.	SO, HI	
	<i>Jovellana sinclairii</i> (Hook.f.) Kranzl.	DP	
	<i>Lepidium sisymbrioides</i> Hook.f. subsp. <i>sisymbrioides</i>	CD	
	<i>Lepidium tenuicaule</i> Kirk	HI	
	<i>Leptinella serrulata</i> (D.G. Lloyd) D.G. Lloyd et C. Webb		
	<i>Libertia peregrinans</i> Cockayne et Allan	HI, CD	
	<i>Mazus arenarius</i> Heenan, P.N. Johnson et C. Webb		
	<i>Melicytus flexuosus</i> Molloy et A.P. Druce	RF	LR/nt
	<i>Mida salicifolia</i> A. Cunn.	RF	
	<i>Montigena novae-zelandiae</i> (Hook.f.) Heenan	DP, RF	
	<i>Myosotis brockiei</i> L.B. Moore et M.J.A. Simpson	CD	
	<i>Myriophyllum robustum</i> Hook.f.	CD	
	<i>Olearia cheesemani</i> Cockayne et Allan	DP	
	<i>Ourisia modesta</i> Diels	DP	
	<i>Pachycladon cheesemani</i> Heenan et A. Mitch.	DP	
	<i>Pachycladon enysii</i> (Cheeseman) Heenan et A. Mitch.		
	<i>Pachycladon fastigiata</i> (Hook.f.) Heenan et A. Mitch.		
	<i>Paspalum orbiculare</i> G. Forst.	SO, HI	
	<i>Pellaea falcata</i> (R. Br.) Fée	SO, HI	
	<i>Peraxilla colensoi</i> (Hook.f.) Tiegh.	CD, HI	
	<i>Peraxilla tetrapetala</i> Tiegh.	CD, HI	
	<i>Pimelea arenaria</i> A. Cunn. s.s.	HI, RF	
	<i>Pimelea lyallii</i> Hook.f.		
	<i>Potamogeton pectinatus</i> L.	DP	
	<i>Pseudopanax laetus</i> (Kirk) Philipson	RF	
	<i>Ranunculus limosella</i> Kirk	HI	
	<i>Ranunculus macropus</i> Hook.f.	HI	
	<i>Ranunculus recens</i> Kirk var. <i>recens</i>	CD	
	<i>Raoulia monroi</i> Hook.f.	EF	
	<i>Raukaua edgerleyi</i> (Hook.f.) Seem.	RF, CD	
	<i>Schoenus carsei</i> Cheeseman	SO, HI	
	<i>Selliera rotundifolia</i> Heenan	CD	
	<i>Sonchus kirkii</i> Hamlin	HI, EF	
	<i>Sophora fulvida</i> (Allan) Heenan et de Lange		
	<i>Teucrium parvifolium</i> Hook.f.	CD	
	<i>Thelypteris confluens</i> (Thunb.) C. Morton	CD, SO	
	<i>Trisetum antarcticum</i> (G. Forst.) Trin.	HI	
	<i>Tupeia antarctica</i> (G. Forst.) Cham. et Schlecht	CD, HI	
	<i>Urtica linearifolia</i> (Hook.f.) Cockayne	CD	
	<i>Utricularia australis</i> R. Br.	HI	
	<i>Utricularia delicatula</i> Cheeseman	HI	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

*Continued on next page*

*Gradual Decline—continued*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
<b>Taxonomically indeterminate</b>			
	<i>Christella aff. dentata</i> (b) (AK 126902; “thermal”)	HI	
	<i>Craspedia</i> (n) (CHR 369978; Henderson)	OL	
	<i>Hoberia</i> aff. <i>sexstylosa</i> (AK 234306; Tararua Ranges)	HI	
	<i>Leptinella ditoica</i> subsp. <i>monoica</i> (AK 200874)		
	<i>Melicytus</i> aff. <i>alpinus</i> (f) (CHR 541566; Waipapa)		
	<i>Ranunculus</i> (b) (CHR 324466; Burgoo Stream)		
	<i>Raoulia</i> aff. <i>bookeri</i> (AK 239529; “coast”)		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

*Continued on next page*

## Sparse

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
<b>BIRDS</b>			
<b>Taxonomically determinate</b>			
South Island fernbird, matata	<i>Bowdleria punctata punctata</i>	HI	NT (for <i>B. punctata</i> s.l.)
North Island fernbird, matata	<i>Bowdleria punctata vealeae</i>	HI	NT (for <i>B. punctata</i> s.l.)
Brown skua	<i>Catharacta antarctica lonnbergi</i>	SO	
Banded rail	<i>Gallirallus philippensis assimilis</i>	DP, HI	
Black shag	<i>Phalacrocorax carbo novaehollandiae</i>	SO	
New Zealand dabchick, wewelia	<i>Poliocephalus rufopectus</i>	ST?	VU C 2a(i)
Marsh crake	<i>Porzana pusilla affinis</i>		
Spotless crake	<i>Porzana tabuensis plumbea</i>	SO	
<b>BRYOPHYTES</b>			
<b>Taxonomically determinate</b>			
Liverwort	<i>Acromastigum cunninghamii</i>	S?O	
Liverwort	<i>Acromastigum marginatum</i>		
Liverwort	<i>Archeophylla schusteri</i>		
Liverwort	<i>Austrolejeunea hispida</i>		
Liverwort	<i>Austroscyphus phoenichorbizus</i>	SO	
Liverwort	<i>Bazzania nova</i>		
Moss	<i>Bryum pallescens</i> Schleich. ex Schwägr.	SO	
Moss	<i>Buxbaumia novae-zelandiae</i> Dixon	SO?	
Liverwort	<i>Cephalozia badia</i>	SO	
Liverwort	<i>Chloranthelia berggrenii</i>		
Moss	<i>Cratoneuron filicinum</i> (Hedw.) Spruce	SO	
Liverwort	<i>Drucella integriflora</i>		
Moss	<i>Encalypta rhaptocarpa</i> Schwaegr.	DP	
Liverwort	<i>Haplomitrium ovalifolium</i>		
Moss	<i>Hylocomitium splendens</i> (Hedw.) B.S.G.	DP	
Liverwort	<i>Isolembidium anomalum</i> var. <i>cucullatum</i>	SO	
Liverwort	<i>Isophyllum attenuata</i>	S?O	
Liverwort	<i>Kurzia tenax</i>	S?O	
Liverwort	<i>Lepidozia acantha</i>	DP	
Liverwort	<i>Metzgeria bartlettii</i>		
Liverwort	<i>Mnioloma fusca</i>	SO	
Liverwort	<i>Pachyschistochila latiloba</i>		
Liverwort	<i>Papillolejeunea pocsii</i>		
Liverwort	<i>Pedinophyllum monoicum</i>	DP	
Liverwort	<i>Porella pulcherrima</i>		
Liverwort	<i>Pseudocephalozia lepidozoides</i>		
Liverwort	<i>Radula ratkowskiana</i>	SO	
Liverwort	<i>Rectolejeunea ocellata</i>		
Liverwort	<i>Riccardia multicorpora</i>		
Moss	<i>Zygodon rufescens</i> (Hampe) Broth. in Paris	SO	
<b>Taxonomically indeterminate</b>			
Moss	<i>Calomnion brownseyi</i> Vitt et H.A. Miller		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Liverwort	<i>Goebelobryum</i> aff. <i>unguiculatum</i>	HI	
Liverwort	<i>Rectolejeunea</i> sp.		
<b>FRESHWATER FISH</b>			
<b>Taxonomically determinate</b>			
Shortjaw kokopu	<i>Galaxias postvectis</i>	HI	VU B1+2bcd
Lamprey	<i>Geotria australis</i>	DP, SO	
<b>FRESHWATER INVERTEBRATES</b>			
<b>Taxonomically determinate</b>			
Freshwater crab	<i>Amarinus lacustris</i>	SO	
Freshwater isopod	<i>Austridotea annectens</i> Nicholls, 1938		
Amphipod	<i>Chaetocorophium lucasi</i> (Hurley, 1954)		
Caddis	<i>Ecnomina zealandica</i> Wise, 1958	DP	
Tadpole shrimps	<i>Lepidurus viridus</i>		
Freshwater polychaete	<i>Namanereis tiriteae</i> (Winterbourn, 1969)	SO	
Caddisfly	<i>Olinga fumosa</i> Wise, 1959		
Flatworm	<i>Prorbynchus baswelli</i> Steinboeck & Reisinger, 1924	DP	
Flatworm	<i>Prorbynchus putealis</i> Haswell, 1898	DP	
Mayfly	<i>Tepakia caligata</i> Towns & Peters, 1996		
Caddisfly	<i>Paroxyethira kimminsi</i> Leader, 1972		
<b>FROGS</b>			
<b>Taxonomically determinate</b>			
Hochstetter's frog	<i>Leiopelma hochstetteri</i>	HI	VU A3e; B1ab(iii)
<b>FUNGI</b>			
<b>Taxonomically determinate</b>			
	<i>Asterinella intensa</i>		
	<i>Lophodermium kaikawakae</i>		
<b>Taxonomically indeterminate</b>			
	<i>Mycosphaerella</i> sp.		
	<i>Patellaria</i> sp.		
	<i>Phyllosticta</i> sp.	EF	
<b>MARINE FISH</b>			
<b>Taxonomically determinate</b>			
Brotula/mottled brotula	<i>Bidenichthys consobrinus</i>		
Frill shark	<i>Chlamydoselachus anguineus</i>	DP, SO	
Mandarin dogfish	<i>Cirrhigaleus barbifer</i>	DP, TO	
Bramble shark	<i>Echinorhinus brucus</i>	DP, SO	
Prickly shark	<i>Echinorhinus cookei</i>	DP, SO	
Lancelet	<i>Epigonichthys hectori</i>		
Smooth lantern shark	<i>Etomopterus pusillus</i>	DP, SO	
Round batfish	<i>Halieutaea maoria</i>	DP	
Sharpnose sevengill shark	<i>Heptranchias perlo</i>	DP, SO	
Sixgill shark	<i>Hexanchus griseus</i>	DP, SO	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

*Sparse—continued*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Goblin shark	<i>Mitsukurina owstoni</i>	DP, SO	
Smalltooth sand tiger	<i>Odontaspis ferox</i>	TO	
Yellow boarfish	<i>Pentaceros decacanthus</i>	SO	
Whitetail dogfish	<i>Scymnodalatias albicauda</i>	DP, SO	
Sherwood's dogfish	<i>Scymnodalatias sherwoodi</i>		
Pacific sleeper shark	<i>Somniosus antarcticus</i>	DP, SO	
Velvet dogfish	<i>Zameus squamulosus</i>	DP, SO	
<b>Taxonomically indeterminate</b>			
Pipefish	<i>Stigmatopora ?argus</i>	SO?	
<b>MARINE INVERTEBRATES</b>			
<b>Taxonomically determinate</b>			
Crab	<i>Elamena momona</i>		
Red coral	<i>Errina novazealandica</i>		
Pin-cushion star	<i>Eurygonias hyalacanthus</i>		
Crab	<i>Halimena aotearoa</i>		
<b>REPTILES</b>			
<b>Taxonomically determinate</b>			
Duvaucel's gecko	<i>Hoplodactylus duvaucelii</i>	HI	
Black-eyed gecko	<i>Hoplodactylus kabutarae</i>		LR/nt
Cloudy gecko	<i>Hoplodactylus nebulosus</i>	HI	
Marlborough green gecko	<i>Naultinus manukanus</i>	DP, HI	
West Coast green gecko	<i>Naultinus tuberculatus</i>	DP	
Long-toed skink	<i>Oligosoma longipes</i>	DP	
Moko skink	<i>Oligosoma moco</i>	HI	
Southern skink	<i>Oligosoma notosaurus</i>	DP	
Brown skink	<i>Oligosoma zelandicum</i>		
Northern tuatara	<i>Sphenodon punctatus punctatus</i>	ST, HI	
<b>Taxonomically indeterminate</b>			
North Cape Pacific gecko	<i>Hoplodactylus</i> “North Cape Pacific gecko”	HI	
North Cape green gecko	<i>Naultinus</i> “North Cape green gecko”	DP	
<b>TERRESTRIAL INVERTEBRATES</b>			
<b>Taxonomically determinate</b>			
Moth	<i>Acroclita discariana</i> Philpott, 1930		
Spider—Amphinectidae	<i>Amphinecta miltna</i> Forster & Wilton, 1973		
Northland tusked weta	<i>Anisoura nicobarica</i> Ander, 1938	DP	
Spider—Lycosidae	<i>Anoteropsis alpina</i> Vink, 2002		
Spider—Lycosidae	<i>Anoteropsis arescens</i> (Goyen, 1887)		
Spider—Lycosidae	<i>Anoteropsis blesti</i> Vink, 2002		
Spider—Lycosidae	<i>Anoteropsis litoralis</i> Vink, 2002		
Spider—Lycosidae	<i>Anoteropsis montana</i> Vink, 2002		
Spider—Lycosidae	<i>Anoteropsis westlandica</i> Vink, 2002		
Large green small-headed bladder fly	<i>Apsona muscaria</i>	DP	
Spider—Lycosidae	<i>Artoria hospita</i> Vink, 2002		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

*Continued on next page*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Carabid	<i>Brullea antarctica</i> (Castelnau, 1867)		
Spider—Linyphiidae	<i>Cassafroneta forsteri</i> Blest, 1979		
Beetle	<i>Cathartocryptus maculosus</i> (Broun)	DP	
Darkling beetle	<i>Chrysopeplus expolitus</i> (Broun, 1880)		
Moth	<i>Circoxena ditrocha</i> Meyrick, 1916		
Moth	<i>Ctenarchis crambooides</i> Dugdale, 1995		
Spider—Cycloctenidae	<i>Cycloctenus centralis</i> Forster, 1979		
Moth	<i>Dasyuris enysti</i> (Butler, 1877)		
Bluff weta	<i>Deinacrida elegans</i> Gibbs, 1999		
Mt Arthur giant weta	<i>Deinacrida tibiospina</i> Salmon, 1950		
Snail	<i>Delouagapia cordelia</i> (Hutton, 1883)		
Assassin bug	<i>Empicoris seorsus</i> (Bergrøth, 1927)	DP	
Moth	<i>Ericodesma cuneata</i> (Clarke, 1926)	ST, HI	
Spider—Araneidae	<i>Eriophora decorosa</i> (Urquhart, 1894)	SO	
Spider—Araneidae	<i>Eriophora heroine</i> (L. Koch, 1871)	SO	
Moth	<i>Gadira petraula</i> (Meyrick, 1883)		
Te Aroha stag beetle	<i>Geodorus auriculatus</i> (Broun, 1903)	DP	
Moth	<i>Glyphipterix euastera</i> Meyrick, 1880	DP, HI	
Cranefly	<i>Gynoplistia troglodyta</i> Alexander, 1962	DP	
Spider—Linyphiidae	<i>Haplitis attenuata</i> Blest & Vink, 2002		
Spider—Linyphiidae	<i>Haplitis fucatnia</i> (Urquhart, 1894)		
Spider—Linyphiidae	<i>Haplitis major</i> (Blest, 1979)		
Spider—Linyphiidae	<i>Haplitis rupicola</i> (Blest, 1979)		
Spider—Linyphiidae	<i>Haplitis subtilis</i> Blest & Vink, 2002		
Hawkes Bay tree weta	<i>Hemideina trewicki</i> Morgan-Richards, 1995		
Ground beetle	<i>Holcaspis falcis</i> Butcher, 1984	DP	
Moth	<i>Isonomeutis restincta</i> Meyrick, 1923		
Moth	<i>Kiwaia pumila</i> (Philpott, 1928)	HI	
Spider—Linyphiidae	<i>Laetesia distincta</i> Millidge, 1988		
Weevil	<i>Lyperobius carinatus</i> Broun, 1881	DP, EF	
Bug	<i>Malpba cockrofti</i> Myers, 1924	DP	
Bug	<i>Malpba mutri</i> Myers, 1924	DP	
Moth	<i>Maoricrambus oncobilus</i> (Meyrick, 1885)		
Ground beetle	<i>Mecodema atrox</i> Britton, 1949	DP	
Giant ground beetle	<i>Mecodema chiltoni</i> Broun, 1917		
Ground beetle	<i>Mecodema costellatum lewisi</i> Broun, 1908	DP, HI	
Ground beetle	<i>Mecodema nitidum</i> Broun, 1903	DP	
Ground beetle	<i>Mecodema proximus</i> Britton, 1949	DP	
Ground beetle	<i>Mecodema pulchellum</i> Townsend, 1965	DP	
Ground beetle	<i>Mecodema rex</i> Britton, 1949	DP	
Ground beetle	<i>Megadromus compressus</i> (Sharp, 1886)	DP	
Ground beetle	<i>Megadromus fultoni</i> (Broun, 1882)		
Ground beetle	<i>Megadromus haplopus</i> (Broun, 1893)		
Darkling beetle	<i>Menimus crinalis</i> Broun, 1880		
Darkling beetle	<i>Menimus curtulus</i> Broun, 1883		
Darkling beetle	<i>Menimus laevicollis</i> Broun, 1895		
Darkling beetle	<i>Menimus oblongus</i> Broun, 1880		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Spider—Linyphiidae	<i>Metamynoglenes flagellata</i> Blest, 1979		
Moth	<i>Meterana pansicolor</i> (Howes, 1912)		
Northern pimelea cutworm moth	<i>Meterana pictula</i> (White in Taylor, 1855)	HI	
Cranefly	<i>Mischoderus neptunus</i> (Edwards, 1923)	DP	
Moth	<i>Mnesarchaea fallax</i> Philpott, 1927		
Longhorn beetle	<i>Nesoptychia simpliceps</i> (Broun, 1880)		
Spider—Linyphiidae	<i>Novafroneta annulipes</i> Blest, 1979		
Spider—Linyphiidae	<i>Novafroneta gladiatrix</i> Blest, 1979		
Ground beetle	<i>Oregus crypticus</i> Pawson, 2003		
Ground beetle	<i>Oregus septentrionalis</i> Pawson, 2003		
Snail	<i>Paracharopa rimu</i> Climo, 1985	DP	
Spider—Linyphiidae	<i>Parafroneta demota</i> Blest & Vink, 2002		
Spider—Linyphiidae	<i>Parafroneta insula</i> Blest, 1979		
Stag beetle	<i>Paralissotess mangonuiensis</i> (Brookes, 1927)		
Stag beetle	<i>Paralissotess oconnori</i>	DP	
Spider	<i>Periegops suterti</i> Simon		
Scarab/chafer beetle	<i>Prodontria grandis</i> Given, 1964	DP	
Scarab/chafer beetle	<i>Prodontria matagouriae</i> Emerson, 1997		
Scarab/chafer beetle	<i>Prodontria minuta</i> Emerson, 1997		
Spider—Linyphiidae	<i>Promynoglenes grandis</i> Blest, 1979		
Spider—Linyphiidae	<i>Promynoglenes nobilis</i> Blest, 1979		
Spider—Linyphiidae	<i>Promynoglenes silvestris</i> Blest, 1979		
Spider—Linyphiidae	<i>Pseudafroneta incerta</i> (Bryant, 1935)		
Spider—Linyphiidae	<i>Pseudafroneta pallida</i> Blest, 1979		
Spider—Linyphiidae	<i>Pseudafroneta perplexa</i> Blest, 1979		
Moth	<i>Pseudocoremia fluminea</i> (Philpott, 1926)		
Moth	<i>Samana acutata</i> Butler, 1877		
Paua slug	<i>Schizoglossa worthyae</i> Powell, 1949		
Bug	<i>Semo barrisi</i> (Myers, 1924)	DP	
Moth	<i>Statimopoda aristodoxa</i> Meyrick, 1926		
Beetle	<i>Tangarona pensus</i> (Broun, 1880)	DP	
Moth	<i>Thambotricha vates</i> Meyrick, 1922	DP	
Moth	<i>Tmetolophota blenheimensis</i> (Fereday, 1883)	DP	
Moth	<i>Trachypepla cyphonias</i> Meyrick, 1927		
Weevil	<i>Tychanopais tuberosus</i> (Broun, 1923)	DP	
Moth	<i>Xanthorhoe loppogramma</i> Meyrick, 1897	DP	
Darkling beetle	<i>Zeadelium aeratum</i> (Broun, 1880)		
Darkling beetle	<i>Zeadelium chalmeri</i> (Broun, 1883)		
<b>Taxonomically indeterminate</b>			
Aphid	<i>Aphis</i> sp. 1		
Snail	<i>Charopidae</i> sp. 33 (NMNZ M.85221)		
Snail	<i>Charopidae</i> sp. 35 (NMNZ M.103006)		
Snail	<i>Charopidae</i> sp. 51 (NMNZ M.86353)		
Aphid	<i>Euschizapbis</i> sp.	DP	
Snail	<i>Flammulina</i> “Akatarawa” (NMNZ M.126198)		
Snail	“Haurakora” “aff. corrugata” (NMNZ M.32092)		
Snail	<i>Iotula</i> “aff. macroaccelerata” (NMNZ M.102700)		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Snail	“Poripotaka” “ponderi” (NMNZ M.88157)		
Snail	“Powellaoma” “arandi” (NMNZ M.151441)		
Snail	<i>Rotadiscus</i> “aff. urquharti Southland group” (NMNZ M.151456)		
Snail	<i>Sinployea</i> “fiordlandica” (NMNZ M.81649)		
Snail	<i>Sinployea</i> “Hikurangi” (NMNZ M.76803)		
Moth	<i>Stigmella</i> sp. “traversia”	DP	
Beetle	<i>Syrphetodes</i> sp. “Te Paki”	DP	
Beetle	<i>Syrphetodes</i> sp. “Waipoua”	DP	
Snail	<i>Therasiella</i> “aff. elevata” (NMNZ M.96613)		
Snail	“Vicariodiscus” “charopiformis” (NMNZ M.29033)		
Snail	<i>Watnuta</i> “Fiordland” (NMNZ M.32755)	DP	
<b>VASCULAR PLANTS</b>			
<b>Taxonomically determinate</b>			
	<i>Aciphylla multisecta</i> Cheeseman		
	<i>Aciphylla subflabellata</i> W.R.B. Oliv.	DP	
	<i>Adelopetalum tuberculatum</i> (Colenso) D.L. Jones, M.A. Clem. et Molloy		
	<i>Agrostis imbecilla</i> Zотов		
	<i>Agrostis oresbia</i> Edgar		
	<i>Anemanthele lessoniana</i> (Steud.) Veldkamp	DP	
	<i>Anemone tenuicaulis</i> (Cheeseman) Parkin et Sledge		
	<i>Anzybas rotundifolius</i> (Hook.f.) D.L. Jones et M.A. Clem.		
	<i>Asplenium cimmeriorum</i> Brownsey et de Lange		
	<i>Asplenium scleroprium</i> Hombr.		
	<i>Atriplex buchananii</i> (Kirk) Cheeseman		
	<i>Blechnum norfolkianum</i> (Heward) C. Chr.	TO	
	<i>Botrychium australe</i> R. Br.	DP, SO, EF	
	<i>Calochilus paludosus</i> R. Br.	SO, EF	
	<i>Calochilus robertsonii</i> Benth.	SO, EF	
	<i>Calystegia marginata</i> R. Br.	SO, EF	
	<i>Carex berggrenii</i> Petrie		
	<i>Carex capillacea</i> Boott		
	<i>Carex chathamica</i> C.B. Clarke	OL	
	<i>Carex edgariae</i> Hamlin	DP	
	<i>Carex enysii</i> Petrie	DP	
	<i>Carex lachenalii</i> subsp. <i>parkeri</i> Petrie	DP	
	<i>Carex muelleri</i> Petrie		
	<i>Carex tenuiculmis</i> (Petrie) Heenan et de Lange		
	<i>Celmisia hookeri</i> Cockayne		
	<i>Centrolepis strigosa</i> (R. Br.) Roem. et Schult.	SO, EF	
	<i>Chionochloa vireta</i> H.E. Connor	DP	
	<i>Clematis petriei</i> Allan		
	<i>Convolvulus fracto-saxosa</i> Petrie		
	<i>Convolvulus verecundus</i> Allan		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
	<i>Coprosma intertexta</i> G. Simpson		
	<i>Corunastylis nuda</i> (Hook.f.) D.L. Jones et M.A. Clem.	SO, EF	
	<i>Corunastylis pumila</i> (Hook.f.) D.L. Jones et M.A. Clem.	SO, EF	
	<i>Crassula helmsii</i> (Kirk) Cockayne	SO	
	<i>Crassula multicaulis</i> (Petrie) A.P. Druce et Given	EF	
	<i>Crassula ruamahanga</i> A.P. Druce		
	<i>Deyeuxia youngii</i> (Hook.f.) Buchanan		
	<i>Doodia mollis</i> Parris		
	<i>Epilobium petraeum</i> Heenan		
	<i>Epilobium purpuratum</i> Hook.f.		
	<i>Euchiton ensifer</i> (D.G. Drury) Holub	DP	
	<i>Euchiton paludosus</i> (Petrie) Holub	DP	
	<i>Fimbristylis velata</i> R. Br.	SO	
	<i>Fuchsia procumbens</i> R. Cunn. ex A. Cunn.		
	<i>Gentiana lineata</i> Kirk		
	<i>Gingidia trifoliolata</i> (Hook.f.)	DP	
	<i>Gnaphalium luteoalbum</i> var. <i>compactum</i> Kirk	DP, EF	
	<i>Grammitis rawlingsii</i> Parris		
	<i>Grammitis rigida</i> Hombr.	DP, SO	
	<i>Halocarpus kirkii</i> (Parl.) Quinn	RF	VU A1a+2b
	<i>Hebe annulata</i> (Petrie) Cockayne et Allan		
	<i>Hebe dilatata</i> G. Simpson et J.S. Thomson		
	<i>Hebe tairawhiti</i> B.D. Clarkson et Garn.-Jones		
	<i>Hierochloe fusca</i> Zотов	DP	
	<i>Hymenochilus tanypodus</i> (D.L. Jones, Molloy et M.A. Clem.) D.L. Jones, M.A. Clem. et Molloy	EF	
	<i>Hymenochilus tristis</i> (Colenso) D.L. Jones, M.A. Clem. et Molloy	EF, HI	
	<i>Hymenophyllum atrovirens</i> Colenso		
	<i>Hypolepis amaurorachis</i> (Kunze) Hook.	SO, EF	
	<i>Hypolepis dicksonioides</i> (Kunze) Hook.	SO, EF	
	<i>Kirkianella novae-zelandiae</i> (Hook.f.) Allan f. <i>novae-zelandiae</i>	HI	
	<i>Korthalsella salicornioides</i> (A. Cunn.) Tiegh.	EF	
	<i>Lagenifera lanata</i> A. Cunn.		
	<i>Lepidium naufragorum</i> Garn.-Jones et D.A. Norton	ST	
	<i>Leptinella dispersa</i> (D.G. Lloyd) D.G. Lloyd et C. Webb subsp. <i>dispersa</i>		
	<i>Leptinella tenella</i> (A. Cunn.) D.G. Lloyd et C. Webb		
	<i>Libocedrus plumosa</i> (D. Don) Sarg.		LR/nt
	<i>Lindsaea viridis</i> Colenso		
	<i>Macrothelypteris torresiana</i> (Gaudich.) Ching	SO, EF	
	<i>Melicytus crassifolius</i> (Hook.f.) F. Muell. s.s.	DP	
	<i>Microlaena carsei</i> Cheeseman		
	<i>Microsorum novae-zelandiae</i> (Baker) Copel.		
	<i>Mimulus repens</i> R. Br.	DP, SO	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
	<i>Muehlenbeckia ephedroides</i> Hook.f.		
	<i>Myosotis saxosa</i> Hook.f.	DP	
	<i>Myosotis spathulata</i> Forst.f. var. <i>spathulata</i>	DP	
	<i>Myosotis tenericaulis</i> Petrie	DP	
	<i>Neopaxia erythrophylla</i> Heenan	DP	
	<i>Olearia capillaris</i> Buchanan		
	<i>Olearia fragrantissima</i> Petrie		LR/nt
	<i>Olearia lineata</i> (Kirk) Cockayne	RF	
	<i>Peperomia tetraphylla</i> (G. Forst.) Hook. et Arn.	SO	
Bird catching plant, Parapara	<i>Picris angustifolia</i> subsp. <i>merxmulleri</i> Lack et S. Holzapfel	SO, EF, HI	
	<i>Pimelea poppelwellii</i> Petrie		
	<i>Pimelea pseudolyallii</i> Allan		
	<i>Pisonia brunoniana</i> Endl.	SO, HI	
	<i>Pittosporum ellipticum</i> Kirk		
	<i>Pittosporum pimeleoides</i> R. Cunn. subsp. <i>pimeleoides</i>		
	<i>Pittosporum virgatum</i> Kirk		LR/nt
	<i>Plantago obconica</i> Sykes		
	<i>Pleurosorus rutifolius</i> (R. Br.) Fée	SO	
	<i>Poa incrassata</i> Petrie		
	<i>Pomaderris hamiltonii</i> L.B. Moore	RC	LR/nt
	<i>Pseudopanax ferox</i> Kirk	CD, RF	LR/nt
	<i>Pseudopanax macintyrei</i> Cheeseman		
	<i>Pterostylis cernua</i> D.L. Jones, Molloy et M.A. Clem.	EF	
	<i>Ranunculus maculatus</i> Cockayne et Allan		
	<i>Ranunculus simulans</i> Garn.-Jones		
	<i>Raoulia beauverdii</i> Cockayne		
	<i>Rytidosperma merum</i> Connor et Edgar	DP	
	<i>Scandia rosifolia</i> (Hook.) J.W. Dawson	DP	
	<i>Schizaea dichotoma</i> (L.) J.E. Sm.	SO	
	<i>Schizeilema pallidum</i> (Kirk) Domin	DP	
	<i>Senecio carnosulus</i> (Kirk) C. Webb		
	<i>Senecio dunedinensis</i> Belcher	EF	
	<i>Senecio marotiri</i> C.J. Webb		
	<i>Senecio repangae</i> de Lange et B.G. Murray subsp. <i>repangae</i>	EF	
	<i>Solanum aviculare</i> f. <i>latifolium</i> G.T.S. Baylis		
	<i>Stegostyla atradenia</i> (D.L. Jones, Molloy et M.A. Clem.) D.L. Jones et M.A. Clem		
	<i>Sticherus flabellatus</i> (R. Br.) H. St. John	SO	
	<i>Streblus banksii</i> (Cheeseman) C. Webb		
	<i>Tetragonia tetragonoides</i> (Pallas) Kuntze	EF	
	<i>Thelymitra formosa</i> Colenso	DP, EF	
	<i>Thelymitra tholiformis</i> Molloy et Hatch	EF	
	<i>Thismia rodwayi</i> F. Muell.	DP, SO, EF	
	<i>Tmesipteris sigmatifolia</i> Chinnock		
	<i>Townsonia deflexa</i> Cheeseman		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

*Sparse—continued*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
	<i>Traversia baccharoides</i> Hook.f.		
	<i>Trichomanes colensoi</i> Hook.f.	DP	
	<i>Trisetum drucei</i> Edgar	DP	
	<i>Uncinia elegans</i> (Kük.) Hamlin		
	<i>Uncinia longifructus</i> (Kük.) Petrie		
	<i>Uncinia purpurata</i> Petrie		
	<i>Uncinia sinclairii</i> Boott		
	<i>Uncinia viridis</i> (C.B. Clarke) Edgar		
	<i>Urtica aspera</i> Petrie	DP	
	<i>Wahlenbergia congesta</i> (Cheeseman) N.E. Brown subsp. <i>congesta</i>		
<b>Taxonomically indeterminate</b>			
	<i>Coriaria</i> (a) (CHR 469745; Rimutaka)		
	<i>Dianella</i> aff. <i>nigra</i> (b) (AK 252911; Kopouatai)	DP	
	<i>Hebe</i> aff. <i>diosmifolia</i> (AK 215221; “summer flowering tetraploid”)		
	<i>Hymenophyllum</i> aff. <i>flexuosum</i> (AK 177370; Mount Burnett)		
	<i>Hypericum</i> aff. <i>japonicum</i> (b) (CHR 140620; “tarn”)	DP	
	<i>Libertia</i> aff. <i>peregrinans</i> (AK 14642; “nonoploid”)		
	<i>Myosotis</i> aff. <i>australis</i> (AK 231051; “small white”)	DP	
	<i>Myosotis spathulata</i> var. <i>radicata</i> L.B. Moore (CHR 87680)		
	<i>Myrsine</i> aff. <i>divaricata</i> (AK 228797; Poor Knights)		
	<i>Oreomyrrhis</i> (CHR 364086; “minute flower”)		
	<i>Oxalis</i> aff. <i>rubens</i> (AK 234308; “scree”)		
	<i>Peperomia</i> aff. <i>urvilleana</i> (AK 206056; “purple vein”)	DP	
	<i>Raoulia</i> (c) (CHR 401140; “M”)		
	<i>Thelymitra</i> aff. <i>ixioides</i> (AK 251348; New Zealand)	DP, SO, EF	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

*Continued on next page*

## Range Restricted

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
<b>BAT</b>			
<b>Taxonomically indeterminate</b>			
Eastern central short-tailed bat	<i>Mystacina tuberculata rhyacobia</i> (eastern)	HI	VU A2c, C2a (for <i>M. tuberculata</i> s.l.)
North-western central short-tailed bat	<i>Mystacina tuberculata rhyacobia</i> (north-western)	HI	VU A2c, C2a (for <i>M. tuberculata</i> s.l.)
<b>BIRDS</b>			
<b>Taxonomically determinate</b>			
Auckland Island teal	<i>Anas aucklandica</i>	HI, RC	NT
White-capped noddy	<i>Anous tenuirostris minutus</i>	SO	
Three Kings bellbird	<i>Anthornis melanura obscura</i>	ST, OL	
Poor Knights bellbird	<i>Anthornis melanura oneho</i>	ST, OL	
Little spotted kiwi	<i>Apteryx owenii</i>	RC, HI	VU D2
Snares Island fernbird, karoti	<i>Bowdleria punctata caudata</i>	ST, OL	NT (for <i>B. punctata</i> s.l.)
Codfish Island fernbird, matata	<i>Bowdleria punctata wilsoni</i>	RC	NT (for <i>B. punctata</i> s.l.)
Auckland Island banded dotterel	<i>Charadrius bicinctus exilis</i>	ST, HI	
Auckland Island snipe	<i>Coenocorypha aucklandica aucklandica</i>		NT (for <i>C. aucklandica</i> s.l. <sup>c</sup> )
Snares Island snipe	<i>Coenocorypha aucklandica huegeli</i>	ST, OL	NT (for <i>C. aucklandica</i> s.l. <sup>c</sup> )
Antipodes Island snipe	<i>Coenocorypha aucklandica meinertzhagenae</i>	ST	NT (for <i>C. aucklandica</i> s.l. <sup>c</sup> )
Chatham Island snipe	<i>Coenocorypha pusilla</i>	ST, HI	VU D2
Reischek's parakeet	<i>Cyanoramphus erythrotis hochstetteri</i>	ST, OL	VU D2
Chatham Island red-crowned kakariki	<i>Cyanoramphus novaezelandiae chathamensis</i>	HI	
Kermadec red-crowned kakariki	<i>Cyanoramphus novaezelandiae cyanurus</i>	HI, EF	
Antipodes Island parakeet	<i>Cyanoramphus unicolor</i>	ST, OL	VU D2
Snares Cape petrel	<i>Daption capense australe</i>		
Antipodes albatross	<i>Diomedea antipodensis</i>	RC	
Southern royal albatross, toroa	<i>Diomedea epomophora</i>	HI, RC	
Gibson's albatross	<i>Diomedea gibsoni</i>	ST	
Auckland Island rail	<i>Dryolimnas muelleri</i>	HI	VU D2
Snares crested penguin	<i>Eudyptes robustus</i>	OL, ST	
Chathams little blue penguin	<i>Eudyptula minor chathamensis</i>		
White-bellied storm petrel	<i>Fregetta grallaria grallaria</i>	TO	
Black-bellied storm petrel	<i>Fregetta tropica</i>	SO, HI, DP	
Grey-backed storm petrel	<i>Garrodia nereis</i>	SO, HI	
Chatham Island warbler	<i>Gerygone albofrontata</i>	ST, HI	
Campbell Island shag	<i>Leucocarbo campbelli</i>	OL	VU D2
New Zealand king shag	<i>Leucocarbo carunculatus</i>	ST	VU D1, D2
Auckland Island shag	<i>Leucocarbo colensoi</i>	HI	VU D2
Northern giant petrel	<i>Macronectes halli</i>	TO	
Fulmar prion	<i>Pachyptila crassirostris crassirostris</i>		
Lesser fulmar prion	<i>Pachyptila crassirostris eatoni</i>	DP, TO	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

<sup>c</sup> Subspecies likely to be elevated to species level in the future.

Continued on next page

*Range Restricted—continued*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Chatham Island fulmar prion	<i>Pachyptila crassirostris pyramidalis</i>	DP	
Snares Island tit	<i>Petroica macrocephala dannefaerdi</i>	ST, OL	
Auckland Island tit	<i>Petroica macrocephala marrineri</i>	ST	
Red-tailed tropicbird	<i>Phaethon rubricauda</i>	SO, HI	
Little black shag	<i>Phalacrocorax sulcirostris</i>	SO	
North Island saddleback, tieke	<i>Philesturnus carunculatus rufusater</i>	RC, HI	LC
White-chinned petrel	<i>Procellaria aequinoctialis</i>	TO, HI, DP	
Black petrel	<i>Procellaria parkinsoni</i>	HI	VU D2
Westland petrel	<i>Procellaria westlandica</i>	ST, OL	VU D2
Grey ternlet	<i>Procelsterna cerulea albivittata</i>	SO	
White-naped petrel	<i>Pterodroma cervicalis</i>	RC, TO, OL	
Mottled petrel	<i>Pterodroma inexpectata</i>	RC, HI	
White-headed petrel	<i>Pterodroma lessonii</i>	SO, HI	
Pycroft's petrel	<i>Pterodroma pycrofti</i>	RC, HI	VU D2
Subantarctic little shearwater	<i>Puffinus assimilis elegans</i>	SO	
North Island little shearwater	<i>Puffinus assimilis baurakiensis</i>	RC, HI	
Kermadec little shearwater	<i>Puffinus assimilis kermadecensis</i>	ST	
Buller's shearwater	<i>Puffinus bulleri</i>	ST, OL	VU D2
Wedge-tailed shearwater	<i>Puffinus pacificus</i>	SO	
Antarctic tern	<i>Sterna vittata bethunei</i>	TO, HI	
Masked booby	<i>Sula dactylatra fuliginea</i>	TO	
Southern Buller's mollymawk	<i>Thalassarche bulleri</i>		VU D2
Campbell mollymawk/New Zealand black-browed mollymawk	<i>Thalassarche impavida</i>	OL, RC	
White-capped (formerly shy) mollymawk	<i>Thalassarche steadi</i>	ST	
<b>Taxonomically indeterminate</b>			
Northern Buller's mollymawk	<i>Thalassarche</i> nov. sp.	DP	
<b>BRYOPHYTES</b>			
<b>Taxonomically determinate</b>			
Moss	<i>Aloina bifrons</i> (De Not.) Delgadillo	SO, OL	
Moss	<i>Bartramia crassinervia</i>	DP	
Moss	<i>Blindia contexta</i> (Hook.f. & Wilson) Mull. Hal.	S?O	
Moss	<i>Blindia lewinskyae</i>	DP	
Moss	<i>Blindia seppeltii</i> J.K. Bartlett & Vitt		
Moss	<i>Brachythecium subpilosum</i> (Hook.f. & Wilson) A. Jaeger	DP, SO, OL	
Moss	<i>Bryum algovicum</i> var. <i>rutheanum</i> (Warnst.) Crundw.	SO	
Moss	<i>Bryum amblyodon</i> (Mull. Hal.)	SO	
Liverwort	<i>Chiloscyphus tuberculatus</i>		
Moss	<i>Crosbya nervosa</i> (Hook.f. & Wilson) Vitt		
Liverwort	<i>Drepanolejeunea vesciculosa</i> subsp. <i>euvesciculosa</i>	SO, OL	
Moss	<i>Ectropothecium sandwichense</i> (Hook. & Arn.) Mitt. in Seem.	SO	
Moss	<i>Entostbodon muehlenbergii</i> (Turner) Fife	SO, DP	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

*Continued on next page*

*Range Restricted—continued*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Moss	<i>Gigaspernum repens</i> (Hook.) Lindb.	SO	
Moss	<i>Grimmia affinis</i> Hornsch.	DP, SO	
Moss	<i>Grimmia longirostris</i> Hook.		
Liverwort	<i>Harpalejeunea filicispis</i>	S?O	
Liverwort	<i>Herzogianthus sanguineus</i>		
Liverwort	<i>Hygrolembidium triquetrum</i>		
Liverwort	<i>Lejeunea antisophylla</i>	SO, OL	
Liverwort	<i>Lepidozia elobata</i>	DP	
Liverwort	<i>Leptolejeunea elliptica</i> subsp. <i>subacuta</i>	SO	
Moss	<i>Macromitrium brevicaule</i>	S?O	
Liverwort	<i>Marchantia macropora</i>		
Liverwort	<i>Metzgeria alpina</i>		
Moss	<i>Muelleriella angustifolia</i> (Hook.f. & Wilson) Dusén	OL	
Moss	<i>Muelleriella aucklandica</i> Vitt	OL	
Moss	<i>Notoligotrichum bellii</i> (Broth.) G.L. Smith.		
Moss	<i>Orthotbeciella varia</i> (Hedw.) Ochyra	SO	
Liverwort	<i>Paracromastigum drucei</i>		
Liverwort	<i>Plagiobchila bazzanioides</i>	OL	
Liverwort	<i>Pseudolophocolea denticulata</i>		
Moss	<i>Pyrrhobryum paramattense</i> (C. Müll.) Manuel	SO	
Moss	<i>Racomitrium crumianum</i>		
Liverwort	<i>Radula javanica</i>	DP, SO	
Liverwort	<i>Riccardia furtiva</i>	SO	
Liverwort	<i>Riccardia pennata</i>		
Liverwort	<i>Schistochila nitidissima</i>		
Moss	<i>Sphagnum perichaetiale</i>	DP, SO	
Liverwort	<i>Stolonivector fiordlandiae</i>		
Moss	<i>Timmia norvegica</i> J.E. Zetterst.	SO	
Moss	<i>Tortella mooreae</i> Sainsbury		
Moss	<i>Tortula mucronifolia</i> Schwaegr.	SO	
Moss	<i>Tortula viridipila</i> Dixon & Sainsbury		
Moss	<i>Verrucidens turpis</i> (Cardot) Cardot	SO	
<b>Taxonomically indeterminate</b>			
Liverwort	<i>Cephaloziella</i> aff. <i>pulcherrima</i>	OL	
Liverwort	<i>Frullania</i> “coastal”		
Moss	<i>Hypnobartlettia fontana</i> Ochyra	OL	
Liverwort	<i>Lejeunea</i> “no lobes”		
Liverwort	<i>Lopbolejeunea</i> “black”		
Liverwort	<i>Pallavicinia</i> aff. <i>rubristipa</i>		
Liverwort	<i>Radula</i> aff. <i>cordiloba</i> “Raoul Island”	DP	
Liverwort	<i>Zoopsis</i> “Croesus Track”		
Liverwort	<i>Zoopsis</i> “Rata Peak”		
<b>FRESHWATER FISH</b>			
<b>Taxonomically determinate</b>			
Tarndale bully	<i>Gobiomorphus alpinus</i>		VU D2
Chatham Island mudfish	<i>Neochanna rekobua</i>	DP	VU D2

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

*Continued on next page*

*Range Restricted—continued*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Stokell's smelt	<i>Stokellia anisodon</i>		
<b>Taxonomically indeterminate</b>			
Possible new non-diadromous galaxias	<i>Galaxias</i> “Northern sp.”	DP, HI	
<b>FRESHWATER INVERTEBRATES</b>			
<b>Taxonomically determinate</b>			
Freshwater isopod	<i>Austridotea benhami</i> Nicholls, 1938		
Snail	<i>Catapyrgus spelaeus</i> Climo, 1974		
Caddisfly	<i>Costachorema notopterum</i> Wise, 1972		
Caddisfly	<i>Costachorema peninsulae</i> Ward, 1995	OL	
Mayfly	<i>Cryopblebia aucklandensis</i>		
Caddis	<i>Cryptobiosella furcata</i> Henderson, 1983		
Caddis	<i>Cryptobiosella spinosa</i> Henderson, 1983	DP	
Mayfly	<i>Deleatidium (Penniketellum) cornutum</i>		
Caddisfly	<i>Edpercivalia dugdalei</i> Ward, 1998	DP	
Snail	<i>Hadopyrgus anops</i> Climo, 1974		
Snail	<i>Hadopyrgus brevis</i> Climo, 1974		
Caddisfly	<i>Helicopsyche torino</i> Johanson		
Snail	<i>Horatia nelsonensis</i> Climo, 1977		
Caddisfly	<i>Hydrobiosella aorere</i> Henderson, 1983	DP	
Caddisfly	<i>Hydrobiosis lindsayi</i> Tillyard, 1925		
Caddisfly	<i>Hydrobiosis taumata</i> Ward, 1997		
Fringed-gill mayfly	<i>Isothraulus abditus</i> Towns & Peters, 1980	DP	
Caddis	<i>Neobiosella irrorata</i> Wise, 1958		
Net-winged midge	<i>Neocurupira chiltoni</i> (Campbell, 1921)	DP	
Mayfly	<i>Nesameletus vulcanus</i> Hitchings?		
Stonefly	<i>Nesoperla johnsi</i> McLellan, 1977	OL	
Caddisfly	<i>Oecetis bathamensis</i> Tillyard, 1925		
Snail	<i>Opacuincola caeca</i> Ponder, 1966		
Snail	<i>Opacuincola kuscheli</i> Climo, 1974		
Caddisfly	<i>Oxyethira waipoua</i> Wise, 1998	DP, OL	
Snail	<i>Paxillostium nanum</i> N. Gardner, 1970		
Snail	<i>Potamopyrgus cresswelli</i> Climo, 1974	OL	
Snail	<i>Potamopyrgus dawbini</i> Powell, 1955		
Snail	<i>Potamopyrgus gardneri</i> Climo, 1974		
Snail	<i>Potamopyrgus manneringi</i> Climo, 1974	DP	
Snail	<i>Potamopyrgus subterraneus</i> (Suter, 1905)		
Snail	<i>Potamopyrgus troglodytes</i> (Climo, 1974)	OL	
Snail	<i>Saganoa inflata</i> (Climo, 1974)		
Snail	<i>Saganoa mica</i> (Climo, 1974)		
Polychaete worm	<i>Scolecolepides freemani</i>		
Caddisfly	<i>Tiphoebiosis chilarella</i> Ward, 1995	DP, OL	
Caddisfly	<i>Traillochorema rakiura</i> McFarlane, 1981		
<b>Taxonomically indeterminate</b>			
Caddisfly	<i>Alloeacentrella</i> sp. C		
Snail	Hydrobiidae sp. 1 (M.174057)	DP	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

*Continued on next page*

*Range Restricted—continued*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Snail	Hydrobiidae sp. 2 (M.174060)	DP	
Snail	Hydrobiidae sp. 3 (M.174062)	DP	
Snail	Hydrobiidae sp. 4 (M.174064)	DP	
Snail	Hydrobiidae sp. 6 (M.174075)	DP	
Snail	Hydrobiidae sp. 7 (M.174081)	DP	
Snail	Hydrobiidae sp. 8 (M.174083)	DP	
Snail	Hydrobiidae sp. 9 (M.174091)	DP	
Snail	Hydrobiidae sp. 10 (M.174095)	DP	
Snail	Hydrobiidae sp. 11 (M.174101)	DP	
Snail	Hydrobiidae sp. 12 (M.174104)	DP	
Snail	Hydrobiidae sp. 13 (M.174115)	DP	
Snail	Hydrobiidae sp. 14 (M.174117)	DP	
Snail	Hydrobiidae sp. 15 (M.174119)	DP	
Snail	Hydrobiidae sp. 16 (M.174121)	DP	
Snail	Hydrobiidae sp. 17 (M.174124)	DP	
Snail	Hydrobiidae sp. 18 (M.174131)	DP	
Snail	Hydrobiidae sp. 19 (M.174133)	DP	
Snail	Hydrobiidae sp. 20 (M.174135)	DP	
Snail	Hydrobiidae sp. 21 (M.174139)	DP	
Snail	Hydrobiidae sp. 22 (M.174051)	DP	
Snail	Hydrobiidae sp. 23 (M.174140)	DP	
Snail	Hydrobiidae sp. 24 (M.174141)	DP	
Snail	Hydrobiidae sp. 25 (M.174142)	DP	
Snail	Hydrobiidae sp. 26 (M.174143)	DP	
Snail	Hydrobiidae sp. 27 (M.174188)	DP	
Snail	Hydrobiidae sp. 28 (M.174106)	DP	
Snail	Hydrobiidae sp. 29 (M.174146)	DP	
Snail	Hydrobiidae sp. 30 (M.174148)	DP	
Snail	Hydrobiidae sp. 31 (M.174150)	DP	
Snail	Hydrobiidae sp. 32 (M.174151)	DP	
Snail	Hydrobiidae sp. 33 (M.174154)	DP	
Snail	Hydrobiidae sp. 34 (M.174193)	DP	
Snail	Hydrobiidae sp. 35 (AK 69073)	DP	
Snail	Hydrobiidae sp. 36 (M.174156)	DP	
Snail	Hydrobiidae sp. 37 (M.174163)	DP	
Snail	Hydrobiidae sp. 38 (M.174165)	DP	
Snail	Hydrobiidae sp. 39 (M.174166)	DP	
Snail	Hydrobiidae sp. 40 (M.174174)	DP	
Snail	Hydrobiidae sp. 41 (M.174175)	DP	
Snail	Hydrobiidae sp. 42 (M.174202)	DP	
Snail	Hydrobiidae sp. 43 (M.174181)	DP	
Snail	Hydrobiidae sp. 44 (M.174197)	DP	
Snail	Hydrobiidae sp. 45 (M.174184)	DP	
Snail	Hydrobiidae sp. 46 (AK48942)	DP	
Caddisfly	<i>Neurochorema</i> sp. A	DP, OL	
Caddisfly	<i>Paroxyethira</i> sp. S		
Caddisfly	<i>Paroxyethira</i> sp. WP		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

*Continued on next page*

*Range Restricted—continued*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
<b>MACROALGAE</b>			
<b>Taxonomically determinate</b>			
Green seaweed—Bryopsidales	<i>Caulerpa racemosa</i>	SO	
Green seaweed—Bryopsidales	<i>Caulerpa sertularioides</i>	SO, OL	
Green seaweed—Bryopsidales	<i>Caulerpa webbiana</i>	SO, OL	
Red seaweed—Erythropheltidales	<i>Chlidophyllum kaspar</i>	OL	
Brown seaweed—Ectocarpales	<i>Chordariopsis capensis</i>	DP, SO	
Red seaweed—Rhodymeniales	<i>Chrysymenia? polydactyla</i>	DP, OL	
Green seaweed—Bryopsidales	<i>Codium geppioporum</i>	SO, OL	
Red seaweed—Gracilariales	<i>Curdiea ballbazar</i>	OL	
Red seaweed—Nemaliales	<i>Dichotomaria marginata</i>	SO, OL	
Brown seaweed—Durvilleales	<i>Durvillaea bathamensis</i>	OL	
Red seaweed—Nemaliales	<i>Galaxaura cobaerens</i>	OL	
Red seaweed—Nemaliales	<i>Galaxaura filamentosa</i>	OL	
Red seaweed—Nemaliales	<i>Galaxaura rugosa</i>	OL	
Red seaweed—Nemaliales	<i>Ganonema farinosa</i>	OL	
Red seaweed—Gelidiales	<i>Gelidium allanti</i>	TO	
Red seaweed—Gelidiales	“ <i>Gelidium</i> ” <i>ceramoides</i>		
Red seaweed—Gelidiales	<i>Gelidium longipes</i>	OL	
Red seaweed—Gigartinales	<i>Gigartina dilatata</i>		
Red seaweed—Gigartinales	<i>Gigartina grandifida</i>	OL	
Brown seaweed—Fucales	<i>Landsburgia ilicifolia</i>	DP, OL	
Brown seaweed—Fucales	<i>Landsburgia myricifolia</i>	OL	
Brown seaweed—Laminariales	<i>Lessonia adamsiae</i>	OL	
Brown seaweed—Laminariales	<i>Lessonia brevifolia</i>		
Brown seaweed—Laminariales	<i>Lessonia tholiformis</i>	OL	
Brown seaweed—Fucales	<i>Marginariella parsonsi</i>		
Green seaweed—Chlorococcales	<i>Palmophyllum umbracola</i>		
Brown seaweed—Sporochnales	<i>Perisporocnus regalis</i>	OL	
Red alga—freshwater— Batrachospermiales	<i>Psilosiphon scoparium</i>	TO	
Red seaweed—Gelidiales	<i>Pterocladia lindaueri</i>	OL	
Red seaweed—Erythropheltidales	<i>Pyropbyllon cameronii</i>	OL	
Brown seaweed—Fucales	<i>Sargassum johnsonii</i>	OL	
Red seaweed—Gigartinales	<i>Sonderopelta coriacea</i>	SO, OL	
Red seaweed—Nemaliales	<i>Tricleocarpa cylindrica</i>	SO, OL	
<b>Taxonomically indeterminate</b>			
Brown seaweed—Durvilleales	<i>Durvillaea</i> sp.—Antipodes WELT A17080	OL	
Red seaweed—Gigartinales	<i>Gigartina</i> sp. Bounty Is.	OL	
Red seaweed—Gigartinales	<i>Gigartina</i> sp. Three Kings	OL	
Red seaweed—Gigartinales	<i>Predaea</i> sp.	OL	
<b>MARINE FISH</b>			
<b>Taxonomically determinate</b>			
Southern sculpin	<i>Antipodocottus galatheae</i>		
Long-lined sculpin	<i>Antipodocottus megalops</i>		
Ocellate triplefin	<i>Apopterygion oculus</i>		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

*Continued on next page*

Range Restricted—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Spotted sole	<i>Aseraggodes babamondei</i>		SO
Flathead	<i>Bembrops morelandi</i>		
Giant triplefin	<i>Blennodon dorsale</i>		
Galapagos shark	<i>Carcarhinus galapagensis</i>		SO
Kermadec demoiselle	<i>Chrysiptera rapanui</i> (Greenfield & Hensley, 1970)		SO
Messmate	<i>Echiodon neotes</i>		
Messmate	<i>Echiodon pegasus</i>		
Messmate	<i>Echiodon pukaki</i>		
Sole	<i>Engyprosopon raoulensis</i> Amaoka & Mihara, 1995		
Kermadec triplefin	<i>Enneapterygius kermadecensis</i> Fricke, 1994		
Fiordland brotula	<i>Fiordichthys slartibartfarsti</i>		
Subantarctic clingfish	<i>Gastrocymba quadriradiata</i>		
Caramel drummer	<i>Girella fimbriata</i> (McCulloch, 1920)		
Clingfish	<i>Kopua nutmata</i>		
Gurnard	<i>Lepidotrigla robinsi</i> Richards, 1997		
Scorpionfish	<i>Maxillicosta raoulensis</i> Eschmeyer & Poss, 1976		
Bellowsfish	<i>Notopogon xenosoma</i>		
Bluefinned butterfish	<i>Odax cyanoallix</i>		
Kermadec scalyfin	<i>Parma kermadecensis</i> Allen, 1987		
Two stripe seaperch	<i>Plectranthias bilaticlavia</i>		
Spottail seaperch	<i>Plectranthias maculicauda</i>		
Yellow goby	<i>Priolepis psygmatophila</i> Winterbottom & Burridge, 1993		
Pearlfish	<i>Pyramodon ventralis</i>		
Subantarctic triplefin	<i>Tripterygion jenningsi</i>		DP
<b>Taxonomically indeterminate</b>			
Sargeant bakers	Aulopodidae—2 undescribed spp.		S?O
Goby	<i>Eviota</i> sp.		
Dragonette	<i>Foetorepus</i> sp.		
Kermadec urchin clingfish	Gobiesocidae, undescribed genus and species		
Batfish	<i>Halieutea</i> spp.		
Stargazer	<i>Kathetostoma</i> sp.		
Longtailed rig	<i>Mustelus</i> sp.		SO
<b>MARINE INVERTEBRATES</b>			
<b>Taxonomically determinate</b>			
Limpet	<i>Actinoleuca campbelli bountyensis</i> Powell, 1956		
Snail	<i>Alcithoe davegibbsi</i> Hart, 2000		
Snail	<i>Alvania kermadecensis</i> (Oliver, 1915)		
Vent shrimp	<i>Alvinocaris longirostris</i> Kikuchi & Ohta, 1995		OL
Vent shrimp	<i>Alvinocaris niwa</i> Webber, 2004		
Snail	<i>Annulobalcis marshalli</i> Warén, 1981		
Black coral	<i>Antipathella fiordensis</i>		DP, ST
Snail	<i>Archiminolia dawsoni</i> (B.A. Marshall, 1979)		DP
Snail	<i>Archiminolia burleyi</i> (B.A. Marshall, 1979)		DP
Snail	<i>Archiminolia tenuiseptum</i> B.A. Marshall, 2000		DP
Coral	<i>Balanophyllia clmous</i> Squires		
Snail	<i>Bathyfautor rapubria</i> B.A. Marshall, 1996		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Range Restricted—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Snail	<i>Bathymopbila valentia</i> B.A. Marshall, 2000	DP	
Mantis shrimp	<i>Bathyssquilla microps</i>	SO, OL	
Clam	<i>Benthocardiella obliquata bountyensis</i> Powell, 1934	DP	
Snail	<i>Brookula stibarochila</i> (Iredale, 1912)		
Snail	<i>Caecum solitarium</i> Oliver, 1916		
Snail	<i>Calliostoma antipodense</i> B.A. Marshall, 1996		
Snail	<i>Calliostoma bentthicola</i> (Dell, 1950)		
Snail	<i>Calliostoma consobrinum</i> (Powell, 1958)		
Snail	<i>Calliostoma eminens</i> B.A. Marshall, 1996		
Snail	<i>Calliostoma gendalli</i> B.A. Marshall, 1980		
Snail	<i>Calliostoma gibbsorum</i> B.A. Marshall, 1996		
Snail	<i>Calliostoma jamiesoni</i> B.A. Marshall, 1996		
Snail	<i>Calliostoma peregrinum</i> B.A. Marshall, 1996	SO?	
Snail	<i>Calliostoma xanthos</i> B.A. Marshall, 1996	SO?	
Snail	<i>Calliotropis crystalophorus</i> B.A. Marshall, 1980	DP	
Snail	<i>Cantharidus burchorum</i> B.A. Marshall, 1999		
Snail	<i>Carinastele coronata</i> B.A. Marshall, 1989	DP	
Snail	<i>Carinastele jugosa</i> B.A. Marshall, 1989	DP	
Snail	<i>Carinastele kristelleae</i> B.A. Marshall, 1989		
Limpet	<i>Cellana craticulata</i> (Suter, 1905)		
Limpet	<i>Cellana strigilis bollonsi</i> Powell, 1955 E		
Limpet	<i>Cellana strigilis flemingi</i> Powell, 1955 E		
Limpet	<i>Cellana strigilis oliveri</i> Powell, 1955 E		
Chiton	<i>Cbiton themeropsis</i> (Iredale, 1914)		
Snail	<i>Cirsonella laxa</i> Powell, 1938		
Snail	<i>Cirsonella maoria</i> (Powell, 1937)		
Snail	<i>Cirsonella paradoxa</i> Powell, 1938		
Snail	<i>Clanculus atypicus</i> Iredale, 1913		
Coral	<i>Coenocyathus brooki</i> Cairns		
Whelk	<i>Cominella quoyana griseicalx</i> Willan, 1979		
Whelk	<i>Cominella regalis</i> Willan, 1979		
Clam	<i>Corculum inexpectatum</i> Crozier, 1967	DP	
Limpet	<i>Cornisepta festiva</i> (Crozier, 1966)		
Hydroid	<i>Crateritbeca novaezelandiae</i> (Thompson, 1879)		
Snail	<i>Crosseola favosa</i> Powell, 1938		
Snail	<i>Crosseola intertexta</i> Powell, 1938		
Scallop	<i>Cyclopecten kermadecensis</i> (E.A. Smith)	OL	
Limpet	<i>Diodora bollonsi</i> (Oliver, 1915)		
Snail	<i>Discotectonica acutissima</i> (G.B. Sowerby III, 1914)	DP, SO? (NZOI TAN107.122)	
Snail	<i>Eatoniella (E.) iredalei</i> (Oliver, 1915)		
Whelk	<i>Etrema hedleyi</i> (Oliver, 1915)		
Coral	<i>Falcatoflabellum raoulensis</i> Cairns		
Snail	<i>Fautrix candida</i> B.A. Marshall, 1996	SO?	
Snail	<i>Fictonoba oliveri</i> (Powell, 1927)		
Limpet	<i>Fissurisepta manawatawhia</i> Powell, 1938		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Range Restricted—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Snail	<i>Fuscapex opbioacanthicola</i> Warén, 1981	OL, DP	
Snail	<i>Fusceulima goodingi</i> Warén, 1981	OL, DP	
Giant vent mussel	<i>Gigantidas gladius</i> Cosel & Marshall, 2003		
Snail	<i>Graphis sculpturata</i> (Oliver, 1915)		
Clam	<i>Grippina acberontis</i> B.A. Marshall, 2002	OL	
Snail	<i>Herpetopoma pruinosa</i> B.A. Marshall, 1980		
Whelk	<i>Hexaplex puniceus</i> Oliver, 1916		
Clam	<i>Hunkydora rakiura</i> B.A. Marshall		
Whelk	<i>Iredalea subtropicalis</i> Oliver, 1916		
Whelk	<i>Kermia benhami</i> Oliver, 1916		
Snail	<i>Laevilitorina antipodum</i> (Filhol, 1880)		
Snail	<i>Laevilitorina bifasciata</i> Suter, 1914		
Snail	<i>Laevilitorina dellii</i> (Powell, 1955)		
Chiton	<i>Leptochiton norfolkensis subtropicalis</i> (Iredale, 1914)		
Crab	<i>Leptomithrax tuberculatus mortenseni</i> Bennett, 1964		
Snail	<i>Leptothyra benthicola</i> B.A. Marshall, 1980		
Snail	<i>Leptothyra kermadecensis</i> B.A. Marshall, 1980		
Whelk	<i>Lienardia roseocincta</i> (Oliver, 1915)		
Black coral	<i>Lillipathes lillei</i> (Totton, 1923)	S?O	
Snail	<i>Lissotesta conoidea</i> Powell, 1938		
Clam	<i>Lutraria bruuni</i> Powell, 1967	OL, DP	
Snail	<i>Margarella antipoda binemoa</i> Powell, 1956		
Snail	<i>Melanella kermadecensis</i> Oliver, 1916		
Snail	<i>Melanella luminosa</i> B.A. Marshall, 1997	DP	
Snail	<i>Melanella perplexa</i> Oliver, 1916		
Snail	<i>Melanella spinosa</i> Oliver, 1916		
Snail	<i>Metaxia kermadecensis</i> B.A. Marshall, 1978		
Snail	<i>Micrelenchus festivus</i> B.A. Marshall, 1999		
Monoplacophoran limpet	<i>Micropilina rakiura</i> B.A. Marshall, 1999	DP	
Monoplacophoran limpet	<i>Micropilina tangaroa</i> B.A. Marshall, 1991	DP	
Whelk	<i>Mitromorpha expeditionis</i> Oliver, 1916		
Snail	<i>Monilea incerta</i> Iredale, 1913		
Snail	<i>Munditia anomala</i> Powell, 1941		
Snail	<i>Munditia aupouria</i> Powell, 1938		
Snail	<i>Munditia delicatula</i> Powell, 1941		
Snail	<i>Munditia ebinata</i> Powell, 1938		
Snail	<i>Munditia manawatawhia</i> Powell, 1938		
Snail	<i>Munditia suteri</i> (Mestayer, 1919)		
Clam	<i>Mysella tellinula</i> (Odhner, 1924)	OL, DP	
Hydroid	<i>Nemertesia elongata</i> Totton, 1930		
Limpet	<i>Notoacmea badia</i> Oliver, 1927	DP	
Crab	<i>Notopodtes latus</i> Henderson, 1888	OL	
Coral	<i>Oculina virgosa</i> Squires, 1958		
Chiton	<i>Onithochiton oliveri</i> (Iredale, 1914)		
Snail	<i>Opbieulima fuscoapicata</i> Warén, 1981	OL, DP	
Clam	<i>Oxyperas belliana</i> (Oliver, 1915)		
Limpet	<i>Patella kermadecensis</i> (Pilsbry, 1894)		
Limpet	<i>Patinigera terroris</i> (Filhol, 1880)		
Caddisfly	<i>Philanisus fasciatus</i> Riek, 1976	OL	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Range Restricted—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Snail	<i>Pbilorene texturata</i> Oliver, 1916		
Clam	<i>Pronucula kermadecensis</i> Oliver, 1916		
Mussel	<i>Pteria avicula</i> (Holten, 1802)	DP, SO	
Snail	<i>Punctifera ophiomoeae</i> Warén, 1981	OL, DP	
Clam	<i>Purpuocardia reinga</i> (Powell, 1933)		
Snail	<i>Pusillina wallacei</i> (Oliver, 1915)		
Snail	<i>Pyramidelloides suteri</i> (Oliver, 1915)		
Snail	<i>Rastodens electra</i> (Oliver, 1915)		
Chiton	<i>Rhyssoplax exasperata</i> Iredale, 1915		
Snail	<i>Ruapukea carolus</i> Dell, 1953	DP	
Snail	<i>Scissurella satrchildi</i> Powell, 1934		
Snail	<i>Selastele kopua</i> (B.A. Marshall, 1995)		
Snail	<i>Selastele limatulum</i> (B.A. Marshall, 1995)		
Snail	<i>Selastele onustum</i> (Odhner, 1924)		
Snail	<i>Serrata raoulica</i> B.A. Marshall, 2004		
Snail	<i>Sinezona pacifica</i> (Oliver, 1915)		
Gravel maggot	<i>Smeagol climot</i> Tillier & Ponder, 1993	DP, OL	
Gravel maggot	<i>Smeagol manneringi</i> Climo, 1981	DP, OL	
Coral	<i>Spbenotrochus squiresi</i> Cairns		
Clam	<i>Spondylus ostreoides</i> E.A. Smith, 1886	DP, OL	
Clam	<i>Spondylus raoulensis</i> Oliver, 1916		
Snail	<i>Stomatella oliveri</i> (Iredale, 1912)		
Snail	<i>Sundaya exquisita</i> Oliver, 1916		
Snail	<i>Tectus royanus</i> (Iredale, 1912)		
Snail	<i>Thoristella polychroma</i> B.A. Marshall, 1999		
Whelk	<i>Trophon subtropicalis</i> Iredale, 1913		
Snail	<i>Zafra fuscolineata</i> Oliver, 1915		
Snail	<i>Zafra kermadecensis</i> Oliver, 1915		
Snail	<i>Zygoceras tropidophora</i> Warén & Bouchet, 1991	OL, DP, S?O	
<b>Taxononomically indeterminate</b>			
Clam	<i>Abra</i> sp. (NMNZ M.225609)	DP	
Scarlet Alcyonium	<i>Alcyonium</i> sp.		
Snail	<i>Amaea</i> sp. (NZOI TAN107.233)	DP, SO?	
Mussel	<i>Amygdalum</i> sp. (NMNZ M.147338)	DP, SO?	
Snail	<i>Anabathbron</i> sp. aff. <i>ovatus</i> (Powell, 1927) (NMNZ M.227089)		
Snail	<i>Ancistrobasis</i> sp. (NZOI TAN107.232)	DP	
Black coral	<i>Antipathes</i> n. sp.		
Snail	<i>Argalista</i> sp. A (NMNZ M.148551)		
Snail	<i>Argalista</i> sp. B (NMNZ M.148552)		
Snail	<i>Asterophila</i> sp. Warén & Lewis, 1994	DP	
Whelk	<i>Bellomitra</i> sp. (NZOI TAN107.127)	DP	
Clam	<i>Benthocardiella</i> sp. A (NMNZ M.148673)		
Clam	<i>Benthocardiella</i> sp. B (NMNZ M.148674)		
Clam	<i>Benthocardiella</i> sp. C (NMNZ M.148675)		
Clam	<i>Benthocardiella</i> sp. D (NMNZ M.148676)		
Crab	<i>Bythograea</i> sp. (NZOI)	OL	
Crab	<i>Calappa</i> sp. (?australasiae)	SO	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Range Restricted—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Snail	<i>Calliostoma</i> sp. (NZOI TAN107.233)	DP	
Snail	<i>Calliotropis</i> sp. A (NMNZ M.152747)	DP	
Snail	<i>Calliotropis</i> sp. B (NMNZ M.152735)	DP	
Snail	<i>Cantrainea</i> sp. A (NZOI TAN107.323)	DP	
Snail	<i>Cantrainea</i> sp. B (NZOI TAN107.323)	DP	
Snail	<i>Cantrainea</i> sp. C (NZOI TAN107.235)	DP	
Clam	<i>Carditella</i> sp. (NMNZ M.20766)		
Vent shrimp	<i>Chorocaris</i> sp. (NIWA specimen, coll. 2001)		
Limpet	<i>Clathrosepta</i> sp. (NZOI U608)	DP	
Clam	<i>Conchocele</i> sp. (NMNZ M.28418)		
Clam	<i>Condylocuna</i> sp. A (NMNZ M.144652)		
Clam	<i>Condylocuna</i> sp. B (NMNZ M.144656)		
Clam	<i>Condylocuna</i> sp. C (NMNZ M.144657)		
Clam	<i>Condylocuna</i> sp. D (NMNZ M.144658)		
Whelk	<i>Coralliphila</i> sp. (NZOI TAN107.234)	DP, SO?	
Limpet	<i>Cranopsis</i> sp. (NZOI TAN107.323)	DP	
Clam	<i>Cyamimactra</i> sp. A (NMNZ M.60854)		
Clam	<i>Cyamimactra</i> sp. B (NMNZ M.33947)		
Scallop	<i>Cycloclamys</i> sp. A (NMNZ M.107588)	DP	
Scallop	<i>Cycloclamys</i> sp. B (NMNZ M.155879)	DP	
Scallop	<i>Cyclopecten</i> sp. A (NZOI P939)	DP	
Scallop	<i>Cyclopecten</i> sp. B (NMNZ M.137443)	DP, SO?	
Snail	<i>Danilia</i> sp. (NZOI U599)	DP	
Scallop	<i>Delectopecten</i> sp. (NMNZ M.137495)	DP, SO?	
Whelk	<i>Eosipho</i> sp. (NMNZ M.150056)	DP	
Limpet	<i>Fissurellidae</i> sp. (NMNZ M.118002)		
Limpet	<i>Fissurisepta</i> sp. (NMNZ M.138467)		
Hydroid	<i>Gonaxia</i> sp. (NZOI)	OL	
Snail	<i>Granata</i> sp. (NMNZ M.148566)		
Snail	<i>Haloceras</i> sp. 1 (NZOI U573)	OL, DP	
Snail	<i>Haloceras</i> sp. 2 (M.147782)	OL, DP	
Snail	<i>Haloceras</i> sp. 3 (NZOI P941)	OL, DP	
Clam	<i>Hamacuna</i> sp. A (NMNZ M.143347)		
Clam	<i>Hamacuna</i> sp. B (NMNZ M.149015)		
Scallop	" <i>Hemipecten</i> " sp. (NZOI TAN107.138)	DP	
Snail	<i>Herpetopoma</i> sp. (NZOI TAN107.233)	DP	
Limpet	<i>Kaiparapelta</i> sp. (NMNZ M.137534)	DP	
Whelk	<i>Kapala</i> sp. (NZOI TAN107.136)	DP	
Clam	<i>Kidderia</i> sp. (NMNZ M.134975)		
Sponge	<i>Lissodendoryx</i> sp. (yellow slimy)		
Snail	<i>Margarella</i> sp. A (NMNZ M.59506)		
Snail	<i>Margarella</i> sp. B (NMNZ M.131607)		
Clam	<i>Metocardia</i> sp. (NZOI T256)	DP, SO?	
Snail	<i>Merelina</i> sp. A (NMNZ M.148669)		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

*Range Restricted—continued*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Snail	<i>Merelina</i> sp. B (NMNZ M.148670)		
Snail	<i>Merelina</i> sp. C (NMNZ M.148671)		
Snail	<i>Merelina</i> sp. D (NMNZ M.148668)		
Whelk	<i>Mitrella</i> sp. A (NZOI TAN107.233)	DP	
Whelk	<i>Mitrella</i> sp. B (NZOI TAN107.323)	DP	
Clam	<i>Mysella</i> sp. (NMNZ M.51502)		
Whelk	<i>Nassarius</i> sp. (NZOI TAN107.218)	DP, SO?	
Snail	<i>Niso</i> sp. (NZOI TAN107.225)	DP	
Clam	<i>Pectunculina</i> sp. (NMNZ M.225313)	DP	
Vent limpet	<i>Peltospiridae</i> sp. A (NZOI Z9504)	DP	
Vent limpet	<i>Peltospiridae</i> sp. B (NZOI Z9504)	DP	
Vent limpet	<i>Peltospiridae</i> sp. C (NZOI Z9504)	DP	
Clam	<i>Perrierina</i> sp. (NMNZ M.96189)		
Whelk	<i>Phymorhynchus</i> sp. (NZOI KAH11.21)	DP	
Clam	<i>Pleuromeris</i> sp. (NMNZ M.148741)		
Limpet	<i>Profundisepta</i> sp. A (NMNZ M.148575)		
Limpet	<i>Profundisepta</i> sp. B (NMNZ M.138462)		
Mussel	<i>Pteria</i> sp. (NMNZ M.158247)	DP	
Limpet	<i>Puncturella</i> sp. (NZOI U601)	DP	
Limpet	<i>Rimulanax</i> sp. (NMNZ M.225598)	DP	
Snail	<i>Serrata</i> sp. A (NMNZ M.227078)		
Vent limpet	<i>Shinkailepas</i> sp. (NZOI TAN107.228)	DP	
Snail	<i>Skeneoides</i> sp. (NMNZ M.148557)		
Snail	<i>Solariella</i> sp. A (NZOI TAN107.233)	DP	
Snail	<i>Solariella</i> sp. B (NZOI TAN107.225)	DP	
Snail	<i>Solariella</i> sp. C (NZOI W672)	DP	
Snail	<i>Solatisonax</i> aff. <i>alleryi</i> (Seguenza, 1876) (NZOI TAN107.053)	DP	
Clam	<i>Solecurtus</i> sp. (NMNZ M.225439)	DP	
Seep limpet	Sp. incertae sedis (NMNZ M.158228)		
Snail	<i>Stilapex</i> sp. (NMNZ M.232084)	OL	
Snail	<i>Stilifer</i> sp. (NMNZ M.150057)	OL, DP	
Snail	<i>Suterilla</i> sp.		
Clam	<i>Talabrida</i> sp. (NMNZ M.137651)		
Chiton	<i>Tegulaplex</i> sp. (NZOI TAN107.235)	DP	
Snail	<i>Thysanodonta</i> sp. (NMNZ M.152736)	DP	
Limpet	<i>Tugali</i> sp. (NMNZ M.36012)		
Whelk	<i>Turbinellid</i> sp. (NZOI TAN107.134)	DP	
Crab	<i>Xenograpsus</i> sp. (McLay photo of NIWA specimen in <i>University of Canterbury Alumni Journal</i> 11(1): 13)	OL	
<b>Marine mammals</b>			
Taxonomically determinate:			
New Zealand sea lion	<i>Phocarctos hookeri</i>	CD, HI	VU D2
Bottlenose dolphin	<i>Tursiops truncatus</i>	SO	DD

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

*Continued on next page*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
<b>REPTILES</b>			
<b>Taxonomically determinate</b>			
Robust skink	<i>Cyclodina alani</i>	ST, HI	VU D1+2
McGregor's skink	<i>Cyclodina macgregori</i>	RC, HI	VU D2
Marbled skink	<i>Cyclodina oliveri</i>	ST	
Fiordland skink	<i>Oligosoma acrinatum</i>	RC, HI	
Three Kings skink	<i>Oligosoma fallai</i>	ST	LR/nt
Chatham Islands skink	<i>Oligosoma nigriplantare nigriplantare</i>	HI	
Small-eared skink	<i>Oligosoma stenotis</i>		
Egg-laying skink	<i>Oligosoma suteri</i>	HI	
<b>Taxonomically indeterminate</b>			
Mokohinau skink	<i>Cyclodina</i> "Mokohinau Island"	ST, HI	
Poor Knights skink	<i>Cyclodina</i> "Poor Knights"	ST	
Anatoki gecko	<i>Hoplodactylus</i> "Anatoki"		
Kaikouras gecko	<i>Hoplodactylus</i> "Kaikouras"	DP	
Mokohinau gecko	<i>Hoplodactylus</i> "Mokohinau gecko"	RC	
Mount Arthur gecko	<i>Hoplodactylus</i> "Mount Arthur"	DP	
Poor Knights gecko	<i>Hoplodactylus</i> "Poor Knights gecko"		
Southern mini gecko	<i>Hoplodactylus</i> "southern mini"		
Three Kings gecko	<i>Hoplodactylus</i> "Three Kings gecko"	ST	
Cook Strait striped gecko	<i>Hoplodactylus stephensi</i> Cook Strait populations	HI	VU D2
Big Bay skink	<i>Oligosoma</i> "Big Bay"	DP	
Grey Valley skink	<i>Oligosoma</i> "Grey Valley"		
Cook Strait tuatara	<i>Spbenodon punctatus</i> "Cook Strait"	ST, HI	
<b>TERRESTRIAL INVERTEBRATES</b>			
<b>Taxonomically determinate</b>			
Spider—Agelenidae	<i>Abua insula</i> Forster & Wilton, 1973	OL	
Snail	<i>Allodiscus basiliratus</i> Gardner, 1967		
Snail	<i>Allodiscus cassandra</i> (Hutton, 1883)		
Snail	<i>Allodiscus cooperi</i> (Suter, 1907)		
Snail	<i>Allodiscus godeti</i> (Suter, 1891)		
Snail	<i>Allodiscus spiritus</i> Powell, 1952		
Snail	<i>Allodiscus turbotti</i> Powell, 1948		
Snail	<i>Amborhytida pycrofti</i> (Powell, 1932)		
Chatham Islands click beetle	<i>Amychus candezei</i> Pascoe, 1876		
Stiletto fly	<i>Anabarhynchus fuscofemoratus</i> Lyneborg, 1992		
Turbott's weevil	<i>Anagotus turbotti</i> (Spiller, 1942)		
Spider—Lycosidae	<i>Anoteropsis arenivaga</i> (Dalmas, 1917)		
Spider—Lycosidae	<i>Anoteropsis cantauria</i> Vink, 2002		
Spider—Lycosidae	<i>Anoteropsis flavescens</i> Vink, 2002		
Spider—Lycosidae	<i>Anoteropsis forsteri</i> Vink, 2002		
Spider—Lycosidae	<i>Anoteropsis insularis</i> Vink, 2002		
Spider—Lycosidae	<i>Anoteropsis okatainiae</i> Vink, 2002		
Spider—Lycosidae	<i>Anoteropsis ralphi</i> (Simon, 1905)		
Spider—Lycosidae	<i>Anoteropsis urquharti</i> (Simon, 1898)		
Tuatara tick	<i>Aponomma sphenodonti</i>		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Range Restricted—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Spider—Segestriidae	<i>Ariadna barbigera</i> Simon, 1905		
Spider—Lycosidae	<i>Artoria segregata</i> Vink, 2002		
Spider—Lycosidae	<i>Artoria separata</i> Vink, 2002		
Darkling beetle	<i>Artystona lata</i> Watt, 1992		
Moth	<i>Asaphodes chlorocapna</i> (Meyrick, 1925)		
Spider—Orsolobidae	<i>Ascuta australis</i> Forster, 1956	OL	
Spider—Orsolobidae	<i>Ascuta insula</i> Forster & Platnick, 1985	OL	
Carabid	<i>Bountya insularis</i> Townsend, 1971		
Spider—Stiphidiidae	<i>Cambridgea annulata</i> Dalmas, 1917		
Spider—Stiphidiidae	<i>Cambridgea australis</i> Blest & Vink, 2000	OL	
Snail	<i>Cavellia delli</i> (Climo, 1969)		
Darkling beetle	<i>Cerodolus curvellus</i> Broun, 1912		
Darkling beetle	<i>Cerodolus genialis</i> Broun, 1893		
Darkling beetle	<i>Cerodolus manepouricus</i> Watt, 1992		
Darkling beetle	<i>Cerodolus sinuatus</i> Watt, 1992		
Darkling beetle	<i>Chrysopeplus triregius</i> Watt, 1992		
Snail	<i>Climocella manawatawhia</i> (Powell, 1935)		
Snail	<i>Climocella reinga</i> Goulstone, 1996		
Spider—Clubionidae	<i>Clubiona chathamensis</i> Simon, 1905		
Cave bug	<i>Configa persephone</i> Fennah, 1975	DP	
Snail	<i>Cyrtora ampla</i> (Powell, 1941)		
Snail	<i>Cyrtora annexens</i> (Powell, 1948)	RC, OL	
Snail	<i>Cyrtora filicosta</i> (Powell, 1948)		
Snail	<i>Cyrtora bispida</i> Gardner, 1967		
Snail	<i>Cyrtora kerrana</i> Gardner, 1968		
Snail	<i>Cyrtora solitaria</i> (Powell, 1935)	RC, OL	
Snail	<i>Cyrtora tepakiensis</i> Gardner, 1967		
Snail	<i>Damonita geminoropiformis</i> Climo, 1981	DP	
Poor Knights giant weta	<i>Deinacrida fallai</i> Salmon, 1950		VU D2
Cook Strait giant weta	<i>Deinacrida rugosa</i> Buller, 1871	RC	VU D2
Giant mole weta	<i>Deinacrida talpa</i> Gibbs, 1999	DP	
Snail	<i>Delos regia</i> Climo, 1973	RC	
Snail	<i>Delos striata</i> Climo, 1973	RC, OL	
Auckland Island weta	<i>Dendroplectron aucklandensis</i>		
Snail	<i>Egestula bicolor</i> (Climo, 1973)		
Snail	<i>Egestula gaza</i> (Suter, 1909)		
Snail	<i>Egestula microgaza</i> (Climo, 1973)		
Snail	<i>Egestula pandora</i> Gardner, 1967		
Snail	<i>Flammocharopa montana</i> (Suter, 1891)		
Snail	<i>Flammoconcha stewartensis</i> Dell, 1952		
Snail	<i>Flammulina lateaperta</i> Dell, 1955		
Spider—Desidae	<i>Gasparia manneringi</i> (Forster, 1964)	OL, DP	
Spider—Desidae	<i>Gobia clarki</i> Forster, 1964	OL, DP	
Spider—Desidae	<i>Gobia falxiata</i> (Hogg, 1909)	OL, DP	
Poor Knights cave weta	<i>Gymnoplectron giganteum</i> Richards, 1962	ST	
Weevil	<i>Hadracalles fuliginosus</i>	DP	
Karo weevil	<i>Hadramphus pittospori</i>	DP	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Range Restricted—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Knobbled weevil	<i>Hadramphus stilbocarpae</i> Kuschel, 1971	HI, EF	
Spider—Linyphiidae	<i>Haplitis antipodiana</i> Blest & Vink, 2002	OL	
Spider—Linyphiidae	<i>Haplitis banksi</i> (Blest, 1979)		
Spider—Linyphiidae	<i>Haplitis chiltoni</i> (Hogg, 1911)	OL	
Spider—Linyphiidae	<i>Haplitis bornungi</i> (Blest, 1979)		
Spider—Linyphiidae	<i>Haplitis minutissima</i> (Blest, 1979)	OL	
Spider—Linyphiidae	<i>Haplitis redacta</i> (Blest, 1979)	OL	
Spider—Linyphiidae	<i>Haplitis ruficeps</i> (Urquhart, 1888)		
Spider—Linyphiidae	<i>Haplitis tegulata</i> (Blest, 1979)		
Spider—Linyphiidae	<i>Haplitis wairarapa</i> Blest & Vink, 2002	OL	
Nematode	<i>Hatterianema hollanderi</i> Chabaud & Dolfus, 1966		
Moth	<i>Helastia angusta</i> Craw, 1987	DP	
Moth	<i>Helastia clandestina</i> (Philpott, 1921)		
Moth	<i>Helastia expolita</i> (Philpott, 1917)	DP	
Moth	<i>Helastia siris</i> (Hawthorne, 1897)		
Ground weta	<i>Hemiandrus bilobatus</i> Ander, 1938		
Ground weta	<i>Hemiandrus subantarcticus</i> (Salmon, 1950)		
Banks Peninsula tree weta	<i>Hemideina ricta</i> (Hutton, 1897)		
Campbell Island ribbed weevil	<i>Heterexis seticostatus</i> (Brookes, 1951)		
Ground beetle	<i>Holcaspis bathana</i> Butcher, 1984		
Spider—Amphinectidae	<i>Huara antarctica</i> (Berland, 1931)	OL	
Spider—Amphinectidae	<i>Huara grossa</i> Forster, 1964	OL	
Spider—Amphinectidae	<i>Huara ovalis</i> (Hogg, 1909)	OL	
Spider—Huttoniidae	<i>Huttonia palpimanoides</i> O.P.-Cambridge, 1879		
Flightless shield bug	<i>Hypsithocus budsonae</i> Bergroth, 1927		
Snares Island weta	<i>Insulanoplectron spinosum</i>		
Bounty Island weta	<i>Ischyroplectron isolatum</i> (Hutton, 1895)	DP, OL	
Spider—Hahniidae	<i>Kapanga bickmani</i> (Forster, 1964)	OL	
Snail	<i>Kokikora mimiwhangata</i> Climo & Goulstone, 1995		
Spider—Desidae	<i>Laestrygones chathamensis</i> Forster, 1970		
Spider—Desidae	<i>Laestrygones minutissimus</i> (Hogg, 1909)	OL	
Spider—Linyphiidae	<i>Laetesia aucklandensis</i> (Forster, 1964)	OL	
Spider—Linyphiidae	<i>Laetesia chathami</i> Millidge, 1988		
Snail	<i>Laoma labyrinthica</i> Powell, 1948	RC	
Snail	<i>Laoma mariae aupouria</i> Cumber, 1967		
Snail	<i>Laoma minuta</i> Climo, 1971		
Snail	<i>Laomarex minuta</i> (Gardner, 1967)		
Snail	<i>Laomarex regia</i> (Gardner, 1968)	RC, OL	
Snail	<i>Laomarex sericea</i> Powell, 1948	RC	
Snail	<i>Liarea aupouria aupouria</i> Powell, 1954		
Snail	<i>Liarea bicarinata</i> (Suter, 1907)		
Snail	<i>Liarea ornata</i> Powell, 1954		
Snail	<i>Liarea partula</i> Powell, 1954		
Snail	<i>Loisthodon benbami</i> (Suter, 1909)		
Spider—Amphinectidae	<i>Mamoea grandiosa</i> Forster & Wilton, 1973		
Spider—Synotaxidae	<i>Mangua forsteri</i> (Brignoli, 1983)	OL	
Orongorongo black cicada	<i>Maoricicada myersi</i> (Fleming, 1971)		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Range Restricted—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Ground beetle	<i>Mecodema angustulum</i> Broun, 1914		DP
Ground beetle	<i>Mecodema dunense</i> Townsend, 1965		DP
Ground beetle	<i>Mecodema litoreum</i> Broun 1886		
Ground beetle	<i>Mecodema pavidum</i> Townsend, 1965		DP
Ground beetle	<i>Mecodema quoinense</i> Broun, 1912		
Ground beetle	<i>Mecodema strictum</i> Britton, 1949		DP
Ground beetle	<i>Megadromus bucolicus</i> (Broun, 1903)		
Darkling beetle	<i>Menimus borealis</i> Watt, 1992		
Darkling beetle	<i>Menimus brouni</i> Watt, 1992		
Darkling beetle	<i>Menimus clarkei</i> Watt, 1992		
Darkling beetle	<i>Menimus dubius</i> Broun, 1880		
Darkling beetle	<i>Menimus elongatus</i> Watt, 1992		
Darkling beetle	<i>Menimus belmorei</i> Watt, 1992		
Darkling beetle	<i>Menimus moehauensis</i> Watt, 1992		
Darkling beetle	<i>Menimus pubiceps</i> Broun, 1921		OL
Spider—Synotaxidae	<i>Meringa australis</i> Forster, 1990		OL
Spider—Linyphiidae	<i>Metafroneta sinuosa</i> Blest, 1979		OL
Spider—Linyphiidae	<i>Metamynoglenes gracilis</i> Blest, 1979		OL
Spider—Linyphiidae	<i>Metamynoglenes helicoides</i> Blest, 1979		
Spider—Linyphiidae	<i>Metamynoglenes magna</i> Blest, 1979		
Spider—Linyphiidae	<i>Metamynoglenes nongotaba</i> Blest & Vink, 2002		OL
Spider—Migidae	<i>Migas insularis</i> Wilton, 1968		OL
Spider—Migidae	<i>Migas sandageri</i> Goyen, 1891		OL
Darkling beetle	<i>Mimopeus clarkei</i> Watt, 1988		
Darkling beetle	<i>Mimopeus costellus</i> (Broun, 1905)		
Darkling beetle	<i>Mimopeus granulosus</i> (Breme, 1842)		
Darkling beetle	<i>Mimopeus insularis</i> Watt, 1988		
Darkling beetle	<i>Mimopeus johnsi</i> Watt, 1988		
Darkling beetle	<i>Mimopeus lateralis</i> (Broun, 1909)		
Darkling beetle	<i>Mimopeus lewisiatus</i> (Sharp, 1903)		
Darkling beetle	<i>Mimopeus parvus</i> Watt, 1988		
Darkling beetle	<i>Mimopeus pascoei</i> (Bates, 1873)		
Darkling beetle	<i>Mimopeus rugosus</i> (Bates, 1873)		
Darkling beetle	<i>Mimopeus subcostatus</i> (Sharp, 1903)		OL
Darkling beetle	<i>Mimopeus turbotti</i> Watt, 1988		OL
Darkling beetle	<i>Mimopeus vallis</i> Watt, 1988		
Cranefly	<i>Mischoderus marginatus</i> (Edwards, 1923)		DP, OL
Spider—Idiopidae	<i>Misgolas abdita</i> (Forster, 1968)		OL
Spider—Idiopidae	<i>Misgolas allani</i> (Forster, 1968)		OL
Spider—Idiopidae	<i>Misgolas dellii</i> (Forster, 1968)		
Spider—Idiopidae	<i>Misgolas insulana</i> (Forster, 1968)		OL
Spider—Idiopidae	<i>Misgolas minor</i> (Forster, 1968)		OL
Spider—Idiopidae	<i>Misgolas stephenensis</i> (Forster, 1968)		OL
Spider—Idiopidae	<i>Misgolas stewarti</i> (Todd, 1945)		OL
New Zealand bat fly	<i>Mystacinobia zelandica</i> Holloway, 1976		
Longhorn	<i>Navomorpha neglecta</i>		DP
Spider—Agelenidae	<i>Neoramia charybdis</i> (Hogg, 1910)		OL

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Range Restricted—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Spider—Agelenidae	<i>Neoramia crucifera</i> (Hogg, 1909)	OL	
Spider—Agelenidae	<i>Neoramia boggi</i> (Forster, 1964)		
Spider—Agelenidae	<i>Neoramia bokina</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Neoramia margaretae</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Neorepukia pilama</i> Forster & Wilton, 1973	OL	
Spider—Amphinectidae	<i>Neororea sorense</i> (Forster, 1955)	OL	
Campbell Island weta	<i>Notoplectron campbellense</i> Richards, 1964	OL	
Cave weta	<i>Novoplectron serratum</i> (Hutton, 1904)		
Darkling beetle	<i>Omedes nitidus</i>	DP	
Peripatus, ngaokeoke, velvetworm	<i>Ooperipatellus nanus</i> Ruhberg, 1977	DP, OL	
Spider—Agelenidae	<i>Oramia chathamensis</i> (Simon, 1899)		
Spider—Agelenidae	<i>Oramia marplesi</i> Forster, 1964	OL	
Spider—Agelenidae	<i>Oramia rubrioides</i> (Hogg, 1909)	OL	
Spider—Agelenidae	<i>Oramia solanderensis</i> Forster & Wilton, 1973	OL	
Ground beetle	<i>Oregus inaequalis</i> Castelnau, 1867		
Spider—Agelenidae	<i>Orepukia nummosa</i> (Hogg, 1909)	OL	
Spider—Agelenidae	<i>Orepukia rakiura</i> Forster & Wilton, 1973	OL	
Leech	<i>Ornitobdella edentula</i>	DP	
Spider—Desidae	<i>Otagoa chathamensis</i> Forster, 1970		
Snail	<i>Otoconcha fiordlandica</i> (Dell, 1952)		
Spider—Miturgidae	<i>Pacificana cockayni</i> Hogg, 1904	OL	
Spider—Amaurobiidae	<i>Pakeha kirki</i> (Hogg, 1909)	OL	
Spider—Amaurobiidae	<i>Pakeha stewartia</i> Forster & Wilton, 1973	OL	
Snail	<i>Palliolopdex verrucosus</i> (Simroth, 1889)		
Aphid	<i>Paradoxaphis plagianthi</i> Eastop, 2001	DP	
Spider—Linyphiidae	<i>Parafroneta marrineri</i> (Hogg, 1909)	OL	
Spider—Linyphiidae	<i>Parafroneta subalpina</i> Blest & Vink, 2002	OL	
Spider—Linyphiidae	<i>Parafroneta subantarctica</i> Blest, 1979	OL	
Snail	<i>Paralaoma buddlei</i> Powell, 1951	RC	
Snail	<i>Paralaoma manawatawhia</i> Goulstone & Brook, 1999	OL	
Snail	<i>Paralaoma pagoda</i> Climo, 1973	RC, OL	
Snail	<i>Paralaoma raki</i> Goulstone & Brook, 1999		
Snail	<i>Paralaoma regia</i> Powell, 1948		
Snail	<i>Paralaoma turbotti</i> Powell, 1948	OL	
Spider—Amphinectidae	<i>Paramamoea insulana</i> Forster & Wilton, 1973	OL, DP	
Three Kings cave weta	<i>Paraneonetus multispinus</i> Salmon	OL	
Darkling beetle	<i>Partystona metallica</i> Watt, 1992		
Sand scarab	<i>Pericopitus frontalis</i> Broun, 1904		
Peripatus, ngaokeoke, velvetworm	<i>Peripatoides indigo</i> Ruhberg, 1977		VU B1+2cd
Peripatus, ngaokeoke, velvetworm	<i>Peripatoides suteri</i>		VU B1+2c
Snail	<i>Phacussa fulminata fulminata</i> (Hutton, 1882)		
Snail	<i>Phenacobelix rusticus</i> (Suter, 1894)		
Snail	<i>Pbrixgnathus blacki</i> Powell, 1951	OL	
Snail	<i>Pbrixgnathus cognata</i> (Suter, 1909)	DP	
Snail	<i>Pbrixgnathus flemingi flemingi</i> Dell, 1950		
Snail	<i>Pbrixgnathus laqueus</i> Dell, 1950	DP	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Range Restricted—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Snail	<i>Pbrixgnathus paralaomiformis</i> (Climo, 1971)		
Snail	<i>Pbrixgnathus rakiura solanderi</i> Dell, 1954		
Snail	<i>Pbrixgnathus sorenseni</i> (Powell, 1955)		
Snail	<i>Pbrixgnathus spiralis</i> (Suter, 1896)		
Snail	<i>Pbrixgnathus subariel</i> Powell, 1948		
Snail	<i>Pbrixgnathus waipoua</i> (Gardner, 1969)	DP	
Spider—Gradungulidae	<i>Pianoa isolata</i> Forster, 1987		
Pupu harakeke	<i>Placostylus (Maoristylus) hongii</i> (Lesson, 1830)	RC	VU C1+2a, D2
Flat beetle	<i>Platisus zelandicus</i> Marris & Klimaszewski, 2001		
Spider—Orsolobidae	<i>Pounamuella australis</i> (Forster, 1964)	OL	
Spider—Orsolobidae	<i>Pounamuella vulgaris</i> (Forster, 1956)	OL, DP	
Large land snail	<i>Powelliphanta fletcheri</i> (Powell, 1938)	DP	
Large land snail	<i>Powelliphanta spedeni lateumbilicata</i> (Powell, 1946)	DP	
Moth	<i>Proditrix chionochloae</i> Dugdale, 1987	DP	
Scarab/chafer beetle	<i>Prodontria jenniferae</i> Emerson, 1997	DP	
Scarab/chafer beetle	<i>Prodontria longitarsus</i>		
Alexandra chafer beetle.	<i>Prodontria modesta</i> (Broun, 1909) s.s.		
Scarab/chafer beetle	<i>Prodontria pinguis</i> Given, 1952		
Scarab/chafer beetle	<i>Prodontria rakiurensis</i> Emerson, 1997	DP	
Scarab/chafer beetle	<i>Prodontria regalis</i> Emerson, 1997		
Bristly cockchafer	<i>Prodontria setosa</i> Given, 1952	DP	
Scarab/chafer beetle	<i>Prodontria truncata</i> Given, 1952?	DP	
Spider—Linyphiidae	<i>Protoerigone obtusa</i> Blest, 1979		
Snail	<i>Protostigmella johnsi</i> Climo, 1971		
Spider—Linyphiidae	<i>Pseudafroneta frigida</i> Blest, 1979		
Snail	<i>Pseudaneitea campbellensis</i> Burton, 1963		
Snail	<i>Pseudaneitea buttoni</i> (Suter, 1909)		
Snail	<i>Pseudaneitea multistriata</i> Burton, 1963		
Slug	<i>Pseudaneitea pallida</i> Climo 1973	ST	
Snail	<i>Pseudaneitea sorenseni</i> Powell, 1955		
Darkling beetle	<i>Pseudbelops capitalis</i> (Broun, 1917)		
Darkling beetle	<i>Pseudbelops chathamensis</i>		
Darkling beetle	<i>Pseudbelops liberalis</i> Watt, 1971		
Darkling beetle	<i>Pseudbelops posticalis</i> Broun, 1909	OL	
Darkling beetle	<i>Pseudbelops quadricollis</i> Broun, 1909		
Darkling beetle	<i>Pseudbelops tuberculatus</i> Guerin-Meneville, 1841		
Three Kings stick insect	<i>Pseudoclitarchus sentus</i> Salmon	OL	
Snail	<i>Ptychodon blacki</i> Dell, 1955		
Moth	<i>Pyrgotis transfixa</i> (Meyrick, 1924)	DP	
Bird louse	<i>Quadraceps coenocoryphae</i> Timmermann, 1955		
Bird louse	<i>Rallicola (Aptericola) pilgrimi</i> Clay, 1972		
Snail	<i>Ranfurlya constanceae</i> Suter, 1903	DP	
Snail	<i>Rhytidarex johnsoni</i> (Powell, 1948)		
Slug mite	<i>Riccardoella (Proriccardoella) novaezealandiae</i>	OL	
	Fain & Barker, 2004		
Spider—Amphinectidae	<i>Roreia aucklandensis</i> Forster & Wilton, 1973	OL	
Snail	<i>Rotadiscus insularis</i> (Climo, 1978)		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Range Restricted—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Snail	<i>Rotadiscus protoinsularis</i> Climo, 1989		
Spider—Synotaxidae	<i>Runga raroa</i> Forster, 1990	OL	
Robber fly	<i>Saropogon chathamensis</i> Hutton, 1901	DP	
Snail	<i>Serpho matthewsi</i> Suter, 1909		
Alpine grasshopper	<i>Sigaus obelisci</i> Bigelow, 1967	DP	
Alpine grasshopper	<i>Sigaus takabe</i> Morris, 2003	DP	
Nelson cave spider	<i>Spelungula cavernicola</i> Forster, 1987	ST	DD
Spider—Nemesiidae	<i>Stanwellia papua</i> (Forster, 1968)	OL	
Spider—Nemesiidae	<i>Stanwellia puna</i> (Forster, 1968)		
Spider—Nemesiidae	<i>Stanwellia taranga</i> (Forster, 1968)	OL	
Spider—Nemesiidae	<i>Stanwellia tuna</i> (Forster, 1968)	OL	
Moth	<i>Stigmella laqueorum</i> (Dugdale, 1971)		
Spider—Orsolobidae	<i>Subantarctia turbotti</i> Forster, 1955	OL	
Snail	<i>Taguabelix antipodensis</i> (Powell, 1955)		
Snail	<i>Taguabelix campbellica</i> (Filhol, 1880)		
Snail	<i>Taguabelix delicatula</i> (Powell, 1955)		
Snail	<i>Taguabelix birsuta</i> (Powell, 1955)	DP	
Cave weta	<i>Talitropsis crassicruris</i> Hutton, 1897		
Cave weta	<i>Talitropsis megatibia</i> Trewick, 1999		
Spider—Orsolobidae	<i>Tangata horningi</i> Forster & Platnick, 1985	OL	
Land amphipod	<i>Tara taranaki</i> Duncan, 1994	OL	
Spider—Micropholcommatidae	<i>Textricella insula</i> Forster, 1959	OL	
Spider—Micropholcommatidae	<i>Textricella plebeia</i> Forster, 1959		
Spider—Micropholcommatidae	<i>Textricella wisei</i> Forster, 1964		
Snail	<i>Thalassobelix antipoda</i> (Hombron & Jacquinot, 1841)		
Snail	<i>Thalassobelix chathamensis</i> (Suter, 1909)		
Snail	<i>Thalassobelix laingi</i> (Suter, 1905)		
Snail	<i>Therastiella pectinifera</i> (Powell, 1935)		
Spider—Cycloctenidae	<i>Toxopsiella horningi</i> Forster, 1979	OL, DP	
Mole cricket	<i>Triamescaptor aotea</i>		
Cave weta	<i>Turbottopleuron unicolor</i> Salmon, 1948	DP	
Spider—Orsolobidae	<i>Waiporia chathamensis</i> Forster & Platnick, 1986		
Spider—Orsolobidae	<i>Waipoua gressitti</i> (Forster, 1964)		
Darkling beetle	<i>Xylochus dentipes</i> Broun, 1886		
Darkling beetle	<i>Xylochus spinifer</i> Broun, 1893		
Darkling beetle	<i>Xylochus triregius</i> Watt, 1992		
Darkling beetle	<i>Zeadelium arthurensis</i> Watt, 1992		
Darkling beetle	<i>Zeadelium australe</i> Watt, 1992		
Darkling beetle	<i>Zeadelium femorale</i> (Broun, 1911)		
Darkling beetle	<i>Zeadelium budsoni</i> (Broun, 1909)		
Darkling beetle	<i>Zeadelium lendum</i> (Broun, 1880)		
Darkling beetle	<i>Zeadelium parvum</i> Watt, 1992		
Darkling beetle	<i>Zeadelium senile</i> Watt, 1992		
Darkling beetle	<i>Zomedes borealis</i> Watt, 1992		
<b>Taxonomically indeterminate</b>			
Weevil	<i>Allanalcis</i> sp. “Poor Knights”	DP, OL	
Snail	<i>Allodiscus</i> “coarse axials” miranda group		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Range Restricted—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Snail	<i>Allodiscus</i> “flat extreme tullia group” (NMNZ M.48100)		
Snail	<i>Allodiscus</i> “flat-ultrafine miranda group” (NMNZ M.80372)		
Snail	<i>Amborhytida</i> sp. “Motukokako” (NMNZ M.151457)		
Three Kings click beetle	<i>Amychus</i> sp.		
Weevil	<i>Anagotus</i> sp. “Three Kings”	DP	
Weevil	<i>Anagotus</i> sp. “Poor Knights”	DP	
Bladder fly	<i>Apsona</i> sp. “Kapiti Island”	DP	
Snail	“Archiphrixgnathus” “hazelwoodi” (NMNZ M.69300)		
Slug	<i>Athoracophorid</i> n. gen., n. sp. “hikurangi”		
Slug	<i>Athoracophorus</i> sp. 5 (NMNZ M.151431)		
Slug	<i>Athoracophorus</i> sp. 8 (NMNZ M.151434)		
Slug	<i>Athoracophorus</i> sp. 9 (NMNZ M.151435)		
Slug	<i>Athoracophorus</i> sp. 11 (NMNZ M.158288)	ST, OL	
	“Warawara 1”		
Snail	“Aureopunctum” “aff. 1 turbotti” (NMNZ M.99823)		
Hunter Hills grasshopper	<i>Brachaspis</i> “Hunter Hills”	DP	
Cave dipluran	<i>Burmjapyx</i> sp.	DP	
Snail	“Calcilaoma” “fordycei” (NMNZ M.88270)		
Snail	“Calcilaoma” “Kaikoura” (NMNZ M.62135)		
Snail	“Calcilaoma” “roscoei” (NMNZ M.88271)		
Snail	“Calcilaoma” “Waipara” (NMNZ M.62137)		
Snail	“Calcimocella” “aff. jacki 1”		
Snail	“Calcimocella” “aff. jacki 2” (NMNZ M.97486)		
Snail	“Calcimocella” “Martinborough limestones”		
Snail	<i>Cavellia</i> “large tapirinopsis”		
Snail	<i>Cavellia</i> “Mangarakau” (NMNZ M.129630)		
Snail	<i>Cavellia</i> “pseudanguicula” (NMNZ M.154805)		
Snail	<i>Cavellia</i> aff. <i>tapirina</i>		
Snail	<i>Charopa</i> “pororari” (NMNZ M.127808)		
Snail	Charopidae sp. 5 (NMNZ M.96786)		
Snail	Charopidae sp. 15 (NMNZ M.127828)		
Snail	Charopidae sp. 23 (NMNZ M.121526)		
Snail	Charopidae sp. 24 (NMNZ M.99795)		
Snail	Charopidae sp. 49 (NMNZ M.113791)		
Snail	Charopidae sp. 53 (NMNZ M.83043)	DP, OL	
Snail	Charopidae sp. 58 (NMNZ M.47453)		
Snail	Charopidae sp. 68 (NMNZ M.85690)		
Snail	Charopidae sp. 73 (NMNZ M.77056)		
Snail	Charopidae sp. 79 (NMNZ M.81650)		
Snail	Charopidae sp. 91 (NMNZ M.127827)		
Snail	Charopidae sp. 92 (NMNZ M.127830)		
Snail	Charopidae sp. 97 (NMNZ M.129356)		
Snail	Charopidae sp. 98 (NMNZ M.127815)		
Snail	Charopidae sp. 99 (NMNZ M.82271)		
Snail	Charopidae sp. 100 (NMNZ M.37918)		
Snail	Charopidae sp. 101 (NMNZ M.89084)		
Snail	Charopidae sp. 105 (NMNZ M.77007)		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Range Restricted—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Snail	Charopidae sp. 123 (NMNZ M.79585)	DP	
Snail	Charopidae sp. 130 (NMNZ M.127912)		
Snail	Charopidae sp. 132 (NMNZ M.127914)		
Snail	Charopidae sp. 135 (NMNZ M.89114)		
Snail	Charopidae sp. 136 (NMNZ M.24481)		
Snail	Charopidae sp. 157 (NMNZ M.85675)		
Snail	Charopidae sp. 167 (NMNZ M.99790)		
Snail	Charopidae sp. 170 (NMNZ M.160258) ( <i>Fectola</i> sp.)	DP	
Snail	Charopidae sp. 173 (NMNZ M.78451)		
Snail	Charopidae sp. 177 (NMNZ M.89841)	DP	
Snail	Charopidae sp. 182 (NMNZ M.106365)		
Snail	Charopidae sp. 184 (NMNZ M.123506)		
Snail	Charopidae sp. 193 (NMNZ M.73285)		
Snail	Charopidae sp. 204 (NMNZ M.104932)		
Snail	Charopidae sp. 205 (NMNZ M.82256)		
Snail	Charopidae sp. 214 (NMNZ M.115249)		
Snail	Charopidae sp. 221 (NMNZ M.124131)		
Snail	<i>Climocella</i> sp. "Poor Knights" (NMNZ M.57501)		
Stick insect	<i>Clitarchus</i> sp. aff. <i>bookeri</i> "Poor Knights"	DP	
Weevil	<i>Crisius</i> sp. "Poor Knights"	DP	
Snail	<i>Cyrtora</i> sp. 4 (NMNZ M.88987)		
Snail	<i>Cyrtora</i> sp. 9 (NMNZ M.110016)		
Snail	<i>Cyrtora</i> sp. 12 (NMNZ M.76290)		
Snail	<i>Cyrtora</i> sp. 13 (NMNZ M.79151)		
Snail	<i>Cyrtora</i> sp. 16 (NMNZ M.154911)	DP	
Snail	"Dellopsis" "peninsularis" (NMNZ M.88182)		
Snail	"Dellopsis" sp. (NMNZ M.89799)		
Snail	<i>Delos</i> sp. 2 (NMNZ M.38250)		
Snail	<i>Delos</i> sp. 12 (NMNZ M.154823)	RC	
Snail	<i>Discocharopa</i> "barrierensis"		
Snail	<i>Discocharopa</i> "biconcava"		
Snail	"Dorilaoma" "elevata" (NMNZ M.109635)		
Weevil	<i>Ectopsis</i> sp. "Poor Knights"	DP	
Snail	"Elsothera" "minuta" (NMNZ M.70337)		
Snail	"Elsothera" sp. (NMNZ M.77368)		
Snail	"Enodiapicis" "australis" (NMNZ M.61605)		
Weevil	<i>Exomesites</i> sp. "Poor Knights"	DP	
Snail	<i>Flammocharopa</i> "giant pilsbryi" (NMNZ M.79044)		
Snail	<i>Flammoconcha</i> "aff. marstoni" (NMNZ M.154804)		
Snail	<i>Flammulina</i> "aff. lateaperta" (NMNZ M.127905)		
Longhorn beetle	<i>Gastrosaurus</i> sp. "Poor Knights"	DP	
Moehau stag beetle	<i>Geodorcus</i> sp. "Moehau"	DP, OL	
Ground weta	<i>Hemiandrus</i> "Cromwell"		
Ground weta	<i>Hemiandrus</i> "disparalis"		
Ground weta	<i>Hemiandrus</i> "evansae"		
Ground weta	<i>Hemiandrus</i> "furoviarius"		
Ground weta	<i>Hemiandrus</i> "Horomaka"		
Ground weta	<i>Hemiandrus</i> "Kapiti"	DP	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Range Restricted—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Moehau weta.	<i>Hemiandrus</i> “Moehau”		DP
Ground weta	<i>Hemiandrus</i> “Okiwi”		
Ground weta	<i>Hemiandrus</i> “Onokis”		
Ground weta	<i>Hemiandrus</i> “Otekauri”		
Ground weta	<i>Hemiandrus</i> “Porters”		DP
Ground weta	<i>Hemiandrus</i> “saxatilis”		
Ground weta	<i>Hemiandrus</i> “vicinus”		
Cuvier Island tree weta	<i>Hemideina thoracica</i> 2n = 11,12		
Snail	“Hylaoma” sp. “globosa group” (NMNZ M.93011)		
Snail	“Hyalolaoma” “elevata” (NMNZ M.88208)		
Snail	“Hyalolaoma” “globosa” (NMNZ M.22290)		
Snail	<i>Iotula</i> “aff. microdegenerata/gracilis” (NMNZ M.62085)		
Snail	<i>Iotula</i> “minute degenerata” (NMNZ M.79488)		
Snail	<i>Iotula</i> “plana” (NMNZ M.84936)		
Snail	“Kaupotaka” “aff. lampra” (NMNZ M.99833)		
Snail	“Korapapa” “aff. aupouria” (NMNZ M.68880)		OL
Snail	“Korukoura” “australis” (NMNZ M.78966)		OL
Snail	“Korukoura” “meadsi” (NMNZ M.151454)		
Snail	<i>Laoma</i> “ballanceae” (NMNZ M.84406)		
Snail	<i>Laoma</i> “decipula” (NMNZ M.88224)		
Snail	<i>Laoma</i> “horingae” (NMNZ M.84947)		
Snail	<i>Laoma</i> “megabartleia” (NMNZ M.56655)		
Snail	<i>Laoma</i> “Mokihinui” (NMNZ M.123541)		
Snail	<i>Laoma</i> “Peraki” (NMNZ M.97024)		
Snail	<i>Laoma</i> “Rapahoe” (NMNZ M.106084)		
Snail	<i>Laoma</i> “tutuhesperus” (NMNZ M.88219)		
Snail	<i>Liarea egea</i> “Poor Knights”		
Snail	<i>Liarea turriculata</i> “Manaia” (NMNZ M.158258)		
Snail	“Litopunctum” “aff. lambata” (NMNZ M.85552)		
Snail	“Litopunctum” “aff. ordinarium” (NMNZ M.106667)		
Snail	“Litopunctum” “brunneum” (NMNZ M.89590)		
Snail	“Litopunctum” “calciphilia” (NMNZ M.106811)		
Snail	“Litopunctum” “mayhillae” (NMNZ M.89589)		
Snail	“Litopunctum” “thomsoni” (NMNZ M.114289)		OL
Snail	“Litopunctum” “viridula” (NMNZ M.89779)		OL
Snail	“Litopunctum” “whakapapaiti” (NMNZ M.82153)		
Snail	“Litopunctum” sp. “tight viridis” (NMNZ M.116438)		
Snail	<i>Magilaoma</i> “Whangaroa” (NMNZ M.160508)		OL
Ground beetle	<i>Mecodema</i> sp. “Poor Knights”		DP
Ground beetle	<i>Mecodema trailli</i> “Big South Cape”		DP
Picton/Port Underwood ground beetle	<i>Megadromus</i> sp. “eastern Sounds”		CD
Ground beetle	<i>Megadromus</i> sp. 1 (species aff. <i>sandageri</i> )		DP
Ground beetle	<i>Megadromus</i> sp. 11 “Benmore”		DP
Snail	“Microbrunnea” “aff. semiminuta/accelerata”		
Snail	“Micromocella” “aff. elliotiae”		
Snail	“Micromocella” “barrierensis” (NMNZ M.57536)		
Darkling beetle	<i>Mimopeus</i> sp. “Poor Knights”		DP

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Range Restricted—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Snail	<i>Mitodon</i> "aff. vortex" (NMNZ M.156140)		
Snail	<i>Mitodon</i> "Akatarawa"		
Snail	<i>Mitodon</i> "Puponga"		
Moth	<i>Mnesarchaea</i> sp. "Stellae"		
Snail	N. gen. "Huonodon/Mitodon" (NMNZ M.127660)		
Snail	N. gen. ( <i>Allodiscus</i> ?) "Hokianga" (NMNZ M.89827)		
Snail	N. gen. ( <i>Allodiscus</i> ?) (NMNZ M.14396)		
Robber fly	<i>Neoitamus</i> "tawahi"	DP	
Robber fly	<i>Neoitamus</i> "tetatus"	DP	
Aphid	<i>Neophyllaphis</i> sp.	DP	
Snail	<i>Obanella</i> sp. (NMNZ M.79272)		
Chafer beetle	<i>Odontria</i> sp. "Poor Knights"	DP	
Snail	"Oparara" "bartleii" (NMNZ M.55905)		
Snail	<i>Papulaoma</i> "aff. monticola" (NMNZ M.61578)		
Snail	<i>Papulaoma</i> "aff. otagoensis" (NMNZ M.99660)		
Snail	<i>Pasmaditta</i> "aff. hamiltoni" (NMNZ M.99639)		
Snail	"Phenacoloma" "sp. 1" (NMNZ M.4095)		
Snail	"Phenacoloma" "sp. 2" (NMNZ M.77762)		
Snail	"Phrixothrix" "depilis" (NMNZ M.88222)		
Weevil	<i>Pbryinxus</i> sp. "Poor Knights"	DP	
King Island turrett snail	<i>Placostylus (Basileostylus) bollonsi</i> "Northeast"	OL	VU D2 (species)
Snail	"Poripotaka" "micropallax" (NMNZ M.83424)		
Snail	"Poripotaka" "Paparoa" (NMNZ M.105749)		
Snail	"Powellaoma" "gigantea" (NMNZ M.79422)		
Snail	"Powellaoma" "peninsularis" (NMNZ M.70358)		
Large land snail	<i>Powelliphanta</i> "Egmont"		
Large land snail	<i>Powelliphanta</i> "Garibaldi"	DP, OL	
Large land snail	<i>Powelliphanta</i> "Kirwans"	DP, OL	
Large land snail	<i>Powelliphanta</i> "Lodestone"	DP	
Large land snail	<i>Powelliphanta</i> "Matakitaki"	DP	
Large land snail	<i>Powelliphanta</i> "Matiri"	DP	
Large land snail	<i>Powelliphanta</i> "Nelson Lakes"	DP	
Large land snail	<i>Powelliphanta</i> "Paparoas"	DP	
Large land snail	<i>Powelliphanta</i> "vittatus"	DP	
Large land snail	<i>Powelliphanta rossiana</i> "Fox"	DP	DD (species)
Weevil	<i>Praolepra</i> sp. "Poor Knights"	DP	
Snail	<i>Pseudaneitea</i> sp. 1 (NMNZ M.151420)		
Snail	<i>Pseudaneitea</i> sp. 4 (NMNZ M.151423)		
Snail	<i>Pseudaneitea</i> sp. 5 (NMNZ M.151424)		
Snail	<i>Pseudaneitea</i> sp. 6 (NMNZ M.151425)		
Snail	<i>Pseudaneitea</i> sp. 7 (NMNZ M.151426)		
Snail	Punctidae sp. 1 (NMNZ M.88223)		
Snail	Punctidae sp. 2 (AIM AK 155577)	OL	
Snail	Punctidae sp. 3 (NMNZ M.99812)		
Snail	Punctidae sp. 5 (NMNZ M.87995)	DP, OL	
Snail	Punctidae sp. 7 (NMNZ M.61762)		
Snail	Punctidae sp. 8 (NMNZ M.89805)		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

*Range Restricted—continued*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Snail	Punctidae sp. 9 (NMNZ M.87994)	OL, DP	
Snail	Punctidae sp. 10 (NMNZ M.62132)		
Snail	Punctidae sp. 11 (NMNZ M.79639)		
Snail	Punctidae sp. 15 (NMNZ M.84467)	DP	
Snail	Punctidae sp. 16 (NMNZ M.87987)	DP	
Snail	Punctidae sp. 17 (NMNZ M.87982)	DP	
Snail	Punctidae sp. 18 (NMNZ M.151443)	DP	
Snail	Punctidae sp. 21 (NMNZ M.62099)	DP	
Snail	Punctidae sp. 22 (NMNZ M.55454)		
Snail	Punctidae sp. 23 (NMNZ M.83503)	DP	
Snail	Punctidae sp. 25 (NMNZ M.54260)		
Snail	Punctidae sp. 28 (NMNZ M.88206)	DP	
Snail	Punctidae sp. 29 (NMNZ M.88205)	DP	
Snail	Punctidae sp. 34 (NMNZ M.37698)	DP	
Snail	Punctidae sp. 38 (NMNZ M.98347)	DP, OL	
Snail	Punctidae sp. 39 (NMNZ M.55868)	DP	
Snail	Punctidae sp. 40 (NMNZ M.78662)	DP	
Snail	Punctidae sp. 41 (NMNZ M.84465)	DP	
Snail	Punctidae sp. 42 (NMNZ M.98351)	DP	
Snail	Punctidae sp. 43 (NMNZ M.83426)	DP	
Snail	Punctidae sp. 63 (NMNZ M.68881)	DP	
Snail	Punctidae sp. 68 (NMNZ M.129971)		
Snail	Punctidae sp. 75 (NMNZ M.84948)		
Snail	Punctidae sp. 121 (NMNZ M.57797)		
Snail	Punctidae sp. 173 (NMNZ M.61625)		
Snail	Punctidae sp. 176 (NMNZ M.30449)		
Snail	Punctidae sp. 182 (NMNZ M.89837)		
Snail	Punctidae sp. 189 (NMNZ M.31873)		
Snail	Punctidae sp. 194 (NMNZ M.92955)		
Snail	Punctidae sp. 219 (NMNZ M.61568)		
Snail	Punctidae sp. 231 (NMNZ M.79385)		
Snail	Punctidae sp. 244 (NMNZ M.75477)		
Snail	Punctidae sp. 252 (NMNZ M.96782)		
Snail	Punctidae sp. 253 (NMNZ M.69812)		
Snail	Punctidae sp. 256 (NMNZ M.38192)		
Snail	Punctidae sp. 267 (NMNZ M.127727)		
Snail	“Rengakora” “aff. microreticulata” (NMNZ M.25117)		
Snail	<i>Rhytidia</i> “Wairau River”	DP	
Snail	“Rohopapa” “cassensis” (NMNZ M.38380)		
Snail	“Roscoepunctum” “embertoni” (NMNZ M.83489)		
Snail	“Roscoepunctum” “Silverstream” (NMNZ M.106729)		
Snail	<i>Rotadiscus</i> “aff. urquharti” (NMNZ M.72132)		
Weevil	<i>Scelodolichus</i> sp. “Poor Knights”	DP, OL	
Snail	“Sericococoncha” “kuscheli” (NMNZ M.88001)		
Snail	“Serratopunctum” “lutulentus” (NMNZ M.115304)	OL	
Alpine grasshopper	<i>Sigaus</i> “Remarkables”	DP, OL	
Chironomid	<i>Tanytarsus</i> sp.	DP, OL	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

*Continued on next page*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Snail	<i>Therasia</i> sp. "chathamensis" (NMNZ M.52076)		
Snail	<i>Therasiella</i> sp. "Unuwaho" (NMNZ M.160257)		
Weevil	<i>Tychanopais</i> sp. "Poor Knights"	DP	
Ground beetle	<i>Zeopoecilus</i> "Pupu Springs"	DP	
Ground beetle	<i>Zeopoecilus</i> "Stephens Island"	OL	
<b>VASCULAR PLANTS</b>			
<b>Taxonomically determinate</b>			
	<i>Abrotanella muscosa</i> Kirk		
	<i>Abrotanella patearoa</i> Heads		
	<i>Abrotanella rostrata</i> Swenson		
	<i>Abrotanella rosulata</i> Hook.f.	OL	
	<i>Abrotanella spathulata</i> Hook.f.		
	<i>Acaena microphylla</i> var. <i>pauciglochidiata</i> Bitter		
	<i>Acaena minor</i> Allan		
	<i>Acaena tesca</i> B.H. Macmill.		
	<i>Achnatherum petriei</i> (Buchanan) S.W.L. Jacobs et J. Everett	CD	
	<i>Aciphylla cartilaginea</i> Petrie		
	<i>Aciphylla congesta</i> Cheeseman		
	<i>Aciphylla crosby-smithii</i> Petrie		
	<i>Aciphylla dissecta</i> (Kirk) W.R.B. Oliv.		
	<i>Aciphylla lecomtei</i> J.W. Dawson		
	<i>Aciphylla leightii</i> Allan		
	<i>Aciphylla montana</i> var. <i>gracilis</i> (W.R.B. Oliv.) J.W. Dawson		
	<i>Aciphylla spedentii</i> Cheeseman		
	<i>Aciphylla squarrosa</i> var. <i>flaccida</i> Kirk		
	<i>Aciphylla stannensis</i> J.W. Dawson		
	<i>Aciphylla takanea</i> W.R.B. Oliv.		
	<i>Aciphylla traillii</i> Kirk		
	<i>Aciphylla trifoliolata</i> Petrie	DP	
	<i>Agrostis subulata</i> Hook.f.		
	<i>Anisotome acutifolia</i> (Kirk) Cockayne	OL	
	<i>Anisotome antipoda</i> Hook.f.	OL	
	<i>Anisotome latifolia</i> Hook.f.		
	<i>Anisotome lyallii</i> Hook.f.		
	<i>Apium prostratum</i> subsp. <i>denticulatum</i> P.S. Short	OL	
	<i>Arachniodes aristata</i> (G. Forst.) Tindale	SO, OL	
	<i>Argyrotegium nitidulum</i> (Hook.f.) J.M. Ward et Breitw.	SO	
	<i>Asplenium chathamense</i> Brownsey	OL	
	<i>Asplenium shuttleworthianum</i> Kunze	SO, OL	
	<i>Astelia nivalis</i> var. <i>morceae</i> L.B. Moore		
	<i>Atriplex billardierei</i> (Moq.) Hook.f.	CD, TO, HI	
	<i>Baumea complanata</i> (Bergg.) Blake	HI	
	<i>Brachyglottis arborescens</i> W.R.B. Oliv.	OL	EN D
	<i>Brachyglottis bifistulosa</i> (Hook.f.) B. Nord.	DP	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Range Restricted—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
	<i>Brachyglottis compacta</i> (Kirk) B. Nord.	OL	
	<i>Brachyglottis myrianthos</i> (Cheeseman) D.G. Drury		
	<i>Brachyglottis pentacopa</i> (D.G. Drury) B. Nord.	OL	VU D2
	<i>Brachyglottis stewartiae</i> (J.B. Armstr.) B. Nord.	DP	
	<i>Brachyglottis turneri</i> (Cheeseman) C. Webb		
	<i>Brachyscome humilis</i> G. Simpson et J.S. Thomson	DP	
	<i>Brachyscome linearis</i> (Petrie) Druce	DP	
	<i>Bulbinella gibbsii</i> Cockayne var. <i>gibbsii</i>		
	<i>Bulbinella rossii</i> (Hook.f.) Cheeseman		
	<i>Bulbinella talbotii</i> L.B. Moore	DP, OL	
	<i>Callitriche aucklandica</i> R. Mason		
	<i>Callitriche petriei</i> subsp. <i>chathamensis</i> R. Mason	OL	
	<i>Canavalia rosea</i> (Sw.) DC.	SO, OL	
	<i>Cardamine lacustris</i> (Garn.-Jones et P.N. Johnson) Heenan	EF	
	<i>Carex albula</i> Allan		
	<i>Carex decurata</i> Cheeseman		
	<i>Carex devia</i> Cheeseman		
	<i>Carex eltingamita</i> Hamlin	RC, OL	
	<i>Carex filamentosa</i> Petrie		
	<i>Carex frettalidis</i> Hamlin	DP	
	<i>Carex impexa</i> K.A. Ford		
	<i>Carex kermadecensis</i> Petrie	OL	
	<i>Carex opbiolithica</i> de Lange et Heenan	OL, CD, ST	
	<i>Carex pleiostachys</i> C.B. Clarke		
	<i>Carex pterocarpa</i> Petrie		
	<i>Carex rubicunda</i> Petrie		
	<i>Carex taboata</i> Hamlin		
	<i>Carex trachycarpa</i> Cheeseman		
	<i>Carex traversii</i> Kirk		
	<i>Carex uncifolia</i> Cheeseman		
	<i>Carex ventosa</i> C.B. Clarke	OL	
	<i>Carmichaelia appressa</i> G. Simpson	OL	
	<i>Carmichaelia astonii</i> G. Simpson		
	<i>Carmichaelia compacta</i> Petrie		
	<i>Carmichaelia torulosa</i> (Kirk) Heenan	ST, RF	LR/nt
	<i>Cassinia amoena</i> Cheeseman	OL, CD, ST	
	<i>Celmisia adamsii</i> var. <i>adamsii</i> Cheeseman		
	<i>Celmisia adamsii</i> var. <i>rugulosa</i> Cheeseman	OL	
	<i>Celmisia clavata</i> G. Simpson et G. Thomson		
	<i>Celmisia cockayneana</i> Petrie		
	<i>Celmisia cordatifolia</i> Buchanan var. <i>cordatifolia</i>		
	<i>Celmisia gibbsii</i> Cheeseman		
	<i>Celmisia glandulosa</i> var. <i>latifolia</i> Cockayne	OL	
	<i>Celmisia inaccessa</i> Given	DP	
	<i>Celmisia insignis</i> Martin	ST	
	<i>Celmisia lindsayi</i> Hook.f.		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Range Restricted—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
	<i>Celmisia mackaui</i> Raoul		
	<i>Celmisia macmabonii</i> var. <i>badfieldii</i> Martin		
	<i>Celmisia major</i> var. <i>brevis</i> Allan	OL	
	<i>Celmisia markii</i> Given et W.R. Lee		
	<i>Celmisia morganii</i> Cheeseman		
	<i>Celmisia philocremna</i> Given		
	<i>Celmisia polyvena</i> G. Simpson et G. Thomson	OL	
	<i>Celmisia rigida</i> (Kirk) Cockayne		
	<i>Celmisia rutlandii</i> Kirk		
	<i>Celmisia spectabilis</i> subsp. <i>lanceolata</i> (Hook.f.) D.R. Given		
	<i>Celmisia spedentii</i> G. Simpson		
	<i>Celmisia thomsonii</i> Cheeseman	OL	
	<i>Cenchrus caliculatus</i> Cav.	OL, TO	
	<i>Centrolepis minima</i> Kirk		
	<i>Chionochloa acicularis</i> Zотов		
	<i>Chionochloa antarctica</i> (Hook.f.) Zотов		
	<i>Chionochloa beddiei</i> Zотов		
	<i>Chionochloa bromoides</i> (Hook.f.) Zотов		
	<i>Chionochloa crassiuscula</i> (Kirk) Zотов subsp. <i>crassiuscula</i>		
	<i>Chionochloa crassiuscula</i> subsp. <i>directa</i> Connor		
	<i>Chionochloa defracta</i> Connor		
	<i>Chionochloa flavicans</i> f. <i>temata</i> Connor	OL	
	<i>Chionochloa juncea</i> Zотов		
	<i>Chionochloa lanea</i> Connor		
	<i>Chionochloa ovata</i> (Buchanan) Zотов		
	<i>Chionochloa rubra</i> subsp. <i>rubra</i> var. <i>inermis</i> Connor	OL	
	<i>Chionochloa spiralis</i> Zотов		
	<i>Chionobebe glabra</i> (Cheeseman) Heads	DP	
	<i>Chionobebe myosotoides</i> (Ashwin) B. Briggs et Ehrend.	DP	
	<i>Colobanthus hookeri</i> Cheeseman		
	<i>Colobanthus squarrosus</i> Cheeseman subsp. <i>squarrosus</i>		
	<i>Colobanthus squarrosus</i> subsp. <i>drucei</i> Sneddon		
	<i>Coprosma distantia</i> (de Lange et R.O. Gardner) de Lange	CD, OL	
	<i>Coprosma neglecta</i> Cheeseman		
	<i>Coprosma talbrockiei</i> L.B. Moore et R. Mason		
	<i>Cordyline kaspar</i> W.R.B. Oliv.		
	<i>Coriaria pottsiana</i> W.R.B. Oliv.		
	<i>Corokia macrocarpa</i> Kirk	CD, OL	LR/nt
	<i>Craspedia robusta</i> var. <i>pedicellata</i> (Kirk) Allan		
	<i>Cyathea kermadecensis</i> W.R.B. Oliv.	RC, OL	
	<i>Cyathea milnei</i> Hook.f.	OL	
	<i>Damnamenia vernicosa</i> (Hook.f.) D.R. Given		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Range Restricted—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
	<i>Deyeuxia lacustris</i> Edgar	DP	
	<i>Dichelachne lautumia</i> Edgar et Connor	ST	
	<i>Dicranopteris linearis</i> (Burm.f.) Underw.	SO	
	<i>Disphyma australe</i> subsp. <i>stricticaule</i> Chinnock	OL	
	<i>Disphyma papillatum</i> Chinnock	OL	
	<i>Doodia milnei</i> (Carr.) Parris	OL	
	<i>Dracophyllum arboreum</i> Cockayne		
	<i>Dracophyllum densum</i> W.R.B. Oliv.		
	<i>Dracophyllum marmoricola</i> S. Venter		
	<i>Dracophyllum ophioliticum</i> S. Venter		
	<i>Dracophyllum patens</i> W.R.B. Oliv.		
	<i>Dracophyllum trimorphum</i> W.R.B. Oliv.		
	<i>Dracophyllum uniflorum</i> var. <i>frondosum</i> G. Simpson		
	<i>Elingamita johnsonii</i> G.T.S. Baylis	OL	VU D2
	<i>Elymus apricus</i> Á. Löve et Connor		
	<i>Elymus falcis</i> Connor		
	<i>Elymus sacandros</i> Connor		
	<i>Epilobium astonii</i> (Allan) Raven et Raven		
	<i>Epilobium brevipes</i> Hook.f.		
	<i>Epilobium confertifolium</i> Hook.f.		
	<i>Epilobium forbesii</i> Allan		
	<i>Epilobium margaretiae</i> Brockie		
	<i>Epilobium wilsonii</i> Petrie		
	<i>Euphrasia drucei</i> M.B. Ashwin		
	<i>Euphrasia repens</i> Hook.f.		
	<i>Euphrasia wettsteintiana</i> du Rietz		
	<i>Ewartiothamnus sinclairii</i> (Hook.f.) Anderb.		
	<i>Festuca coxii</i> (Petrie) Hack.	OL	
	<i>Festuca luciarum</i> Connor	DP	
	<i>Festuca mattbewsii</i> subsp. <i>pisamontis</i> Connor		
	<i>Festuca ultramafica</i> Connor	OL	
	<i>Geniostoma ligustrifolium</i> var. <i>crassum</i> Cheeseman	OL, CD, ST	
	<i>Geniostoma ligustrifolium</i> var. <i>matus</i> Cheeseman	RC	
	<i>Gentiana antarctica</i> Kirk	OL	
	<i>Gentiana antipoda</i> Kirk	OL	
	<i>Gentiana astontii</i> Petrie	ST	
	<i>Gentiana bellidifolia</i> var. <i>magnifica</i> Kirk	ST	
	<i>Gentiana cerina</i> Hook.f.		
	<i>Gentiana chathamica</i> Cheeseman		
	<i>Gentiana concinna</i> Hook.f.		
	<i>Gentiana filipes</i> Cheeseman		
	<i>Gentiana gibbsii</i> Petrie	OL	
	<i>Gentiana lilliputiana</i> C. Webb	EF	
	<i>Geranium traversii</i> Hook.f.	OL	
	<i>Geum divergens</i> Cheeseman		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Range Restricted—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
	<i>Gingidia enystii</i> (Kirk) J.W. Dawson	DP	
	<i>Gingidia flabellata</i> (Kirk) J.W. Dawson		
	<i>Haastia pulvinaris</i> var. <i>minor</i> Laing		
	<i>Haastia recurva</i> var. <i>wallii</i> Cockayne	DP	
	<i>Haloragis erecta</i> subsp. <i>cartilaginea</i> (Cheeseman) Orchard	OL, CD, ST	
	<i>Hebe acutiflora</i> Cockayne		
	<i>Hebe adamsii</i> (Cheeseman) Cockayne et Allan	OL	
	<i>Hebe amplexicaulis</i> (J.B. Armstr.) Cockayne et Allan f. <i>amplexicaulis</i>		
	<i>Hebe amplexicaulis</i> f. <i>birta</i> Garn.-Jones et Molloy		
	<i>Hebe arganthera</i> Garn.-Jones, Bayly, W.G. Lee et Rance		
	<i>Hebe benthamii</i> (Hook.f.) Cockayne et Allan		
	<i>Hebe biggarii</i> Cockayne		
	<i>Hebe brevifolia</i> (Cheeseman) de Lange	OL, CD, ST	
	<i>Hebe calcicola</i> Bayly et Garn.-Jones		
	<i>Hebe carnosula</i> (Hook.f.) Cockayne		
	<i>Hebe chathamica</i> (Buchanan) Cockayne et Allan		
	<i>Hebe dieffenbachii</i> (Benth.) Cockayne et Allan		
	<i>Hebe elliptica</i> var. <i>crassifolia</i> Cockayne et Allan		
	<i>Hebe evenosa</i> (Petrie) Cockayne et Allan		
	<i>Hebe gibbsii</i> (Kirk) Cockayne et Allan	DP	
	<i>Hebe insularis</i> (Cheeseman) Cockayne et Allan		
	<i>Hebe macrocalyx</i> (J.B. Armstr.) G. Simpson var. <i>macrocalyx</i>		
	<i>Hebe obtusata</i> (Cheeseman) Cockayne et Allan		
	<i>Hebe ochracea</i> M.B. Ashwin		
	<i>Hebe pareora</i> Garn.-Jones et Molloy		
	<i>Hebe pauciflora</i> G. Simpson et J.S. Thomson	DP	
	<i>Hebe ramosissima</i> G. Simpson et J.S. Thomson		
	<i>Hebe rigidula</i> (Cheeseman) Cockayne et Allan var. <i>rigidula</i>		
	<i>Hebe rigidula</i> var. <i>sulcata</i> Bayly et Kellow	DP	
	<i>Hebe stenophylla</i> var. <i>besperia</i> Bayly et Garn.-Jones		
	<i>Hebe stenophylla</i> var. <i>oliveri</i> Bayly et Garn.-Jones	OL	
	<i>Hebe townsonii</i> (Cheeseman) Cockayne et Allan		
	<i>Hebe tumida</i> (Kirk) Cockayne et Allan		
	<i>Hebe urvilleana</i> W.R.B. Oliv.		
	<i>Helichrysum plumeum</i> Allan		
	<i>Helichrysum selago</i> var. <i>tumidum</i> Cheeseman		
	<i>Hierochloe brunonis</i> Hook.f.		
	<i>Hoberia equitum</i> Heads		
	<i>Homalanthus polyandrus</i> (Hook.f.) Cheeseman	OL	VU D2
	<i>Imperata cheesemanii</i> Hack.	OL	
	<i>Ipomoea pes-caprae</i> subsp. <i>brasiliensis</i> (L.) Ooststr.	SO	
	<i>Isolepis crassiuscula</i> Hook.f.	SO	
	<i>Juncus scheuchzeroides</i> Gaudich.	SO	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Range Restricted—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
	<i>Kirkianella novae-zelandiae</i> f. <i>glauca</i> Allan	ST, HI	
	<i>Kunzea ericoides</i> var. <i>microflora</i> (G. Simpson) W. Harris		
	<i>Kunzea sinclairii</i> (Kirk) W. Harris		
	<i>Lachnagrostis ammobia</i> Edgar	DP	
	<i>Lachnagrostis leptostachys</i> (Hook.f.) Zотов		
	<i>Lachnagrostis pilosa</i> subsp. <i>nubifera</i> Edgar		
	<i>Lachnagrostis tenuis</i> (Cheeseman) Edgar		
	<i>Leptecophylla robusta</i> (Hook.f.) Weiller	OL	
	<i>Leptinella albida</i> (D.G. Lloyd) D.G. Lloyd et C. Webb		
	<i>Leptinella atrata</i> subsp. <i>luteola</i> (D.G. Lloyd) D.G. Lloyd et C. Webb		
	<i>Leptinella calcarea</i> (D.G. Lloyd) D.G. Lloyd et C. Webb		
	<i>Leptinella dispersa</i> subsp. <i>rupestris</i> (D.G. Lloyd) D.G. Lloyd et C. Webb	RF	
	<i>Leptinella lanata</i> Hook.f.		
	<i>Leptinella minor</i> Hook.f.		
	<i>Leptinella plumosa</i> Hook.f.	SO	
	<i>Leptinella potentillina</i> F. Muell.		
	<i>Leptinella pyrethrifolia</i> var. <i>linearifolia</i> (Cheeseman) D.G. Lloyd et C. Webb	OL	
	<i>Leptinella traillii</i> (Kirk) D.G. Lloyd et C. Webb subsp. <i>traillii</i>		
	<i>Leptinella traillii</i> subsp. <i>puchella</i> (Kirk) D.G. Lloyd et C. Webb		
	<i>Leucogenes neglecta</i> Molloy		
	<i>Leucopogon parviflorus</i> (Andrews) Lindl.	OL, SO	
	<i>Lignocarpa diversifolia</i> (Cheeseman) J.W. Dawson		
	<i>Luzula crenulata</i> Buchanan		
	<i>Luzula traversii</i> var. <i>tenuis</i> Edgar	DP	
	<i>Macropiper excelsum</i> subsp. <i>peltatum</i> f. <i>delangei</i> R.O. Gardner	OL, RC	
	<i>Macropiper excelsum</i> subsp. <i>peltatum</i> R.O. Gardner f. <i>peltatum</i>		
	<i>Macropiper excelsum</i> subsp. <i>psittacorum</i> (Endl.) Sykes	OL, SO	
	<i>Macropiper melchior</i> Sykes	OL	
	<i>Melicytus chathamicus</i> (F. Muell.) Garn.-Jones	OL	
	<i>Melicytus drucei</i> Molloy et B.D. Clarkson	CD, RC, OL	
	<i>Melicytus obovatus</i> (Kirk) Garn.-Jones s.s.		
	<i>Meryta sinclairii</i> (Hook.f.) Seem		VU D2
	<i>Metrosideros kermadecensis</i> W.R.B. Oliv.	OL	
	<i>Mitrasacme montana</i> var. <i>helmsii</i> Kirk		
	<i>Myoporum kermadecense</i> Sykes	OL	
	<i>Myosotis antarctica</i> Hook.f.	OL	
	<i>Myosotis arnoldii</i> L.B. Moore		
	<i>Myosotis capitata</i> Hook.f.	DP	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Range Restricted—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
	<i>Myosotis concinna</i> Cheeseman		
	<i>Myosotis eximia</i> Petrie		
	<i>Myosotis explanata</i> Cheeseman	DP	
	<i>Myosotis laeta</i> Cheeseman	OL	
	<i>Myosotis matthewsii</i> L.B. Moore	DP, EF	
	<i>Myosotis monroi</i> Cheeseman	OL	
	<i>Myosotis oreophila</i> Petrie	HI, EF	
	<i>Myosotis raktura</i> L.B. Moore		
	<i>Myrsine coxii</i> Cockayne	OL, RF	
	<i>Myrsine kermadecensis</i> Cheeseman	OL	
	<i>Myrsine oliveri</i> Allan	OL	VU D2
	<i>Neopaxia racemosa</i> (Buchanan) Heenan	OL	
	<i>Nephrolepis birtsutula</i> (G. Forst.) C. Presl	SO, OL	
	<i>Olearia allomii</i> Kirk	OL	
	<i>Olearia bathamica</i> Kirk	OL	LR/nt
	<i>Olearia crosby-smithiana</i> Petrie		
	<i>Olearia semidentata</i> Decne.	OL	
	<i>Olearia traversii</i> (F. Muell.) Hook.f.	CD, RC, OL	LR/nt
	<i>Oreoporanthera alpina</i> (Cheeseman) Orchard		
	<i>Ourisia confertifolia</i> Arroyo		
	<i>Ourisia goulandiana</i> Arroyo	DP	
	<i>Ourisia remotifolia</i> Arroyo		
	<i>Ourisia spathulata</i> Arroyo	DP	
	<i>Pachycladon crenatus</i> Phillipson		
	<i>Pachycladon wallii</i> (Carse) Heenan et A. Mitch.		
	<i>Pachystegia rufa</i> Molloy	CD, ST	
	<i>Parsonsia praeruptis</i> Heads et de Lange	CD, OL	
	<i>Petalochilus alatus</i> (R. Br.) D.L. Jones et M.A. Clem.	DP, TO	
	<i>Pimelea crosby-smithiana</i> Petrie		
	<i>Pimelea microphylla</i> Colenso		
	<i>Pimelea suteri</i> Kirk		
	<i>Pittosporum fairchildii</i> Cheeseman	OL	VU D2
	<i>Pittosporum pimeleoides</i> subsp. <i>matius</i> (Cheeseman) R.C. Cooper	CD, OL, ST	
	<i>Plantago aucklandica</i> Hook.f.	OL	
	<i>Plantago spathulata</i> subsp. <i>picta</i> (Colenso) Sykes		
	<i>Plantago triantha</i> Spreng.	SO	
	<i>Pleurophyllum criniferum</i> Hook.f.		
	<i>Pleurophyllum hookeri</i> Buchanan		
	<i>Pleurophyllum speciosum</i> Hook.f.		
	<i>Poa acicularifolia</i> Buchanan subsp. <i>acicularifolia</i>		
	<i>Poa acicularifolia</i> subsp. <i>ophitalis</i> Edgar		
	<i>Poa anceps</i> subsp. <i>polypylla</i> (Hack.) Edgar		
	<i>Poa antipoda</i> Petrie		
	<i>Poa aucklandica</i> Petrie subsp. <i>aucklandica</i>		
	<i>Poa aucklandica</i> subsp. <i>campbellensis</i> (Petrie) Edgar		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Range Restricted—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
	<i>Poa aucklandica</i> subsp. <i>rakiura</i> Edgar	DP, OL	
	<i>Poa chathamica</i> Petrie		
	<i>Poa foliosa</i> (Hook.f.) Hook.f.		
	<i>Poa pygmaea</i> Buchanan		
	<i>Poa ramosissima</i> Hook.f.		
	<i>Poa senex</i> Edgar		
	<i>Poa tenuantiana</i> Petrie		
	<i>Poa xenica</i> Edgar et Connor		
	<i>Pomaderris paniculosa</i> subsp. <i>novaeczelandiae</i> (L.B. Moore) N.G. Walsh		
	<i>Pomaderris rugosa</i> Cheeseman		
	<i>Poranthera microphylla</i> Brong.	SO	
	<i>Pseudopanax chathamicus</i> Kirk		
	<i>Pseudopanax gilliesii</i> Kirk		LR/nt
	<i>Pseudopanax kermadecensis</i> (W.R.B. Oliv.) Phillipson	RC, OL	
	<i>Pterostylis silvicolatrix</i> (F. Muell.) Molloy, D.L. Jones et M.A. Clem.	OL	
	<i>Puccinellia walkeri</i> (Kirk) Allan subsp. <i>walkeri</i>		
	<i>Puccinellia walkeri</i> subsp. <i>antipoda</i> (Petrie) Edgar	OL	
	<i>Rachelia glaria</i> J.M. Ward et Breitw.	DP	
	<i>Ranunculus brevis</i> Garn.-Jones		
	<i>Ranunculus godleyanus</i> Hook.f.	CD	
	<i>Ranunculus grahamii</i> Petrie		
	<i>Ranunculus haastii</i> subsp. <i>piliferus</i> F.J.F. Fisher		
	<i>Ranunculus kirkitii</i> Petrie		
	<i>Ranunculus mirus</i> Garn.-Jones		
	<i>Ranunculus pinguis</i> Hook.f.		
	<i>Ranunculus recens</i> var. <i>lacustris</i> G. Simpson		
	<i>Ranunculus scribalis</i> Garn.-Jones		
	<i>Ranunculus stylosus</i> H.D. Wilson et Garn.-Jones		
	<i>Ranunculus subscapus</i> Hook.f.	OL	
	<i>Ranunculus viridis</i> H.D. Wilson et Garn.-Jones	DP, OL	
	<i>Raoulia cinerea</i> Petrie		
	<i>Raoulia goyenii</i> Kirk		
	<i>Raoulia hectorii</i> var. <i>mollis</i> Buchanan		
	<i>Raoulia petriensis</i> Kirk		
	<i>Raoulia rubra</i> Buchanan		
	<i>Rhopalostylis cheesemanii</i> Becc.		
	<i>Rumex neglectus</i> Kirk	HI	
	<i>Rytidosperma nudum</i> (Hook.f.) Connor et Edgar	DP	
	<i>Rytidosperma petrosum</i> Connor et Edgar		
	<i>Scaevola gracilis</i> Hook.f.	TO, OL	
	<i>Schizolema</i> (a) (CHR 190698; Ruahine)		
	<i>Schizolema reniforme</i> (Hook.f.) Domin		
	<i>Schoenus fluitans</i> Hook.f.	SO	
	<i>Senecio glaucophyllus</i> Cheeseman subsp. <i>glaucophyllus</i>		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
	<i>Senecio glaucocephalus</i> subsp. <i>basinudus</i> Ornduff		
	<i>Senecio hawaii</i> Sykes	ST	
	<i>Senecio lautus</i> var. <i>esperensis</i> Sykes	EF, OL	
	<i>Senecio radiolatus</i> F. Muell. subsp. <i>radiolatus</i>	EF	
	<i>Senecio radiolatus</i> subsp. <i>antipodus</i> (Kirk) C. Webb	OL, EF	
	<i>Senecio repandae</i> subsp. <i>pokohinuensis</i> de Lange et B.G. Murray	OL, EF	
	<i>Senecio sterquilinus</i> Ornduff	HI, EF	
	<i>Simplicia buchananii</i> (Zotov) Zotov	DP	
	<i>Sophora longicarinata</i> G. Simpson et J.S. Thomson		
	<i>Sophora molloyi</i> Heenan et de Lange		
	<i>Sporadanthus ferrugineus</i> de Lange, Heenan et B.D. Clarkson	CD, ST, HI	
	<i>Sporadanthus traversii</i> (F. Muell.) F. Muell.	CD, ST	
	<i>Sprengelia incarnata</i> Sm.		
	<i>Stellaria decipiens</i> Hook.f. var. <i>decipiens</i>		
	<i>Stellaria decipiens</i> var. <i>angustata</i> Kirk		
	<i>Stenostachys deceptorix</i> Connor	DP	
	<i>Stilbocarpa lyallii</i> J.B. Armstr.	RC	
	<i>Stilbocarpa polaris</i> (Hombr. et Jacq.) Gray		
	<i>Stilbocarpa robusta</i> (Kirk) Cockayne	OL	
	<i>Streblus smithii</i> (Cheeseman) Corner		
	<i>Trisetum serpentinum</i> Edgar et A.P. Druce		
	<i>Wahlenbergia akaroa</i> J.A. Petterson		
	<i>Wahlenbergia albomarginata</i> subsp. <i>flexilis</i> (Petrie) J.A. Petterson		
	<i>Wahlenbergia albomarginata</i> subsp. <i>olivina</i> J.A. Petterson		
	<i>Wahlenbergia cartilaginea</i> Hook.f.		
	<i>Wahlenbergia matthewsi</i> Cockayne	DP	
	<i>Wahlenbergia pygmaea</i> subsp. <i>drucei</i> J.A. Petterson		
	<i>Xeronema callistemon</i> f. <i>bracteosa</i> (L.B. Moore) de Lange et E.K. Cameron	OL	
	<i>Xeronema callistemon</i> W.R.B. Oliv. f. <i>callistemon</i>		
<b>Taxonomically indeterminate</b>			
	<i>Ascarina lucida</i> var. <i>lanceolata</i> (Hook.f.) Allan (AK 233919)		
	<i>Astelia</i> aff. <i>graminea</i> (CHR 129122; Red Hills)	OL	
	<i>Astelia</i> aff. <i>nervosa</i> (a) (AK 108205; Mount Stokes)	DP, OL	
	<i>Astelia</i> aff. <i>nervosa</i> (b) (CHR 355412; Stewart Island)		
	<i>Brachyscome</i> aff. <i>humilis</i> (AK 231703; West Dome)	DP, OL	
	<i>Cardamine</i> (e) (AK 231673; West Dome)	DP, OL	
	<i>Carex</i> (a) (CHR 395744; Takaka)		
	<i>Carex</i> (b) (AK 232856; Matiri)		
	<i>Celmisia</i> aff. <i>major</i> (AK 255352; Pupu)		
	<i>Chionochloa</i> (a) (OTA 18879; Mt Burns)	DP	
	<i>Colobanthus</i> (b) (AK 232645; "Red Hills")		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Range Restricted—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
	<i>Colobanthus</i> aff. <i>wallii</i> (AK 232551; “serpentine”)		
	<i>Coprosma</i> aff. <i>acerosa</i> (AK 36799; Taranaki)		
	<i>Coprosma</i> aff. <i>neglecta</i> (AK 250769; Whangaroa)		
	<i>Craspedia</i> (c) (CHR 529115; Kaitorete)	CD, HI, OL	
	<i>Craspedia</i> (d) (CHR 245893; Otakeho)	DP	
	<i>Craspedia</i> (f) (CHR 514362; Hackett)	OL	
	<i>Craspedia</i> (g) (CHR 469764; Pikikiruna)	OL	
	<i>Craspedia</i> (h) (CHR 260312; Gouland Downs)	DP	
	<i>Craspedia</i> (k) (CHR 283173; “coast”)		
	<i>Craspedia</i> (l) (CHR 479212; Charleston)	DP, OL	
	<i>Craspedia</i> (o) (CHR 471883; Loveridge)	DP, OL	
	<i>Craspedia</i> (p) (CHR 469073; Havelock River)	OL	
	<i>Craspedia</i> (q) (AK 251905; Anglem)	DP, OL	
	<i>Craspedia</i> (r) (CHR 313349; Punakaiki)	DP, OL	
	<i>Craspedia</i> (s) (CHR 401645; “serpentine”)		
	<i>Craspedia</i> (t) (CHR 365392; Chalk)		
	<i>Craspedia</i> aff. <i>minor</i> (AK 228074; Chatham Island)		
	<i>Deyeuxia</i> aff. <i>quadrisetata</i> (AK 252511; Volcanic Plateau)		
	<i>Dianella</i> aff. <i>nigra</i> (a) (AK 256873; Hauturu)		
	<i>Epilobium</i> aff. <i>glabellum</i> (CHR 387893; “pink”)	DP	
	<i>Gentiana</i> (a) (CHR 395723; Lookout Range)		
	<i>Gentiana</i> aff. <i>astonii</i> (c) (CHR 519113; Awahokomo)	EF, OL	
	<i>Gentiana</i> aff. <i>astonii</i> (d) (CHR 529114; Ward)	ST	
	<i>Gentiana</i> aff. <i>astonii</i> (f) (CHR 279272; Chalk Range)		
	<i>Gentiana</i> aff. <i>tenuifolia</i> (CHR 387194; “stellar”)		
	<i>Geranium</i> (a) (CHR 518296; Pareora River)	OL	
	<i>Geranium</i> (b) (CHR 469918; Red Hills)	ST, OL	
	<i>Gingidia</i> aff. <i>montana</i> (b) (CHR 103349; North Otago)	DP	
	<i>Hebe</i> aff. <i>ligustrifolia</i> (AK 207101; Surville Cliffs)		
	<i>Hebe</i> aff. <i>pinguifolia</i> (CHR 461354; “high flyer”)	DP	
	<i>Hebe pubescens</i> subsp. <i>rebuarum</i> Bayly et de Lange		
	<i>Hebe pubescens</i> subsp. <i>sejuncta</i> Bayly et de Lange		
	<i>Hebe</i> sp. ( <i>Veronica salicifolia</i> var. <i>angustissima</i> Cockayne) (AK 233637)		
	<i>Helichrysum</i> aff. <i>aggregatum</i> (AK 54473; Surville Cliffs)	CD, ST, OL	
	<i>Helichrysum</i> aff. <i>intermedium</i> (CHR 274826; Chalk Range)		
	<i>Lastreopsis</i> aff. <i>glabella</i> (AK 242151; Kermadecs)	OL	
	<i>Leucopogon xerampelinus</i> de Lange, Heenan et M.I. Dawson	OL, CD, ST	
	<i>Melicytus</i> aff. <i>alpinus</i> (g) (CHR 541569; Blondin)		
	<i>Melicytus</i> aff. <i>crassifolius</i> (CHR 279358; “cliff”)		
	<i>Melicytus</i> aff. <i>novae-zelandiae</i> (CHR 89907; “maritime”)	ST	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

*Range Restricted—continued*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
	<i>Melicytus ramiflorus</i> subsp. (a) (AK 207155; Three Kings)	DP	
	<i>Melicytus ramiflorus</i> subsp. (b) (AK 234207; Raoul)	OL	
	<i>Myosotis</i> (a) (CHR 320240; Mossburn)		
	<i>Myosotis</i> aff. <i>brockiei</i> (CHR 497375; Lake Otuhié)	OL	
	<i>Nephrolepis</i> (a) (AK 232904; “thermal”)	TO	
	<i>Notothlaspi</i> (a) (CHR 363071; Red Hills)	OL	
	<i>Ourisia</i> aff. <i>caespitosa</i> (CHR 395703; Hope Range)		
	<i>Parabebe</i> aff. <i>catarractae</i> (CHR 324810; “hairy”)	DP	
	<i>Phyllocladus</i> aff. <i>trichomanoides</i> (AK 138493; Surville Cliffs)	OL, CD, ST	
	<i>Pimelea</i> (b) (AK 165780; Mt Manaia)	ST	
	<i>Pimelea</i> (c) (CHR 511713; “tarn”)	DP	
	<i>Pimelea</i> (d) (CHR 472016; Pisa)		
	<i>Pimelea</i> aff. <i>aridula</i> (c) (CHR 402249; Moawhango)	ST, OL	
	<i>Pimelea</i> aff. <i>aridula</i> (d) (CHR 221089; Maungaharuru)	OL	
	<i>Pimelea</i> aff. <i>sericeovillosa</i> (CHR 467766; Cobb)		
	<i>Pimelea</i> aff. <i>tomentosa</i> (b) (AK 130893; Surville Cliffs)	OL, CD, ST	
	<i>Pimelea</i> aff. <i>tomentosa</i> (c) (AK 228145; Three Kings)	OL	
	<i>Pseudopanax</i> aff. <i>lessonii</i> (AK 46066; Surville Cliffs)	CD	
	<i>Pteris</i> aff. <i>macilenta</i> (AK 210045; Punakaiki)	DP	
	<i>Raoulia</i> aff. <i>bryoides</i> (AK 323119; “L”)	DP, ST, OL	
	<i>Rubus</i> aff. <i>schmidelioides</i> (CHR 325720; “strawberry”)		
	<i>Stellaria</i> aff. <i>parviflora</i> (AK 169580; Poor Knights)		
	<i>Thelymitra</i> (b) (CHR 518036; “darkie”)	EF	
	<i>Thelymitra</i> (c) (CHR 518036; “rough leaf”)	EF	
	<i>Wahlenbergia congesta</i> subsp. <i>haastii</i> J.A. Pettersson (CHR 201850)		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

*Continued on next page*

## Data Deficient

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
<b>BATS</b>			
<b>Taxonomically determinate</b>			
Greater short-tailed bat	<i>Mystacina robusta</i>		EX
<b>BIRDS</b>			
<b>Taxonomically indeterminate</b>			
New Zealand storm-petrel	<i>Oceanites maorianus</i>		DD
<b>BRYOPHYTES</b>			
<b>Taxonomically determinate</b>			
Liverwort	<i>Andrewsianthus confusus</i>		
Liverwort	<i>Cephalozia pachygyna</i>		
Liverwort	<i>Cephaloziella aenigmatica</i>		
Liverwort	<i>Cephaloziella exigua</i>		
Liverwort	<i>Cephaloziella grandiretis</i>		
Liverwort	<i>Cephaloziella bispidissima</i>	S?O	
Liverwort	<i>Cephaloziella invisa</i>		
Liverwort	<i>Cephaloziella muelleriana</i>		
Liverwort	<i>Cephaloziella nothogena</i>		
Liverwort	<i>Cephaloziella pellucida</i>		
Liverwort	<i>Cephaloziella pseudocrassigyna</i>		
Liverwort	<i>Cephaloziella subspinosa</i>		
Liverwort	<i>Cheilolejeunea imbricata</i>	SO, OL	
Liverwort	<i>Chiloscyphus erosus</i> J.J. Engel	OL	
Liverwort	<i>Chiloscyphus battortii</i> J.J. Engel		
Liverwort	<i>Cryptostipula inundata</i>	OL	
Liverwort	<i>Eoisotachis nigella</i>	SO	
Moss	<i>Fissidens oblongifolius</i> var. <i>oblongifolius</i>	SO	
Liverwort	<i>Gymnomitrion strictum</i> var. <i>inequalis</i>		
Liverwort	<i>Haplomitrium minutum</i>		
Moss	<i>Hymenostylium recurvirostre</i> (Hedw.) Dixon	SO, OL	
Liverwort	<i>Lepidozia fugax</i>		
Liverwort	<i>Paracromastigum fissifolium</i>		
Liverwort	<i>Paracromastigum kirkii</i>		
Liverwort	<i>Radula parviretis</i>	OL	
Liverwort	<i>Rectolejeunea denudata</i>	OL	
Liverwort	<i>Riccia fluitans</i>	SO	
Liverwort	<i>Stolonivector waipouensis</i>		
Moss	<i>Trichostomum imshaugii</i> (Vitt) R.H. Zander		
Moss	<i>Willia calobolax</i> (Mull. Hal.) Lightowers	S?O	
<b>Taxonomically indeterminate</b>			
Liverwort	<i>Acromastigum aff. tenax</i>	OL	
Liverwort	<i>Aneura aff. rodwayi</i>	OL	
Liverwort	<i>Chiloscyphus beckettianus</i>		
Liverwort	<i>Cololejeunea</i> sp. 2		
Moss	<i>Dicranella temperata</i>		
Liverwort	<i>Drepanolejeunea aff. pentadactyla</i>	SO, OL	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Liverwort	<i>Drepanolejeunea</i> aff. <i>ternatensis</i>		
Liverwort	<i>Frullania rostrata</i> var. "draco"	OL	
Liverwort	<i>Hyalolepidozia</i> "microphylla"		
Liverwort	<i>Lejeunea</i> AK253889		
Liverwort	<i>Lejeunea</i> sp. 768	OL	
Liverwort	<i>Lejeunea</i> sp. 799	OL	
Liverwort	<i>Nephelolejeunea</i> aff. <i>bamata</i>	OL	
Liverwort	<i>Radula</i> "big Dave"		
Liverwort	<i>Radula</i> aff. <i>retroflexa</i>		
Liverwort	<i>Radula</i> sp.	OL	
Liverwort	<i>Scapania</i> cf. <i>nemorosa</i>		
Liverwort	<i>Scapania</i> cf. <i>undulata</i>		
Liverwort	<i>Telaranea</i> aff. <i>pseudozoopsis</i>	OL	
Liverwort	<i>Treubia</i> aff. <i>tasmanica</i>		
Liverwort	<i>Triandrophylloides symmetricum</i>		
<b>FRESHWATER FISH</b>			
<b>Taxonomically indeterminate</b>			
Possible new non-diadromous galaxias	<i>Galaxias</i> "Von sp."		HI
Lower Shag galaxias	<i>Galaxias</i> sp. E.		
Possible new non-migratory bully	<i>Gobiomorphus</i> sp. aff. <i>basalis</i>		
<b>FRESHWATER INVERTEBRATES</b>			
<b>Taxonomically determinate</b>			
Freshwater mussel	<i>Cucumerunio websteri delli</i> McMichael & Hiscock, 1958		
Freshwater mussel	<i>Cucumerunio websteri websteri</i> (Simpson, 1902)		
Mayfly	<i>Deleatidium (Penniketellum) insolitum</i>		
Mayfly	<i>Deleatidium magnum</i>		
Caddisfly	<i>Helicopsyche cuvieri</i> Johanson	OL	
Caddisfly	<i>Helicopsyche baurapango</i> Johanson		
Caddisfly	<i>Hydrochorema lyfordi</i> Ward, 1997		
Snail	<i>Latia climoi</i> Starobogatov, 1986		
Mayfly	<i>Mauiulus aquilus</i>		
Nematode	<i>Miconchus rex</i> (Cobb, 1904) Andrassy, 1958	OL	
Amphipod	<i>Neopbreatoicus assimilis</i>		
Amphipod	<i>Notamphisopus benbami</i>		
Amphipod	<i>Notamphisopus dunedinensis</i>		
Amphipod	<i>Notamphisopus flavius</i>		
Amphipod	<i>Notamphisopus kirktii</i>	OL	
Amphipod	<i>Notamphisopus littoralis</i>		
Amphipod	<i>Notamphisopus percevali</i>		
Net-winged midge	<i>Notoboraiia micrognathia</i> Craig, 1969	OL	
Caddisfly	<i>Paroxyethira hintoni</i> Leader, 1972		
Amphipod	<i>Pbreatoicus orarii</i>		
Snail	<i>Potamopyrgus pupoides</i> Hutton, 1882		
Caddisfly	<i>Psilochorema foliobarpax</i> McFarlane, 1956		
Mayfly	<i>Siphlaenigma janae</i> Penniket, 19??		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

*Data Deficient—continued*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
<b>Taxonomically indeterminate</b>			
Caddisfly	<i>Edpercivalia</i> sp. G		
Caddisfly	<i>Paroxyethira</i> sp. B		
Caddisfly	<i>Paroxyethira</i> sp. H		
Caddisfly	<i>Paroxyethira</i> sp. W		
<b>FUNGI</b>			
The list of Data Deficient fungi has not changed in the last 3 years; for the full list see Hitchmough (comp. 2002).			
<b>MACROALGAE</b>			
<b>Taxonomically determinate</b>			
Green seaweed—Ulvales	<i>Acrochaete endostraca</i>		
Red seaweed—Acrochaetales	<i>Acrochaetium leptonemoides</i>	OL	
Red seaweed—Acrochaetales	<i>Acrochaetium neozealandicum</i>	OL	
Red seaweed—Rhodymeniales	<i>Cephalocystis furcellata</i>		
Green seaweed—Bryopsidales	<i>Codium perriniae</i>		
Green seaweed—Bryopsidales	<i>Codium platyclados</i>	T?O	
Brown seaweed—Cutleriales	<i>Cutleria mollis</i>		
Green seaweed—Ulvales	<i>Epicladia testarum</i>		
Red seaweed—Erythropeltidales	<i>Erythrotrichia bangtoides</i>	OL	
Red seaweed—Erythropeltidales	<i>Erythrotrichia bunterae</i>	OL	
Golden brown alga	<i>Giraudyopsis stellifera</i>	SO	
Brown seaweed—Ectocarpales	<i>Nemacystus novae-zelandiae</i>		
Green seaweed—Ulvales	<i>Ochlochaete bystrix</i>		
Green seaweed—Ulvales	<i>Ovillaria catenata</i>		
Red algae—Porphyridiales	<i>Porphyridium purpuream</i>	SO	
Green seaweed—Bryopsidales	<i>Pseudobryopsis planktonica</i>	OL	
Red seaweed—Halymeniales	<i>Sebdenia lindaueri</i>	OL	
Green seaweed—Ulvales	<i>Syncoryne reinkei</i>		
<b>Taxonomically indeterminate</b>			
Red seaweed—Bangiales	<i>Bangia</i> spp.		
Green seaweed—Bryopsidales	<i>Caulerpa fastigiata</i>		
Red seaweed—Rhodymeniales	<i>Champiocolax</i> sp.		
Green seaweed—Ulvales	<i>Entocladia</i> spp.		
Red seaweed—Bangiales	<i>Porphyra</i> spp.		
<b>MARINE FISH</b>			
<b>Taxonomically determinate</b>			
Richardson's skate	<i>Bathyraja richardsoni</i>	SO	
Harrison's dogfish	<i>Centrophorus harringtoni</i>	TO	
Largespine velvet dogfish	<i>Centroscymnus macracanthus</i>	T?O	
Dusky rattail	<i>Coelorinchus infuscus</i>		
Bluntnose weefish	<i>Cologrammus flavescens</i>		
Longsnout dogfish	<i>Deania quadrispinosum</i>	SO	
Smooth deepsea skate	<i>Notoraja asperula</i>		
Prickly deepsea skate	<i>Notoraja spinifera</i>		
Redband weever	<i>Parapercis binivirgata</i>	SO	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

*Continued on next page*

*Data Deficient—continued*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
McMillan's cat shark	<i>Parmaturus macmillani</i>	S?O	
Fourspine boxfish	<i>Polyplacapros tyleri</i>	SO	
Wreckfish	<i>Polyprion americanus</i>		DD
Crocodile shark	<i>Pseudocarcharias kamoharai</i>	SO	
False cat shark	<i>Pseudotriakis microdon</i>	SO	
Little sleeper shark	<i>Somniosus rostratus</i>	SO	
Longsnout/smooth pipefish	<i>Stigmatopora macropterygia</i>		
Black pipefish	<i>Stigmatopora nigra</i>	SO	
<b>Taxonomically indeterminate</b>			
Fragile dogfish	<i>Centroscyllium ?kamoharai</i>		
Swell shark/banded carpetshark	<i>Cephaloscyllium</i> sp. B sensu Last & Stevens 1994		
Green-eyes	Chlorophthalmidae—sp.		
Pipefish	? <i>Cosmocampus</i> sp.		
Giant black ghostshark	<i>Hydrolagus</i> sp. D (giant black ghostshark)		
Codlings	<i>Laemonema</i> spp.		
Fangtooth hoki/bluebelly	<i>Lyconus</i> spp.	SO	
Triangle batfishes	<i>Malbopsis</i> spp.		
Deepsea skate	<i>Notoraja</i> sp. A		
Deepsea skate	<i>Notoraja</i> sp. B		
Deepsea skate	<i>Notoraja</i> sp. C		
Deepsea skate	<i>Notoraja</i> sp. D		
Deepsea skate	<i>Notoraja</i> sp. E		
Rough-backed cat shark	<i>Parmaturus</i> sp.		
Knifetooth dogfish	<i>Scymnodon</i> cf. <i>ringens</i>	SO	
Greeneeye spiny dogfish	<i>Squalus</i> cf. <i>mitsukurii</i>		
Longnose spiny dogfish	<i>Squalus</i> sp. 2		
Paddletail spiny dogfish	<i>Squalus</i> sp. 3		
Kermadec spiny dogfish	<i>Squalus</i> sp. 4		
Giant spiny dogfish	<i>Squalus</i> sp. 5		
<b>MARINE INVERTEBRATES</b>			
<b>Taxonomically determinate</b>			
Black coral	<i>Antipathes fruticosa</i> Gray, 1857		
Polychaete worm	<i>Hartmanonuphis pectinata</i>		
Polychaete worm	<i>Scoletoma lynnei</i>		
<b>Taxonomically indeterminate</b>			
Monoplacophoran limpet	<i>Micropilina</i> sp. A (NMNZ M.171278)	OL	
Monoplacophoran limpet	<i>Micropilina</i> sp. B (NMNZ M.171273)	OL	
Monoplacophoran limpet	<i>Micropilina</i> sp. C (NMNZ M.171275)	OL	
Monoplacophoran limpet	<i>Rokopella</i> sp. (NMNZ M.171277)	OL	
Monoplacophoran limpet	<i>Vema</i> sp. (NMNZ M.171272)	OL	
<b>MARINE MAMMALS</b>			
<b>Taxonomically determinate</b>			
Arnoux's beaked whale	<i>Berardius arnouxi</i>		LR/cd
Pygmy right whale	<i>Caperea marginata</i>		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

*Continued on next page*

*Data Deficient—continued*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Southern bottlenose whale	<i>Hyperoodon planifrons</i>		LR/cd
Pygmy sperm whale	<i>Kogia breviceps</i>	SO	
Andrews' beaked whale	<i>Mesoplodon bowdoini</i>		DD
Dense-beaked whale	<i>Mesoplodon densirostris</i>		DD
Gray's beaked whale	<i>Mesoplodon grayi</i>		DD
Hector's beaked whale	<i>Mesoplodon Hectori</i>		DD
Strap-toothed whale	<i>Mesoplodon layardii</i>		DD
Pygmy beaked whale	<i>Mesoplodon peruvianus</i>		DD
Spade-toothed whale	<i>Mesoplodon traversii</i>		
Shepherd's beaked whale	<i>Tasmacetus shepherdi</i>		DD
Goose-beaked whale	<i>Ziphius cavirostris</i>		DD
<b>Taxonomically indeterminate</b>			
Bottlenose dolphin "offshore"	<i>Tursiops truncatus</i> "offshore"	SO	
<b>REPTILES</b>			
<b>Taxonomically determinate</b>			
Takitimus gecko	<i>Hoplodactylus cryptozoicus</i>		
Striped skink	<i>Oligosoma striatum</i>		VU B1+2bcd
<b>Taxonomically indeterminate</b>			
Cascades forest gecko	<i>Hoplodactylus</i> "Cascades"		
Cupola Basin gecko	<i>Hoplodactylus</i> "Cupola"		
Okarito forest gecko	<i>Hoplodactylus</i> "Okarito forest gecko"	OL	
Roys Peak gecko	<i>Hoplodactylus</i> "Roys Peak"		
Rangitata skink	<i>Oligosoma</i> "Rangitata"	OL	
Sinbad skink	<i>Oligosoma</i> "Sinbad Valley"	OL	
Te Kakahu skink	<i>Oligosoma</i> "Te Kakahu"	OL	
Whirinaki skink	<i>Oligosoma</i> "Whirinaki"	OL	
<b>TERRESTRIAL INVERTEBRATES</b>			
<b>Taxonomically determinate</b>			
Earthworm	<i>Acanthodrilus kermadecensis</i>	OL	
Nematode	<i>Acrobeloides ellesmerensis</i> Yeates, 1967	OL	
Spider—Agelenidae	<i>Abua dentata</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Abua kaituna</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Abua vulgaris</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Akatorea gracilis</i> (Marples, 1959)		
Spider—Amphinectidae	<i>Akatorea otagoensis</i> Forster & Wilton, 1973		
Nematode	<i>Alaimus bimatangiensis</i> Yeates, 1967	OL	
Spider—Hahniidae	<i>Alistra centralis</i> (Forster, 1970)		
Spider—Hahniidae	<i>Alistra tnanga</i> (Forster, 1970)	OL	
Spider—Hahniidae	<i>Alistra mangareia</i> (Forster, 1970)	OL	
Spider—Hahniidae	<i>Alistra napua</i> (Forster, 1970)	OL	
Spider—Hahniidae	<i>Alistra opina</i> (Forster, 1970)	OL	
Spider—Hahniidae	<i>Alistra reinga</i> (Forster, 1970)		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

*Continued on next page*

Data Deficient—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Spider—Hahniidae	<i>Alistra tuna</i> (Forster, 1970)	OL	
Spider—Anyphaenidae	<i>Amaurobioides major</i> Forster, 1970	OL	
Spider—Anyphaenidae	<i>Amaurobioides maritimus</i> O.P.-Cambridge, 1883		
Spider—Anyphaenidae	<i>Amaurobioides minor</i> Forster, 1970		
Spider—Anyphaenidae	<i>Amaurobioides pallidus</i> Forster, 1970		
Spider—Anyphaenidae	<i>Amaurobioides picinus</i> Forster, 1970		
Spider—Anyphaenidae	<i>Amaurobioides piscator</i> Hogg, 1909		
Spider—Anyphaenidae	<i>Amaurobioides pletus</i> Forster, 1970		
Spider—Anyphaenidae	<i>Amaurobioides pocharus</i> Forster, 1970		
Spider—Amphinectidae	<i>Amphinecta decemmaculata</i> Simon, 1898		
Spider—Amphinectidae	<i>Amphinecta dejecta</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Amphinecta luta</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Amphinecta mara</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Amphinecta mula</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Amphinecta pika</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Amphinecta pila</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Amphinecta puka</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Amphinecta tama</i> Forster & Wilton, 1973	OL	
Spider—Amphinectidae	<i>Amphinecta tula</i> Forster & Wilton, 1973		
Stiletto fly	<i>Anabarhynchus albipennis</i> Lyneborg, 1992		
Stiletto fly	<i>Anabarhynchus atratus</i> Lyneborg, 1992		
Stiletto fly	<i>Anabarhynchus embersoni</i> Lyneborg, 1992		
Stiletto fly	<i>Anabarhynchus flaviventris</i> Lyneborg, 1992		
Stiletto fly	<i>Anabarhynchus fluviatilis</i> Lyneborg, 1992		
Stiletto fly	<i>Anabarhynchus budsoni</i> Lyneborg, 1992		
Stiletto fly	<i>Anabarhynchus indistinctus</i> Lyneborg, 1992		
Stiletto fly	<i>Anabarhynchus olivaceus</i> Lyneborg, 1992		
Stiletto fly	<i>Anabarhynchus simplex</i> Lyneborg, 1992		
Stiletto fly	<i>Anabarhynchus waitarerensis</i> Lyneborg, 1992		
Stiletto fly	<i>Anabarhynchus wisei</i> Lyneborg, 1992		
Spider—Cycloctenidae	<i>Anaua unica</i> Forster, 1970	OL	
Spider—Orsolobidae	<i>Anopsolobus subterraneus</i> Forster & Platnick, 1985	OL	
Spider—Lycosidae	<i>Anoteropsis ballae</i> Vink, 2002		
Spider—Amphinectidae	<i>Aorangia agama</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Aorangia fiordensis</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Aorangia isolata</i> Forster & Wilton, 1973	OL	
Spider—Amphinectidae	<i>Aorangia kapitiensis</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Aorangia mauii</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Aorangia muscicola</i> Forster & Wilton, 1973	OL	
Spider—Amphinectidae	<i>Aorangia obscura</i> Forster & Wilton, 1973	OL	
Spider—Amphinectidae	<i>Aorangia otria</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Aorangia pilgrimi</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Aorangia poppelwelli</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Aorangia pudica</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Aorangia semita</i> Forster & Wilton, 1973	OL	
Spider—Amphinectidae	<i>Aorangia silvestris</i> Forster & Wilton, 1973	OL	
Spider—Amphinectidae	<i>Aorangia singularis</i> Forster & Wilton, 1973		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Data Deficient—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Spider—Amphinectidae	<i>Aorangia tumida</i> Forster & Wilton, 1973		
Spider—Mecysmauchenidae	<i>Aotearoa magna</i> (Forster, 1949)		
Aphid	<i>Aphis nelsonensis</i> Cottier		
Nematode	<i>Aporcelaimellus taylori</i> Yeates, 1967		OL
Spider—Dictynidae	<i>Arangina cornigera</i> (Dalmas, 1917)		
Spider—Dictynidae	<i>Arangina pluva</i> Forster, 1970		
Moth	<i>Archyala lindsayi</i> (Philpott, 1927)		
Spider—Theridiidae	<i>Argyrodes lepida</i> O.P.-Cambridge, 1879		
Spider—Theridiidae	<i>Argyrodes triangulatus</i> (Urquhart, 1887)		
Spider—Theridiidae	<i>Argyrodes urquhartii</i> (Bryant, 1933)		
Spider—Segestriidae	<i>Ariadna bellatoria</i> Dalmas, 1917		OL
Flatworm	<i>Artburdendyus albidus</i> Jones & Gerard, 1999		
Darkling beetle	<i>Artystona richmondiana</i> Watt, 1992		OL
Nematode	<i>Ascaris apterycis</i> Chatin, 1884		
Spider—Orsolobidae	<i>Ascuta cantuaria</i> Forster & Platnick, 1985		OL
Spider—Orsolobidae	<i>Ascuta inopinata</i> Forster, 1956		
Spider—Orsolobidae	<i>Ascuta leith</i> Forster & Platnick, 1985		
Spider—Orsolobidae	<i>Ascuta media</i> Forster, 1956		
Spider—Orsolobidae	<i>Ascuta monowai</i> Forster & Platnick, 1985		OL
Spider—Orsolobidae	<i>Ascuta montana</i> Forster & Platnick, 1985		OL
Spider—Orsolobidae	<i>Ascuta musca</i> Forster & Platnick, 1985		
Spider—Orsolobidae	<i>Ascuta ornata</i> Forster, 1956		
Spider—Orsolobidae	<i>Ascuta parornata</i> Forster & Platnick, 1985		
Spider—Orsolobidae	<i>Ascuta taupo</i> Forster & Platnick, 1985		OL
Spider—Orsolobidae	<i>Ascuta tongariro</i> Forster & Platnick, 1985		
Spider—Orsolobidae	<i>Ascuta univa</i> Forster & Platnick, 1985		OL
Spider—Salticidae	<i>Attus albibreviatus</i> Walckenaer, 1837		
Spider—Salticidae	<i>Attus aquilis</i> Urquhart, 1886		
Spider—Salticidae	<i>Attus cookii</i> Walckenaer, 1837		
Spider—Salticidae	<i>Attus monticulus</i> Urquhart, 1891		
Spider—Salticidae	<i>Attus montinus</i> Urquhart, 1891		
Spider—Salticidae	<i>Attus pullus</i> Urquhart, 1890		
Spider—Salticidae	<i>Attus scindus</i> Urquhart, 1890		
Spider—Salticidae	<i>Attus subfuscus</i> Urquhart, 1887		
Spider—Salticidae	<i>Attus tenebrosus</i> Urquhart, 1893		
Spider—Salticidae	<i>Attus valentulus</i> Urquhart, 1891		
Spider—Amaurobiidae	<i>Aubunga pectinata</i> Forster & Wilton, 1973		
Moth	<i>Austrocidaria liturga</i> (Meyrick, 1911)		
Moth	<i>Bascantis sirenica</i> Meyrick, 1914		
Spider—Orsolobidae	<i>Bealeyia unicolor</i> Forster & Platnick, 1985		OL
Spider—Salticidae	<i>Bianor compactus</i> (Urquhart, 1885)		
Beetle	<i>Brounia thboracica</i>		
Spider—Stiphidiidae	<i>Cambridgea agrestis</i> Forster & Wilton, 1973		
Spider—Stiphidiidae	<i>Cambridgea decorata</i> Blest & Vink, 2000		
Spider—Stiphidiidae	<i>Cambridgea elegans</i> Blest & Vink, 2000		OL
Spider—Stiphidiidae	<i>Cambridgea elongata</i> Blest & Vink, 2000		
Spider—Stiphidiidae	<i>Cambridgea inaequalis</i> Blest & Vink, 2000		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Data Deficient—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Spider—Stiphidiidae	<i>Cambridgea insulana</i> Blest & Vink, 2000	OL	
Spider—Stiphidiidae	<i>Cambridgea longipes</i> Blest & Vink, 2000		
Spider—Stiphidiidae	<i>Cambridgea mercurialis</i> Blest & Vink, 2000	OL	
Spider—Stiphidiidae	<i>Cambridgea obscura</i> Blest & Vink, 2000		
Spider—Stiphidiidae	<i>Cambridgea occidentalis</i> Forster & Wilton, 1973	OL	
Spider—Stiphidiidae	<i>Cambridgea ordishi</i> Blest & Vink, 2000	OL	
Spider—Stiphidiidae	<i>Cambridgea pallidula</i> Blest & Vink, 2000		
Spider—Stiphidiidae	<i>Cambridgea peculiaris</i> Forster & Wilton, 1973	OL	
Spider—Stiphidiidae	<i>Cambridgea peelensis</i> Blest & Vink, 2000		
Spider—Stiphidiidae	<i>Cambridgea plagiata</i> Forster & Wilton, 1973		
Spider—Stiphidiidae	<i>Cambridgea quadromaculata</i> Blest & Taylor, 1995		
Spider—Stiphidiidae	<i>Cambridgea ramsayi</i> Forster & Wilton, 1973		
Spider—Stiphidiidae	<i>Cambridgea reinga</i> Forster & Wilton, 1973	OL	
Spider—Stiphidiidae	<i>Cambridgea secunda</i> Forster & Wilton, 1973		
Spider—Stiphidiidae	<i>Cambridgea solanderensis</i> Blest & Vink, 2000	OL	
Spider—Stiphidiidae	<i>Cambridgea sylvatica</i> Forster & Wilton, 1973	OL	
Spider—Stiphidiidae	<i>Cambridgea tuiiae</i> Blest & Vink, 2000	OL	
Spider—Stiphidiidae	<i>Cambridgea turbotti</i> Forster & Wilton, 1973	OL	
Moth	<i>Cateristis eustyla</i> Meyrick, 1889	TO, OL	
Spider—Araneidae	<i>Celaenia atkinsoni</i> (O.P.-Cambridge, 1879)	SO	
Spider—Araneidae	<i>Celaenia bectori</i> (O.P.-Cambridge, 1879)		
Spider—Araneidae	<i>Celaenia olivacea</i> (Urquhart, 1885)		
Spider—Araneidae	<i>Celaenia penna</i> (Urquhart, 1886)		
Spider—Araneidae	<i>Celaenia tuberosa</i> (Urquhart, 1887)		
Darkling beetle	<i>Cerodolus tuberculatus</i> Broun, 1917		
Snail	<i>Charopa macgillivrayana</i> Iredale, 1913	OL	
Fly	<i>Chersodromia zealandica</i>	OL	
Mirid bug	<i>Chinamiris niculatus</i> Eyles & Carvalho, 1991		
Spider—Clubionidae	<i>Clubiona blesti</i> Forster, 1979		
Spider—Clubionidae	<i>Clubiona producta</i> Forster, 1979		
Spider—Clubionidae	<i>Clubiona scatula</i> Forster, 1979		
Spider—Clubionidae	<i>Clubiona torta</i> Forster, 1979	OL	
Spider—Salticidae	<i>Clynotis barresi</i> Hogg, 1909		
Spider—Salticidae	<i>Clynotis knoxi</i> Forster, 1964		
Spider—Salticidae	<i>Clynotis saxatilis</i> (Urquhart, 1886)		
Nematode	<i>Cobbonchus chauloidus</i> Clark, 1961	OL	
Spider—Araneidae	<i>Colaranea brunnea</i> Court & Forster, 1988		
Spider—Araneidae	<i>Colaranea melanoviridis</i> Court & Forster, 1988		
Spider—Salticidae	<i>Cosmophasis archeyi</i> Berland, 1931		
Nematode	<i>Cryptbodera coxi</i> (Wouts, 1973) Luc, Taylor & Cadet, 1978	OL	
Nematode	<i>Cryptbodera nothofagi</i> (Wouts, 1973) Luc, Taylor & Cadet, 1978	OL	
Spider—Araneidae	<i>Cryptaranea stewartensis</i> Court & Forster, 1988		
Spider—Araneidae	<i>Cryptaranea subalpina</i> Court & Forster, 1988		
Spider—Cycloctenidae	<i>Cycloctenus agilis</i> Forster, 1979		
Spider—Cycloctenidae	<i>Cycloctenus duplex</i> Forster, 1979		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Data Deficient—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Spider—Cycloctenidae	<i>Cycloctenus fiordensis</i> Forster, 1979		
Spider—Cycloctenidae	<i>Cycloctenus fugax</i> Goyen, 1890		
Spider—Cycloctenidae	<i>Cycloctenus lepidus</i> Urquhart, 1890		
Spider—Cycloctenidae	<i>Cycloctenus nelsonensis</i> Forster, 1979		
Spider—Cycloctenidae	<i>Cycloctenus patura</i> Forster, 1979		
Spider—Cycloctenidae	<i>Cycloctenus pulcher</i> Urquhart, 1891		
Spider—Thomisidae	<i>Cymbachina albobrunnea</i> (Urquhart, 1893)		
Nematode	<i>Cyrnea (Cyrnea) apterycis</i> Harris, 1975		
Earthworm	<i>Decachaetus minor</i>	OL	
Earthworm	<i>Decachaetus violaceus</i>	OL	
Earthworm	<i>Deinodrilus agilis</i>	OL	
Earthworm	<i>Deinodrilus benhami</i>		
Earthworm	<i>Deinodrilus gracilis</i>		
Earthworm	<i>Deinodrilus lateralis</i>		
Earthworm	<i>Deinodrilus parvus</i>	OL	
Earthworm	<i>Deinodrilus suteri</i>		
Spider—Thomisidae	<i>Diaeal alboltimbata</i> L. Koch, 1875		
Spider—Thomisidae	<i>Diae a ambara</i> (Urquhart, 1885)		
Spider—Thomisidae	<i>Diae a spherooides</i> (Urquhart, 1885)		
Spider—Dictynidae	<i>Dictyna urquharti</i> Roewer, 1951		
Spider—Linyphiidae	<i>Diploplecta adjacens</i> Millidge, 1988		
Spider—Linyphiidae	<i>Diploplecta duplex</i> Millidge, 1988		
Spider—Linyphiidae	<i>Diploplecta nuda</i> Millidge, 1988	OL	
Spider—Linyphiidae	<i>Diploplecta opaca</i> Millidge, 1988		
Spider—Linyphiidae	<i>Diploplecta proxima</i> Millidge, 1988		
Spider—Linyphiidae	<i>Diploplecta simplex</i> Millidge, 1988		
Earthworm	<i>Diporochaeta aquatica</i>		
Earthworm	<i>Diporochaeta brachysoma</i>	OL	
Earthworm	<i>Diporochaeta caswelli</i>	OL	
Earthworm	<i>Diporochaeta chatbamensis</i>	OL	
Earthworm	<i>Diporochaeta duodecimalis</i>	OL	
Earthworm	<i>Diporochaeta heterochaeta</i>	OL	
Earthworm	<i>Diporochaeta intermedia</i>		
Earthworm	<i>Diporochaeta minima</i>	OL	
Earthworm	<i>Diporochaeta obtusa</i>		
Earthworm	<i>Diporochaeta punctata</i>		
Spider—Linyphiidae	<i>Drapetisca australis</i> Forster, 1955	OL	
Spider—Orsolobidae	<i>Dugdalea oculata</i> Forster & Platnick, 1985		
Spider—Linyphiidae	<i>Dunedinia decolor</i> Millidge, 1988		
Spider—Linyphiidae	<i>Dunedinia pullata</i> Millidge, 1988		
Spider—Amphinectidae	<i>Dunstania angustiae</i> (Marples, 1959)	OL	
Spider—Amphinectidae	<i>Dunstania besperis</i> Forster & Wilton, 1973	OL	
Spider—Amphinectidae	<i>Dunstania binawa</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Dunstania bova</i> Forster & Wilton, 1973	OL	
Spider—Amphinectidae	<i>Dunstania kochi</i> Forster & Wilton, 1973	OL	
Spider—Amphinectidae	<i>Dunstania mira</i> Forster & Wilton, 1973	OL	
Spider—Amphinectidae	<i>Dunstania montana</i> Forster & Wilton, 1973	OL	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Data Deficient—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Spider—Amphinectidae	<i>Dunstania nuntia</i> (Marples, 1959)	OL	
Spider—Amphinectidae	<i>Dunstania salmoni</i> Forster & Wilton, 1973	OL	
Spider—Orsolobidae	<i>Duripelta alta</i> Forster & Platnick, 1985	OL	
Spider—Orsolobidae	<i>Duripelta australis</i> Forster, 1956	OL, DP	
Spider—Orsolobidae	<i>Duripelta egmont</i> Forster & Platnick, 1985		
Spider—Orsolobidae	<i>Duripelta bunua</i> Forster & Platnick, 1985	OL	
Spider—Orsolobidae	<i>Duripelta koomaa</i> Forster & Platnick, 1985	OL	
Spider—Orsolobidae	<i>Duripelta mawhero</i> Forster & Platnick, 1985		
Spider—Orsolobidae	<i>Duripelta minutu</i> Forster, 1956		
Spider—Orsolobidae	<i>Duripelta monowai</i> Forster & Platnick, 1985	OL	
Spider—Orsolobidae	<i>Duripelta otara</i> Forster & Platnick, 1985	OL	
Spider—Orsolobidae	<i>Duripelta pallida</i> (Forster, 1956)		
Spider—Orsolobidae	<i>Duripelta paringa</i> Forster & Platnick, 1985		
Spider—Orsolobidae	<i>Duripelta peha</i> Forster & Platnick, 1985	OL	
Spider—Orsolobidae	<i>Duripelta scuta</i> Forster & Platnick, 1985		
Spider—Orsolobidae	<i>Duripelta totara</i> Forster & Platnick, 1985	OL	
Spider—Orsolobidae	<i>Duripelta townsendi</i> Forster & Platnick, 1985		
Spider—Orsolobidae	<i>Duripelta wattii</i> Forster & Platnick, 1985		
Nematode	<i>Echinuria australis</i> Clark, 1979		
Stiletto fly	<i>Ectinorhynchus furcatus</i> Lyneborg, 1992	OL	
Moth	<i>Elachista belonoma</i> (Meyrick, 1889)		
Moth	<i>Elachista melanura</i> Meyrick, 1889		
Assassin bug	<i>Empicoris aculeatus</i> (Berghroth, 1927)	OL	
Assassin bug	<i>Empicoris angulipennis</i> (Berghroth, 1927)	OL	
Earthworm	<i>Eodrilus annectens</i>	HI, OL	
Earthworm	<i>Eodrilus fallax</i>	OL	
Earthworm	<i>Eodrilus haplocystis</i>	OL	
Earthworm	<i>Eodrilus micros</i>	OL	
Earthworm	<i>Eodrilus montanus</i>		
Earthworm	<i>Eodrilus pallidus</i>		
Earthworm	<i>Eodrilus paludosus</i>		
Earthworm	<i>Eodrilus parvus</i>	OL	
Earthworm	<i>Eodrilus rossi</i>	OL	
Moth	<i>Epichorista lindsayi</i> Philpott, 1928		
Spider—Theridiidae	<i>Episinus antipodianus</i> O.P.-Cambridge, 1879		
Spider—Theridiidae	<i>Episinus similanus</i> Urquhart, 1893		
Spider—Theridiidae	<i>Episinus similitudus</i> Urquhart, 1893		
Spider—Tetragnathidae	<i>Eryciniolia purpurapunctata</i> (Urquhart, 1890)		
Beetle	<i>Euconnus microciliipes</i> Franz, 1985		
Beetle	<i>Euconnus paraciliipes</i> Franz, 1985		
Earthworm	<i>Eudinodriloides forsteri</i>		
Moth	<i>Eudonia ustiramis</i> (Meyrick, 1931)	OL	
Darkling beetle	<i>Exadellium rufilabrum</i> (Broun, 1886)		
Snail	<i>Fanulum expositum</i> (Mousson, 1873)		
Spider—Zodariidae	<i>Forsterella faceta</i> Jocqué, 1991		
Spider—Nicodamidae	<i>Forstertyna marplesi</i> (Forster, 1970)		
Spider—Desidae	<i>Gasparia busa</i> Forster, 1970	OL	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Data Deficient—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Spider—Desidae	<i>Gasparia coriacea</i> Forster, 1970		
Spider—Desidae	<i>Gasparia delli</i> (Forster, 1955)	OL	
Spider—Desidae	<i>Gasparia dentata</i> Forster, 1970	OL	
Spider—Desidae	<i>Gasparia edwardsi</i> Forster, 1970	OL	
Spider—Desidae	<i>Gasparia katangaroa</i> Forster, 1970	OL	
Spider—Desidae	<i>Gasparia littoralis</i> Forster, 1970		
Spider—Desidae	<i>Gasparia lomasi</i> Forster, 1970	OL	
Spider—Desidae	<i>Gasparia mangamuka</i> Forster, 1970	OL	
Spider—Desidae	<i>Gasparia montana</i> Forster, 1970	OL	
Spider—Desidae	<i>Gasparia nava</i> Forster, 1970	OL	
Spider—Desidae	<i>Gasparia nebulosa</i> Marples, 1956	OL	
Spider—Desidae	<i>Gasparia nelsonensis</i> Forster, 1970		
Spider—Desidae	<i>Gasparia nuntia</i> Forster, 1970		
Spider—Desidae	<i>Gasparia oparara</i> Forster, 1970	OL	
Spider—Desidae	<i>Gasparia parva</i> Forster, 1970	OL	
Spider—Desidae	<i>Gasparia pluta</i> Forster, 1970	OL	
Spider—Desidae	<i>Gasparia rupicola</i> Forster, 1970	OL	
Spider—Desidae	<i>Gasparia rustica</i> Forster, 1970		
Spider—Desidae	<i>Gasparia tepakia</i> Forster, 1970		
Spider—Desidae	<i>Gasparia tuaiensis</i> Forster, 1970		
Nematode	<i>Globodera zealandica</i> Wouts, 1984	OL?	
Spider—Desidae	<i>Gobia isolata</i> Forster, 1970		
Spider—Desidae	<i>Gobia parisolata</i> Forster, 1970	OL	
Spider—Desidae	<i>Goyenia electa</i> Forster, 1970		
Spider—Desidae	<i>Goyenia fresa</i> Forster, 1970		
Spider—Desidae	<i>Goyenia gratiosa</i> Forster, 1970		
Spider—Desidae	<i>Goyenia lucrosa</i> Forster, 1970		
Spider—Desidae	<i>Goyenia marplesi</i> Forster, 1970	OL	
Spider—Desidae	<i>Goyenia multidentata</i> Forster, 1970	OL	
Spider—Desidae	<i>Goyenia ornata</i> Forster, 1970	OL	
Spider—Desidae	<i>Goyenia sana</i> Forster, 1970	OL	
Spider—Desidae	<i>Goyenia scitula</i> Forster, 1970	OL	
Spider—Desidae	<i>Goyenia sylvatica</i> Forster, 1970		
Moth	<i>Graphania omicron</i> (Hudson, 1898)	OL	
Moth	<i>Gymnobathra ambigua</i> (Philpott, 1926)		
Moth	<i>Gymnobathra origenes</i> Meyrick, 1936		
Moth	“ <i>Gymnobathra</i> ” <i>rufopunctella</i> Hudson, 1950		
Cranefly	<i>Gynoplistia ambulator</i> Alexander, 1924		
Cranefly	<i>Gynoplistia beimalis</i> (Alexander, 1924)		
Fly	<i>Halteripborus mirabilis</i> Parent, 1933	OL	
Spider—Linyphiidae	<i>Haplitis alticola</i> Blest & Vink, 2002	OL	
Spider—Linyphiidae	<i>Haplitis anomala</i> Blest & Vink, 2003	OL	
Spider—Linyphiidae	<i>Haplitis brevipes</i> (Blest, 1979)	OL, DP	
Spider—Linyphiidae	<i>Haplitis contorta</i> (Blest, 1979)	OL	
Spider—Linyphiidae	<i>Haplitis dunstani</i> (Blest, 1979)	OL	
Spider—Linyphiidae	<i>Haplitis exigua</i> Blest & Vink, 2002	OL	
Spider—Linyphiidae	<i>Haplitis fluviatilis</i> (Blest, 1979)		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Data Deficient—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Spider—Linyphiidae	<i>Haplitis fulvolineata</i> Blest & Vink, 2002	OL	
Spider—Linyphiidae	<i>Haplitis innotabilis</i> (Blest, 1979)		
Spider—Linyphiidae	<i>Haplitis insignis</i> (Blest, 1979)		
Spider—Linyphiidae	<i>Haplitis marplesi</i> Blest & Vink, 2003	OL	
Spider—Linyphiidae	<i>Haplitis moratnicola</i> Blest & Vink, 2002		
Spider—Linyphiidae	<i>Haplitis mundenia</i> (Urquhart, 1894)		
Spider—Linyphiidae	<i>Haplitis paradoxa</i> (Blest, 1979)		
Spider—Linyphiidae	<i>Haplitis silvicola</i> (Blest, 1979)		
Spider—Linyphiidae	<i>Haplitis similis</i> (Blest, 1979)		
Spider—Linyphiidae	<i>Haplitis subclathrata</i> Simon, 1894		
Spider—Linyphiidae	<i>Haplitis taranakii</i> (Blest, 1979)	OL	
Spider—Linyphiidae	<i>Haplitis titan</i> (Blest, 1979)		
Spider—Linyphiidae	<i>Haplitis tokaanuae</i> Blest & Vink, 2002	OL	
Spider—Desidae	<i>Hapona amira</i> Forster, 1970	OL	
Spider—Desidae	<i>Hapona aucklandensis</i> (Forster, 1964)	OL	
Spider—Desidae	<i>Hapona crypta</i> (Forster, 1964)	OL	
Spider—Desidae	<i>Hapona insula</i> (Forster, 1964)	OL	
Spider—Desidae	<i>Hapona marplesi</i> (Forster, 1964)	OL	
Spider—Desidae	<i>Hapona moana</i> Forster, 1970	OL	
Spider—Desidae	<i>Hapona momona</i> Forster, 1970	OL	
Spider—Desidae	<i>Hapona muscicola</i> (Forster, 1964)		
Spider—Desidae	<i>Hapona otagoa</i> (Forster, 1964)		
Spider—Desidae	<i>Hapona paibia</i> Forster, 1970	OL	
Spider—Desidae	<i>Hapona reinga</i> Forster, 1970	OL	
Spider—Desidae	<i>Hapona salmoni</i> (Forster, 1964)	OL	
Spider—Desidae	<i>Hapona tararua</i> Forster, 1970	OL	
Spider—Tengellidae	<i>Haurokoia filicicola</i> Forster & Wilton, 1973	OL	
Nematode	<i>Hedruris minuta</i> Andrews, 1974		
Nematode	<i>Helicotylenchus depressus</i> Yeates, 1967	OL	
Spider—Salticidae	<i>Helpis minitabunda</i> (L. Koch, 1880)	SO	
Spider—Desidae	<i>Helsonia plata</i> Forster, 1970	OL	
Ground weta	<i>Hemiandrus lanceolatus</i> (Walker, 1869)	OL	
Nematode	<i>Heterakis gracilicauda</i> Harris, 1975		
Nematode	<i>Heterodera aucklandica</i> Wouts & Sturhan, 1995	OL	
Spider—Hexathelidae	<i>Hexatbele cantuaria</i> Forster, 1968		
Spider—Hexathelidae	<i>Hexatbele cavernicola</i> Forster, 1968		
Spider—Hexathelidae	<i>Hexatbele exemplar</i> Parrott, 1960		
Spider—Hexathelidae	<i>Hexatbele buka</i> Forster, 1968		
Spider—Hexathelidae	<i>Hexatbele buttoni</i> Hogg, 1908		
Spider—Hexathelidae	<i>Hexatbele maitai</i> Forster, 1968	OL	
Spider—Hexathelidae	<i>Hexatbele nigra</i> Forster, 1968		
Spider—Hexathelidae	<i>Hexatbele otira</i> Forster, 1968		
Spider—Hexathelidae	<i>Hexatbele para</i> Forster, 1968		
Spider—Hexathelidae	<i>Hexatbele pukea</i> Forster, 1968	OL	
Spider—Hexathelidae	<i>Hexatbele putuna</i> Forster, 1968	OL	
Spider—Hexathelidae	<i>Hexatbele ramsayi</i> Forster, 1968	OL	
Spider—Hexathelidae	<i>Hexatbele rupicola</i> Forster, 1968		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Data Deficient—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Spider—Hexathelidae	<i>Hexatbele taumara</i> Forster, 1968	OL	
Spider—Hexathelidae	<i>Hexatbele waipa</i> Forster, 1968		
Spider—Hexathelidae	<i>Hexatbele waita</i> Forster, 1968	OL	
Spider—Hexathelidae	<i>Hexatbele wiltoni</i> Forster, 1968		
Spider—Salticidae	<i>Hinewata embolica</i> Zabka & Pollard, 2002		
Spider—Amphinectidae	<i>Holomamoea foveata</i> Forster & Wilton, 1973		
Spider—Salticidae	<i>Holoplatys appressus</i> Dalmas, 1917		
Moth	<i>Homoeosoma ischnomorpha</i> Meyrick, 1931		
Earthworm	<i>Hoplochaetina durvilleana</i>	OL	
Earthworm	<i>Hoplochaetina pallida</i>		
Earthworm	<i>Hoplochaetina polycystis</i>		
Earthworm	<i>Hoplochaetina robusta</i>	OL	
Earthworm	<i>Hoplochaetina rossii</i>		
Earthworm	<i>Hoplochaetina rubra</i>		
Earthworm	<i>Hoplochaetina spirilla</i>		
Earthworm	<i>Hoplochaetina subtilis</i>		
Spider—Amphinectidae	<i>Huara chapmanae</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Huara decorata</i> Forster & Wilton, 1973	OL	
Spider—Amphinectidae	<i>Huara dolosa</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Huara bastata</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Huara inflata</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Huara kikkawa</i> Forster & Wilton, 1973	OL	
Spider—Amphinectidae	<i>Huara marplesi</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Huara mura</i> Forster & Wilton, 1973	OL	
Spider—Amphinectidae	<i>Huara pudica</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Huka alba</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Huka lobata</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Huka minima</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Huka minutula</i> Forster & Wilton, 1973		
Spider—Desidae	<i>Hulua convoluta</i> Forster & Wilton, 1973		
Spider—Desidae	<i>Hulua manga</i> Forster & Wilton, 1973		
Spider—Desidae	<i>Hulua minima</i> Forster & Wilton, 1973		
Spider—Desidae	<i>Hulua pana</i> Forster & Wilton, 1973		
Spider—Linyphiidae	<i>Hyperafroneta obscura</i> Blest, 1979		
Spider—Salticidae	<i>Hypoblemum albovittatum</i> (Keyserling, 1882)	SO	
Spider—Gnaphosidae	<i>Hypodrassodes apicus</i> Forster, 1979	OL	
Spider—Gnaphosidae	<i>Hypodrassodes courti</i> Forster, 1979		
Spider—Gnaphosidae	<i>Hypodrassodes crassus</i> Forster, 1979		
Spider—Gnaphosidae	<i>Hypodrassodes insulanus</i> Forster, 1979	OL	
Spider—Gnaphosidae	<i>Hypodrassodes isopus</i> Forster, 1979	OL	
Spider—Theridiidae	<i>Icona alba</i> Forster, 1955		
Spider—Theridiidae	<i>Icona drama</i> Forster, 1964		
Nematode	<i>Idiodylaimus novaezealandiae</i> (Cobb, 1904)	OL	
	Andrássy, 1969		
Moth	<i>Izatba caustopa</i> Meyrick, 1892		
Spider—Salticidae	<i>Jotus rarus</i> (Urquhart, 1893)		
Spider—Gnaphosidae	<i>Kaitawa insulare</i> (Marples, 1956)	OL	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Data Deficient—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Spider—Hahniidae	<i>Kapanga alta</i> Forster, 1970	OL	
Spider—Hahniidae	<i>Kapanga festiva</i> Forster, 1970	OL	
Spider—Hahniidae	<i>Kapanga grana</i> Forster, 1970	OL	
Spider—Hahniidae	<i>Kapanga isulata</i> (Forster, 1970)	OL	
Spider—Hahniidae	<i>Kapanga luana</i> Forster, 1970	OL	
Spider—Hahniidae	<i>Kapanga mana</i> Forster, 1970	OL	
Spider—Hahniidae	<i>Kapanga manga</i> Forster, 1970	OL	
Spider—Hahniidae	<i>Kapanga solitaria</i> (Bryant, 1935)		
Spider—Hahniidae	<i>Kapanga wiltoni</i> Forster, 1970		
Spider—Oonopidae	<i>Kapitia obscura</i> Forster, 1956	OL	
Snail	<i>Kermarion kermadecensis</i> (E.A. Smith, 1873)	OL	
Snail	<i>Kermodon royanum</i> (Iredale, 1913)	OL	
Snail	<i>Kieconcha kermadeci</i> (Pfeiffer, 1856)	OL	
Spider—Desidae	<i>Laestrygones albiceres</i> Urquhart, 1894		
Spider—Desidae	<i>Laestrygones otagoensis</i> Forster, 1970		
Spider—Desidae	<i>Laestrygones westlandicus</i> Forster, 1970	OL	
Spider—Linyphiidae	<i>Laetesia amoena</i> Millidge, 1988		
Spider—Linyphiidae	<i>Laetesia bellissima</i> Millidge, 1988		
Spider—Linyphiidae	<i>Laetesia germana</i> Millidge, 1988		
Spider—Linyphiidae	<i>Laetesia intermedia</i> Blest & Vink, 2003		
Spider—Linyphiidae	<i>Laetesia minor</i> Millidge, 1988		
Spider—Linyphiidae	<i>Laetesia olvidada</i> Blest & Vink, 2003	OL	
Spider—Linyphiidae	<i>Laetesia paragermana</i> Blest & Vink, 2003	OL	
Spider—Linyphiidae	<i>Laetesia prominens</i> Millidge, 1988		
Spider—Linyphiidae	<i>Laetesia pseudamoena</i> Blest & Vink, 2003	OL	
Spider—Linyphiidae	<i>Laetesia pulcherrima</i> Blest & Vink, 2003	OL	
Spider—Linyphiidae	<i>Laetesia weburdi</i> (Urquhart, 1890)		
Spider—Desidae	<i>Lamina minor</i> Forster, 1970		
Spider—Desidae	<i>Lamina montana</i> Forster, 1970		
Spider—Desidae	<i>Lamina parana</i> Forster, 1970		
Spider—Desidae	<i>Lamina ulva</i> Forster, 1970	OL	
Moth	<i>Lathicrossa prophetica</i> Meyrick, 1927		
Spider—Salticidae	<i>Laufeta aeribirta</i> (Urquhart, 1888)		
Moth	“ <i>Leptocroca</i> ” <i>xyrias</i> Meyrick, 1931		
Earthworm	<i>Leucodrilus digitocystis</i>		
Earthworm	<i>Leucodrilus fuscus</i>	OL	
Earthworm	<i>Leucodrilus robustus</i>		
Nematode	<i>Longibulbophora ammophilae</i> Yeates, 1967	OL	
Nematode	<i>Longidorus orongorongensis</i> Yeates, van Etteger & Hooper, 1992	OL	
Spider—Agelenidae	<i>Mabura accola</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Mabura bainhamensis</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Mabura crypta</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Mabura detrita</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Mabura binua</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Mabura musca</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Mabura rubella</i> Forster & Wilton, 1973	OL	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Data Deficient—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Spider—Agelenidae	<i>Mabura rufula</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Mabura scuta</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Mabura spinosoides</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Mabura takabea</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Mabura tarsa</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Mabura vella</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Makora calypso</i> (Marples, 1959)		
Spider—Amphinectidae	<i>Makora detrita</i> Forster & Wilton, 1973	OL	
Spider—Amphinectidae	<i>Makora diversa</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Makora figurata</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Makora minica</i> Forster & Wilton, 1973		
Spider—Amaurobiidae	<i>Maloides cavernicola</i> (Forster & Wilton, 1973)	OL	
Spider—Amphinectidae	<i>Mamoea assimilis</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Mamoea bicolor</i> (Bryant, 1935)		
Spider—Amphinectidae	<i>Mamoea cantuaria</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Mamoea cooki</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Mamoea florae</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Mamoea hesperis</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Mamoea hughsoni</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Mamoea inornata</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Mamoea mandibularis</i> (Bryant, 1935)		
Spider—Amphinectidae	<i>Mamoea maorica</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Mamoea montana</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Mamoea monticola</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Mamoea otira</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Mamoea pilosa</i> (Bryant, 1935)		
Spider—Amphinectidae	<i>Mamoea rakirua</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Mamoea rufa</i> (Berland, 1931)		
Spider—Amphinectidae	<i>Mamoea unica</i> Forster & Wilton, 1973		
Spider—Desidae	<i>Manawa frondosa</i> Forster, 1970	OL	
Spider—Desidae	<i>Manawa setosa</i> Forster, 1970	OL	
Spider—Desidae	<i>Manawa solitaria</i> Forster, 1970		
Spider—Desidae	<i>Mangareia maculata</i> Forster, 1970	OL	
Spider—Desidae	<i>Mangareia motu</i> Forster, 1970	OL	
Spider—Synotaxidae	<i>Mangua caswell</i> Forster, 1990		
Spider—Synotaxidae	<i>Mangua convoluta</i> Forster, 1990		
Spider—Synotaxidae	<i>Mangua flora</i> Forster, 1990		
Spider—Synotaxidae	<i>Mangua gunni</i> Forster, 1990		
Spider—Synotaxidae	<i>Mangua hughsoni</i> Forster, 1990		
Spider—Synotaxidae	<i>Mangua kapiti</i> Forster, 1990	OL	
Spider—Synotaxidae	<i>Mangua makarora</i> Forster, 1990		
Spider—Synotaxidae	<i>Mangua medialis</i> Forster, 1990		
Spider—Synotaxidae	<i>Mangua oparara</i> Forster, 1990	OL	
Spider—Synotaxidae	<i>Mangua otira</i> Forster, 1990	OL	
Spider—Synotaxidae	<i>Mangua paringa</i> Forster, 1990		
Spider—Synotaxidae	<i>Mangua sana</i> Forster, 1990	OL	
Spider—Synotaxidae	<i>Mangua secunda</i> Forster, 1990	OL	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Data Deficient—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Spider—Amphinectidae	<i>Manibo australis</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Manibo cantuarius</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Manibo centralis</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Manibo insulanus</i> Forster & Wilton, 1973	OL	
Spider—Amphinectidae	<i>Manibo meridionalis</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Manibo ngaitabu</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Manibo otagoensis</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Manibo pumilio</i> Forster & Wilton, 1973	OL	
Spider—Amphinectidae	<i>Manibo tigris</i> Marples, 1959		
Spider—Amphinectidae	<i>Manibo vulgaris</i> Forster & Wilton, 1973		
Spider—Orsolobidae	<i>Maoriata magna</i> (Forster, 1956)		
Spider—Orsolobidae	<i>Maoriata montana</i> Forster & Platnick, 1985		
Earthworm	<i>Maoridrilus alpinus</i>	OL	
Earthworm	<i>Maoridrilus carnosus</i>		
Earthworm	<i>Maoridrilus dissimilis</i>		
Earthworm	<i>Maoridrilus fuscus</i>	OL	
Earthworm	<i>Maoridrilus gravus</i>		
Earthworm	<i>Maoridrilus megacystis</i>	OL	
Earthworm	<i>Maoridrilus michaelseni</i>	OL	
Earthworm	<i>Maoridrilus minor</i>	OL	
Earthworm	<i>Maoridrilus modestus</i>		
Earthworm	<i>Maoridrilus montanus</i>		
Earthworm	<i>Maoridrilus nelsoni</i>		
Earthworm	<i>Maoridrilus pallidus</i>		
Earthworm	<i>Maoridrilus parkeri</i>	OL	
Earthworm	<i>Maoridrilus plumbeus</i>		
Earthworm	<i>Maoridrilus purus</i>		
Earthworm	<i>Maoridrilus ruber</i>		
Earthworm	<i>Maoridrilus rubicundus</i>	OL	
Earthworm	<i>Maoridrilus smithi</i>		
Earthworm	<i>Maoridrilus suteri</i>	OL	
Earthworm	<i>Maoridrilus tetragonurus</i>	OL	
Earthworm	<i>Maoridrilus transalpinus</i>		
Earthworm	<i>Maoridrilus ultiginosus</i>		
Earthworm	<i>Maoridrilus ultimus</i>		
Earthworm	<i>Maoridrilus volutus</i>	OL	
Earthworm	<i>Maoridrilus wilkini</i>		
Spider—Linyphiidae	<i>Maorineta acerba</i> Millidge, 1988		
Spider—Linyphiidae	<i>Maorineta gentilis</i> Millidge, 1988		
Spider—Linyphiidae	<i>Maorineta sulcata</i> Millidge, 1988		
Spider—Linyphiidae	<i>Maorineta tibialis</i> Millidge, 1988		
Spider—Linyphiidae	<i>Maorineta tumida</i> Millidge, 1988		
Beetle	<i>Maorinus bunuaeformis</i> (Franz, 1985)		
Beetle	<i>Maorinus toronouti</i> (Franz, 1985)		
Spider—Salticidae	<i>Marpissa arenaria</i> Urquhart, 1888		
Spider—Salticidae	<i>Marpissa armifera</i> Urquhart, 1892		
Spider—Salticidae	<i>Marpissa cineracea</i> Urquhart, 1891		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Data Deficient—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Spider—Salticidae	<i>Marpissa leucophaina</i> Urquhart, 1888		
Spider—Salticidae	<i>Marpissa marina</i> (Goyen, 1892)		
Spider—Salticidae	<i>Marpissa nemoralis</i> Urquhart, 1892		
Spider—Amphinectidae	<i>Marplesia dugdalei</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Marplesia pobara</i> Forster & Wilton, 1973	OL	
Spider—Desidae	<i>Matachbia magna</i> Forster, 1970	OL	
Spider—Desidae	<i>Matachbia marplesi</i> Forster, 1970	OL	
Spider—Desidae	<i>Matachbia ramulicola</i> Dalmas, 1917		
Spider—Desidae	<i>Matachbia similis</i> Forster, 1970		
Spider—Gnaphosidae	<i>Matua festiva</i> Forster, 1979	OL	
Spider—Gnaphosidae	<i>Matua valida</i> Forster, 1979		
Fern weevil	<i>Megacolabus bifurcatus</i> May, 1973		
Fern weevil	<i>Megacolabus obesus</i> May, 1963	OL	
Spider—Linyphiidae	<i>Megafroneta dugdaleae</i> Blest & Vink, 2002	OL	
Spider—Linyphiidae	<i>Megafroneta elongata</i> Blest, 1979	OL	
Spider—Linyphiidae	<i>Megafroneta gigas</i> Blest, 1979	OL	
Earthworm	<i>Megascolex animae</i>	OL	
Earthworm	<i>Megascolex novae-zealandiae</i>		
Earthworm	<i>Megascolides albus</i>	OL	
Earthworm	<i>Megascolides equestris</i>	OL	
Earthworm	<i>Megascolides esculentus</i>	OL	
Earthworm	<i>Megascolides fuscus</i>		
Earthworm	<i>Megascolides buttoni</i>	OL	
Earthworm	<i>Megascolides irregularis</i>		
Earthworm	<i>Megascolides kirki</i>	OL	
Earthworm	<i>Megascolides majoricus</i>		
Earthworm	<i>Megascolides mortensenii</i>	OL	
Earthworm	<i>Megascolides napierensis</i>	OL	
Earthworm	<i>Megascolides neglectus</i>	OL	
Earthworm	<i>Megascolides parvus</i>		
Earthworm	<i>Megascolides raglani</i>		
Earthworm	<i>Megascolides reptans</i>		
Earthworm	<i>Megascolides ruber</i>		
Earthworm	<i>Megascolides rubicundus</i>	OL	
Earthworm	<i>Megascolides sapidus</i>	OL	
Earthworm	<i>Megascolides suteri</i>		
Earthworm	<i>Megascolides tasmani</i>	OL	
Earthworm	<i>Megascolides unipapillatus</i>	OL	
Earthworm	<i>Megascolides urewerae</i>		
Earthworm	<i>Megascolides viridis</i>		
Darkling beetle	<i>Menimus obscurus</i> Broun, 1880		
Darkling beetle	<i>Menimus puncticeps</i> Broun, 1880		
Darkling beetle	<i>Menimus thoracicus</i> Broun, 1880		
Spider—Synotaxidae	<i>Meringa borealis</i> Forster, 1990		
Spider—Synotaxidae	<i>Meringa centralis</i> Forster, 1990		
Spider—Synotaxidae	<i>Meringa conway</i> Forster, 1990	OL	
Spider—Synotaxidae	<i>Meringa leith</i> Forster, 1990		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

*Data Deficient—continued*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Spider—Synotaxidae	<i>Meringa nelsoni</i> Forster, 1990	OL	
Spider—Synotaxidae	<i>Meringa otago</i> Forster, 1990		
Spider—Synotaxidae	<i>Meringa tetragnyna</i> Forster, 1990	OL	
Nematode	<i>Mermis novaezealandiae</i> Cobb, 1904	OL	
Nematode	<i>Mesodorylaimus profundis</i> (Cobb, 1904) Goodey, 1963	OL	
Spider—Tetragnathidae	<i>Meta rufolineata</i> (Urquhart, 1889)		
Spider—Linyphiidae	<i>Metafroneta minima</i> Blest, 1979	OL	
Spider—Linyphiidae	<i>Metafroneta subversa</i> Blest & Vink, 2002		
Spider—Linyphiidae	<i>Metamynoglenes absurdus</i> Blest & Vink, 2002	OL	
Spider—Linyphiidae	<i>Metamynoglenes attenuata</i> Blest, 1979		
Spider—Linyphiidae	<i>Metamynoglenes incurvata</i> Blest, 1979	OL	
Nematode	<i>Miconchus kirikiri</i> Yeates, 1967	OL	
Fly	<i>Microsania tonnoiri</i> Collart, 1934	OL	
Earthworm	<i>Microscolex aucklandicus</i>		
Earthworm	<i>Microscolex campbellianus</i>		
Earthworm	<i>Microscolex phosphoreus</i>	OL	
Nematode	<i>Microtetrameres nestoris</i> Clark, Black & Rutherford, 1979		
Spider—Migidae	<i>Migas australis</i> Wilton, 1968		
Spider—Migidae	<i>Migas borealis</i> Wilton, 1968	OL	
Spider—Migidae	<i>Migas cambridgei</i> Wilton, 1968	OL	
Spider—Migidae	<i>Migas cantuarius</i> Wilton, 1968	OL	
Spider—Migidae	<i>Migas centralis</i> Wilton, 1968	OL	
Spider—Migidae	<i>Migas cumberi</i> Wilton, 1968	OL	
Spider—Migidae	<i>Migas distinctus</i> O.P.-Cambridge, 1879		
Spider—Migidae	<i>Migas gatenbyi</i> Wilton, 1968		
Spider—Migidae	<i>Migas giveni</i> Wilton, 1968		
Spider—Migidae	<i>Migas goyenii</i> Wilton, 1968	OL	
Spider—Migidae	<i>Migas hesperus</i> Wilton, 1968		
Spider—Migidae	<i>Migas hollowayi</i> Wilton, 1968	OL	
Spider—Migidae	<i>Migas kirki</i> Wilton, 1968		
Spider—Migidae	<i>Migas kochii</i> Wilton, 1968	OL	
Spider—Migidae	<i>Migas linburnensis</i> Wilton, 1968	OL	
Spider—Migidae	<i>Migas lomasi</i> Wilton, 1968	OL	
Spider—Migidae	<i>Migas marplesi</i> Wilton, 1968		
Spider—Migidae	<i>Migas minor</i> Wilton, 1968	OL	
Spider—Migidae	<i>Migas otari</i> Wilton, 1968	OL	
Spider—Migidae	<i>Migas paradoxus</i> L. Koch, 1873		
Spider—Migidae	<i>Migas quintus</i> Wilton, 1968	OL	
Spider—Migidae	<i>Migas saxtilis</i> Wilton, 1968	OL	
Spider—Migidae	<i>Migas secundus</i> Wilton, 1968	OL	
Spider—Migidae	<i>Migas solitarius</i> Wilton, 1968	OL	
Spider—Migidae	<i>Migas taterii</i> Todd, 1945	OL	
Spider—Migidae	<i>Migas tasmani</i> Wilton, 1968	OL	
Spider—Migidae	<i>Migas toddae</i> Wilton, 1968	OL	
Spider—Migidae	<i>Migas tuboe</i> Wilton, 1968	OL	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

*Continued on next page*

Data Deficient—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Spider—Mimetidae	<i>Mimetes mendicus</i> O.P.-Cambridge, 1879		
Spider—Mimetidae	<i>Mimetes sennio</i> (Urquhart, 1891)		
Darkling beetle	<i>Mimopeus neglectus</i> Watt, 1988		
Spider—Idiopidae	<i>Misgolas aperta</i> (Forster, 1968)		
Spider—Idiopidae	<i>Misgolas apica</i> (Forster, 1968)		
Spider—Idiopidae	<i>Misgolas assimilis</i> (Forster, 1968)		
Spider—Idiopidae	<i>Misgolas borealis</i> (Forster, 1968)	OL	
Spider—Idiopidae	<i>Misgolas cattinensis</i> (Forster, 1968)	OL	
Spider—Idiopidae	<i>Misgolas cognata</i> (Forster, 1968)	OL	
Spider—Idiopidae	<i>Misgolas collensis</i> (Todd, 1945)	OL	
Spider—Idiopidae	<i>Misgolas dendyi</i> (Hogg, 1901)		
Spider—Idiopidae	<i>Misgolas depressa</i> (Forster, 1968)	OL	
Spider—Idiopidae	<i>Misgolas dunedinensis</i> (Forster, 1968)	OL	
Spider—Idiopidae	<i>Misgolas gilliesi</i> (O.P.-Cambridge, 1877)		
Spider—Idiopidae	<i>Misgolas grandis</i> (Forster, 1968)		
Spider—Idiopidae	<i>Misgolas buttoni</i> (O.P.-Cambridge, 1879)		
Spider—Idiopidae	<i>Misgolas isolata</i> (Forster, 1968)	OL	
Spider—Idiopidae	<i>Misgolas johnsi</i> (Forster, 1968)		
Spider—Idiopidae	<i>Misgolas kakabuensis</i> (Forster, 1968)	OL	
Spider—Idiopidae	<i>Misgolas kakanuiensis</i> (Forster, 1968)	OL	
Spider—Idiopidae	<i>Misgolas lomasi</i> (Forster, 1968)	OL	
Spider—Idiopidae	<i>Misgolas magna</i> (Forster, 1968)		
Spider—Idiopidae	<i>Misgolas marplesi</i> (Todd, 1945)		
Spider—Idiopidae	<i>Misgolas maxima</i> (Forster, 1968)	OL	
Spider—Idiopidae	<i>Misgolas medialis</i> (Forster, 1968)	OL	
Spider—Idiopidae	<i>Misgolas myersi</i> (Forster, 1968)		
Spider—Idiopidae	<i>Misgolas napua</i> (Forster, 1968)		
Spider—Idiopidae	<i>Misgolas orepukiensis</i> (Forster, 1968)		
Spider—Idiopidae	<i>Misgolas parrotti</i> (Forster, 1968)		
Spider—Idiopidae	<i>Misgolas pilama</i> (Forster, 1968)	OL	
Spider—Idiopidae	<i>Misgolas prina</i> (Forster, 1968)		
Spider—Idiopidae	<i>Misgolas reducta</i> (Forster, 1968)		
Spider—Idiopidae	<i>Misgolas secunda</i> (Forster, 1968)	OL	
Spider—Idiopidae	<i>Misgolas sinclairi</i> (Forster, 1968)		
Spider—Idiopidae	<i>Misgolas sylvatica</i> (Forster, 1968)	OL	
Spider—Idiopidae	<i>Misgolas toddae</i> (Forster, 1968)	OL	
Spider—Idiopidae	<i>Misgolas vellosa</i> (Forster, 1968)		
Spider—Idiopidae	<i>Misgolas wanganuiensis</i> (Todd, 1945)		
Stag beetle	<i>Mitophyllus gibbosus</i> (Broun, 1886)		
Darkling beetle	<i>Mitua triangularis</i> Watt, 1992		
Darkling beetle	<i>Mitua tuberculicostata</i> (White, 1846)		
Mirid bug	<i>Monopharsus annulatus</i> Eyles & Carvalho, 1995		
Spider—Amaurobiidae	<i>Muritaia kaituna</i> Forster & Wilton, 1973	OL	
Spider—Amaurobiidae	<i>Muritaia longispinata</i> Forster & Wilton, 1973	OL	
Spider—Amaurobiidae	<i>Muritaia orientalis</i> Forster & Wilton, 1973	OL	
Spider—Amaurobiidae	<i>Muritaia parabusa</i> Forster & Wilton, 1973	OL	
Nematode	<i>Mylonchulus arenicolus</i> Clark, 1961	OL	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Data Deficient—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Spider—Desidae	<i>Myro marinus</i> (Goyen, 1890)		
Spider—Stiphidiidae	<i>Nanocambridgea grandis</i> Blest & Vink, 2000	OL	
Snail	<i>Negulus senex</i> Iredale, 1913	OL	
Spider—Ctenidae	<i>Nemoctenus aurens</i> (Hogg, 1911)		
Earthworm	<i>Neochaeta forsteri</i>		
Earthworm	<i>Neochaeta salmoni</i>	OL	
Earthworm	<i>Neodrilus agilis</i>		
Earthworm	<i>Neodrilus campestris</i>		
Earthworm	<i>Neodrilus dissimilis</i>	OL	
Earthworm	<i>Neodrilus edwardsi</i>		
Earthworm	<i>Neodrilus polycystis</i>		
Spider—Neolanidae	<i>Neolana pallida</i> Forster & Wilton, 1973		
Spider—Neolanidae	<i>Neolana septentrionalis</i> Forster & Wilton, 1973	OL	
Spider—Desidae	<i>Neomyro amplius</i> Forster & Wilton, 1973		
Spider—Desidae	<i>Neomyro circe</i> Forster & Wilton, 1973		
Spider—Desidae	<i>Neomyro scitulus</i> (Urquhart, 1891)		
Spider—Agelenidae	<i>Neoramia allanae</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Neoramia alta</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Neoramia finschi</i> (L. Koch, 1872)	OL	
Spider—Agelenidae	<i>Neoramia fiordensis</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Neoramia koba</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Neoramia komata</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Neoramia mamoea</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Neoramia marama</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Neoramia matua</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Neoramia minuta</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Neoramia nana</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Neoramia oroua</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Neoramia otagoa</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Neoramia raua</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Neoramia setosa</i> (Bryant, 1935)		
Spider—Agelenidae	<i>Neorepukia bama</i> Forster & Wilton, 1973	OL	
Spider—Amphinectidae	<i>Neororea homerica</i> Forster & Wilton, 1973	OL	
Spider—Araneidae	<i>Neoscona orientalis</i> (Urquhart, 1887)		
Spider—Synotaxidae	<i>Nomaua arborea</i> Forster, 1990	OL	
Spider—Synotaxidae	<i>Nomaua cauda</i> Forster, 1990		
Spider—Synotaxidae	<i>Nomaua crinifrons</i> (Urquhart, 1891)		
Spider—Synotaxidae	<i>Nomaua nelson</i> Forster, 1990		
Spider—Synotaxidae	<i>Nomaua perdita</i> Forster, 1990		
Spider—Synotaxidae	<i>Nomaua waikaremoana</i> Forster, 1990	OL	
Peace's weevil	<i>Notbaldonus peacei</i> (Broun, 1880)		
Spider—Gnaphosidae	<i>Notiodrassus distinctus</i> Bryant, 1935		
Spider—Gnaphosidae	<i>Notiodrassus fiordensis</i> Forster, 1979		
Spider—Desidae	<i>Notomatachia cantuaria</i> Forster, 1970		
Spider—Desidae	<i>Notomatachia birsuta</i> (Marples, 1962)		
Spider—Desidae	<i>Notomatachia wiltoni</i> Forster, 1970		
Earthworm	<i>Notoscolex bakeaphilus</i>		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Data Deficient—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Spider—Linyphiidae	<i>Novafroneta nova</i> Blest & Vink, 2003	OL	
Spider—Linyphiidae	<i>Novafroneta parmulata</i> Blest, 1979	OL	
Spider—Linyphiidae	<i>Novafroneta truncata</i> Blest & Vink, 2003	OL	
Spider—Araneidae	<i>Novakiella trituberculosa</i> (Roewer, 1942)	SO	
Spider—Linyphiidae	<i>Novalaestesia anceps</i> Millidge, 1988	OL	
Spider—Linyphiidae	<i>Novalaestesia atra</i> Blest & Vink, 2003	OL	
Spider—Salticidae	<i>Ocrisiona cinerea</i> (L. Koch, 1879)	SO	
Spider—Salticidae	<i>Ocrisiona leucocomis</i> (L. Koch, 1879)	SO	
Earthworm	<i>Octochaetus ?levis</i>	OL	
Earthworm	<i>Octochaetus ?microchaetus</i>	OL	
Earthworm	<i>Octochaetus antarcticus</i> (Beddard, 1889)	OL	
Earthworm	<i>Octochaetus brucei</i>		
Earthworm	<i>Octochaetus buttoni</i>		
Earthworm	<i>Octochaetus kapitiensis</i>	OL	
Earthworm	<i>Octochaetus michaelseni</i> Benham, 1904		
Earthworm	<i>Octochaetus pelorus</i>		
Earthworm	<i>Octochaetus ravus</i>	OL	
Earthworm	<i>Octochaetus sylvestris</i>		
Earthworm	<i>Octocbaetus thomasi</i>		
Earthworm	<i>Octochaetus trocystis</i>		
Nematode	<i>Ogma campbelli</i> Wouts, Yeates & Loof, 1999	OL	
Nematode	<i>Ogma crenulatum</i> Wouts, Yeates & Loof, 1999	OL	
Spider—Amphinectidae	<i>Oparara karamea</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Oparara vallus</i> (Marples, 1959)		
Spider—Agelenidae	<i>Oramia littoralis</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Oramia mackerowii</i> (Marples, 1959)	OL	
Spider—Agelenidae	<i>Oramia occidentalis</i> (Marples, 1959)	OL	
Spider—Agelenidae	<i>Oramiella wisei</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Orepukia alta</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Orepukia catlinsensis</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Orepukia dugdalei</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Orepukia egmontensis</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Orepukia florae</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Orepukia geophila</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Orepukia grisea</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Orepukia insula</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Orepukia orophila</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Orepukia pallida</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Orepukia prina</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Orepukia redacta</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Orepukia riparia</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Orepukia sabua</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Orepukia similis</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Orepukia simplex</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Orepukia sorenseni</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Orepukia tanea</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Orepukia tonga</i> Forster & Wilton, 1973	OL	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Spider—Agelenidae	<i>Orepukia virtuta</i> Forster & Wilton, 1973		
Moth	<i>Orocrambus ornatus</i> (Philpott, 1927)		
Moth	<i>Orocrambus punctellus</i> (Hudson, 1950)		
Moth	<i>Orocrambus sopbronellus</i> (Meyrick, 1885)	OL	
Spider—Orsolobidae	<i>Orongia medialis</i> Forster & Platnick, 1985	OL	
Spider—Orsolobidae	<i>Orongia motueka</i> Forster & Platnick, 1985	OL	
Spider—Orsolobidae	<i>Orongia whangamoa</i> Forster & Platnick, 1985	OL	
Spider—Desidae	<i>Otagoa nova</i> Forster, 1970		
Spider—Desidae	<i>Otagoa wiltoni</i> Forster, 1970	OL	
Spider—Amaurobiidae	<i>Otira canasta</i> Forster & Wilton, 1973	OL	
Spider—Amaurobiidae	<i>Otira indura</i> Forster & Wilton, 1973		
Spider—Amaurobiidae	<i>Otira liana</i> Forster & Wilton, 1973		
Spider—Amaurobiidae	<i>Otira parva</i> Forster & Wilton, 1973	OL	
Spider—Amaurobiidae	<i>Otira satula</i> Forster & Wilton, 1973		
Spider—Amaurobiidae	<i>Otira terricola</i> Forster & Wilton, 1973		
Spider—Synotaxidae	<i>Pabora cantuaria</i> Forster, 1990	OL	
Spider—Synotaxidae	<i>Pabora kaituna</i> Forster, 1990	OL	
Spider—Synotaxidae	<i>Pabora media</i> Forster, 1990		
Spider—Synotaxidae	<i>Pabora montana</i> Forster, 1990	OL	
Spider—Synotaxidae	<i>Pabora raktura</i> Forster, 1990		
Spider—Synotaxidae	<i>Pabora taranaki</i> Forster, 1990		
Spider—Synotaxidae	<i>Pabora wiltoni</i> Forster, 1990	OL	
Spider—Synotaxidae	<i>Paboroides courti</i> Forster, 1990		
Spider—Amaurobiidae	<i>Pakeha buechlerae</i> Forster & Wilton, 1973		
Spider—Amaurobiidae	<i>Pakeha duplex</i> Forster & Wilton, 1973		
Spider—Amaurobiidae	<i>Pakeha biloa</i> Forster & Wilton, 1973	OL	
Spider—Amaurobiidae	<i>Pakeha inornata</i> Forster & Wilton, 1973		
Spider—Amaurobiidae	<i>Pakeha insignita</i> Forster & Wilton, 1973	OL	
Spider—Amaurobiidae	<i>Pakeha lobata</i> Forster & Wilton, 1973	OL	
Spider—Amaurobiidae	<i>Pakeha manapouri</i> Forster & Wilton, 1973	OL	
Spider—Amaurobiidae	<i>Pakeha maxima</i> Forster & Wilton, 1973		
Spider—Amaurobiidae	<i>Pakeha media</i> Forster & Wilton, 1973		
Spider—Amaurobiidae	<i>Pakeha minima</i> Forster & Wilton, 1973		
Spider—Amaurobiidae	<i>Pakeha paratecta</i> Forster & Wilton, 1973		
Spider—Amaurobiidae	<i>Pakeha parrotti</i> Forster & Wilton, 1973		
Spider—Amaurobiidae	<i>Pakeha pula</i> Forster & Wilton, 1973		
Spider—Amaurobiidae	<i>Pakeha subiecta</i> Forster & Wilton, 1973		
Spider—Amaurobiidae	<i>Pakeha tecta</i> Forster & Wilton, 1973		
Spider—Desidae	<i>Panoa contorta</i> Forster, 1970		
Spider—Desidae	<i>Panoa fiordensis</i> Forster, 1970	OL	
Spider—Desidae	<i>Panoa mora</i> Forster, 1970		
Spider—Desidae	<i>Panoa tapanuiensis</i> Forster, 1970	OL	
Spider—Dictynidae	<i>Paradictyna ilamia</i> Forster, 1970	OL	
Spider—Dictynidae	<i>Paradictyna rufoflava</i> (Chamberlain, 1946)		
Spider—Linyphiidae	<i>Parafroneta ambigua</i> Blest, 1979		
Spider—Linyphiidae	<i>Parafroneta baurokoae</i> Blest & Vink, 2002	OL	
Spider—Linyphiidae	<i>Parafroneta birsuta</i> Blest & Vink, 2003	OL	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Data Deficient—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Spider—Linyphiidae	<i>Parafroneta minuta</i> Blest, 1979		
Spider—Linyphiidae	<i>Parafroneta monticola</i> Blest, 1979	OL	
Spider—Linyphiidae	<i>Parafroneta persimilis</i> Blest, 1979		
Spider—Linyphiidae	<i>Parafroneta pilosa</i> Blest & Vink, 2003	OL	
Spider—Linyphiidae	<i>Parafroneta westlandica</i> Blest & Vink, 2002		
Spider—Orsolobidae	<i>Paralobus salmoni</i> (Forster, 1956)		
Spider—Amphinectidae	<i>Paramamoea aquilonalis</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Paramamoea arawa</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Paramamoea incerta</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Paramamoea incertoides</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Paramamoea pandora</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Paramamoea paradisica</i> Forster & Wilton, 1973	OL	
Spider—Amphinectidae	<i>Paramamoea parva</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Paramamoea urewera</i> Forster & Wilton, 1973		
Spider—Amphinectidae	<i>Paramamoea waipoua</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Paramyro apicus</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Paramyro parapicus</i> Forster & Wilton, 1973	OL	
Spider—Anapidae	<i>Paranapis insula</i> (Forster, 1951)		
Spider—Anapidae	<i>Paranapis isolata</i> Platnick & Forster, 1989	OL	
Spider—Micropholcommatidae	<i>Parapua punctata</i> Forster, 1959		
Spider—Pararchaeidae	<i>Pararchaea alba</i> Forster, 1955		
Spider—Pararchaeidae	<i>Pararchaea rubra</i> (Forster, 1949)		
Cylindrical rove beetle	<i>Paratorbus atifer</i> (McColl, 1982)		
Cylindrical rove beetle	<i>Paratorbus flexuosus</i> (McColl, 1982)		
Nematode	<i>Paratylenchus halophilus</i> Wouts, 1966	OL	
Spider—Amaurobiidae	<i>Paravoca opaca</i> Forster & Wilton, 1973		
Spider—Amaurobiidae	<i>Paravoca otagoensis</i> Forster & Wilton, 1973		
Scarab beetle	<i>Pericoptus nitidulus</i> Broun, 1880	OL	
Earthworm	<i>Perieodrilus lateralis</i>		
Earthworm	<i>Perieodrilus montanus</i>		
Earthworm	<i>Perieodrilus plunketi</i>	OL	
Earthworm	<i>Perieodrilus ricardi</i>	OL	
Earthworm	<i>Perionyx egmonti</i>		
Earthworm	<i>Perionyx belophilus</i>	OL	
Earthworm	<i>Perionyx perionychopsis</i>	OL	
Earthworm	<i>Perionyx shoceanus</i>		
Moth	<i>Petasactis technica</i> Meyrick, 1888	OL	
Moth	<i>Phaeosaces lindsayae</i>	OL	
Spider—Philodromidae	<i>Philodromus rubrofrontus</i> Urquhart, 1891		
Spider—Theridiidae	<i>Pholcomma antipodianum</i> (Forster, 1955)		
Spider—Theridiidae	<i>Pholcomma bickmani</i> Forster, 1964		
Spider—Theridiidae	<i>Pholcomma turbotti</i> (Marples, 1956)		
Spider—Theridiidae	<i>Phoroncidia pukeiwa</i> (Marples, 1955)		
Spider—Theridiidae	<i>Phoroncidia puketoru</i> (Marples, 1955)		
Spider—Theridiidae	<i>Phoroncidia quadrata</i> (O.P.-Cambridge, 1879)		
Flax snail (pupuharakeke)	<i>Placostylus (Maoristylus) ambagiosus bancoxii</i>	RF, OL	VU B1+2abcde (for <i>P. ambagiosus</i> s.l.)

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Data Deficient—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Earthworm	<i>Plagiochaeta lineata</i>		
Earthworm	<i>Plagiochaeta stewartensis</i>		
Earthworm	<i>Plagiochaeta sylvestris</i>		
Moth	<i>Platyptilia bokowbitalis</i> Hudson, 1939		
Spider—Cycloctenidae	<i>Plectophanes alta</i> Forster, 1964		
Spider—Cycloctenidae	<i>Plectophanes archeyi</i> Forster, 1964		
Spider—Cycloctenidae	<i>Plectophanes frontalis</i> Bryant, 1935		
Spider—Cycloctenidae	<i>Plectophanes hollowayae</i> Forster, 1964	OL	
Spider—Cycloctenidae	<i>Plectophanes pilgrimi</i> Forster, 1964		
Spider—Salticidae	<i>Plexippus sylvanus</i> Urquhart, 1892	SO	
Earthworm	<i>Plutellus aucklandicus</i>	OL	
Earthworm	<i>Plutellus parvus</i>		
Earthworm	<i>Plutellus stewartensis</i>	OL	
Spider—Linyphiidae	<i>Poecilafroneta caudata</i> Blest, 1979		
Earthworm	<i>Pontodrilus lacustris</i>	OL	
Earthworm	<i>Pontodrilus matsushimensis</i> var. <i>chatbamianus</i>	OL	
Spider—Hahniidae	<i>Poroides rima</i> (Forster, 1970)		
Spider—Hahniidae	<i>Poroides tasmani</i> (Forster, 1970)	OL	
Spider—Agelenidae	<i>Porotaka detrita</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Porotaka florae</i> Forster & Wilton, 1973	OL	
Spider—Hexathelidae	<i>Porrhothele blanda</i> Forster, 1968		
Spider—Hexathelidae	<i>Porrhothele modesta</i> Forster, 1968		
Spider—Hexathelidae	<i>Porrhothele quadrigyna</i> Forster, 1968		
Spider—Orsolobidae	<i>Pounamuella complexa</i> (Forster, 1956)		
Spider—Orsolobidae	<i>Pounamuella bauroko</i> Forster & Platnick, 1985	OL	
Spider—Orsolobidae	<i>Pounamuella hollowayae</i> (Forster, 1956)	OL	
Spider—Orsolobidae	<i>Pounamuella insula</i> Forster & Platnick, 1985	OL	
Spider—Orsolobidae	<i>Pounamuella kuscheli</i> Forster & Platnick, 1985	OL	
Spider—Orsolobidae	<i>Pounamuella ramsayi</i> (Forster, 1956)		
Spider—Araneidae	<i>Prasonica discolor</i> (Urquhart, 1887)		
Spider—Araneidae	<i>Prasonica plagiata</i> (Simon, 1896)		
Nematode	<i>Procyrnea kea</i> Clark, 1978		
Scarab/chafer beetle	<i>Prodontria patricki</i> Emerson, 1997		
Nematode	<i>Proleptonchus saccatus</i> (Clark, 1962) Andrassy, 1963	OL	
Spider—Linyphiidae	<i>Promynoglenes miniscula</i> Blest & Vink, 2003	OL	
Spider—Linyphiidae	<i>Promynoglenes minuta</i> Blest & Vink, 2002	OL	
Spider—Linyphiidae	<i>Promynoglenes parvula</i> Blest, 1979		
Spider—Linyphiidae	<i>Protoerigone otagoa</i> Blest, 1979		
Spider—Linyphiidae	<i>Pseudafroneta lineata</i> Blest, 1979		
Spider—Linyphiidae	<i>Pseudafroneta maxima</i> Blest, 1979	OL	
Spider—Linyphiidae	<i>Pseudafroneta prominula</i> Blest, 1979	OL	
Moth	<i>Pseudocoremia albafasciata</i> (Philpott, 1915)		
Moth	<i>Pseudocoremia lutea</i> (Philpott, 1914)		
Spider—Amphinectidae	<i>Rangitata peelensis</i> Forster & Wilton, 1973	OL	
Spider—Desidae	<i>Rapua australis</i> Forster, 1970		
Spider—Amphinectidae	<i>Reinga apica</i> Forster & Wilton, 1973	OL	
Spider—Amphinectidae	<i>Reinga aucklandensis</i> (Marples, 1959)	OL	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Data Deficient—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Spider—Amphinectidae	<i>Reinga grossa</i> Forster & Wilton, 1973	OL	
Spider—Amphinectidae	<i>Reinga media</i> Forster & Wilton, 1973	OL	
Spider—Amphinectidae	<i>Reinga waipoua</i> Forster & Wilton, 1973		
Nematode	<i>Rbigonema zealandica</i> Clark, 1978	OL	
Earthworm	<i>Rhododrilus aduncocystis</i>		
Earthworm	<i>Rhododrilus agathis</i>	OL	
Earthworm	<i>Rhododrilus albidus</i>		
Earthworm	<i>Rhododrilus aquaticus</i>		
Earthworm	<i>Rhododrilus attenuatus</i>		
Earthworm	<i>Rhododrilus benhami</i>		
Earthworm	<i>Rhododrilus besti</i>	OL	
Earthworm	<i>Rhododrilus cockaynei</i>		
Earthworm	<i>Rhododrilus disparatus</i>		
Earthworm	<i>Rhododrilus dobsoni</i>	OL	
Earthworm	<i>Rhododrilus edulis</i>		
Earthworm	<i>Rhododrilus buttoni</i>	OL	
Earthworm	<i>Rhododrilus insularis</i>	OL	
Earthworm	<i>Rhododrilus intermedius</i>		
Earthworm	<i>Rhododrilus kermadecensis</i>	OL	
Earthworm	<i>Rhododrilus leptomerus</i>		
Earthworm	<i>Rhododrilus macroseptus</i>		
Earthworm	<i>Rhododrilus microgaster</i>	OL	
Earthworm	<i>Rhododrilus minimus</i>		
Earthworm	<i>Rhododrilus minutus</i>		
Earthworm	<i>Rhododrilus papaensis</i>		
Earthworm	<i>Rhododrilus parvus</i>	OL	
Earthworm	<i>Rhododrilus ravus</i>	OL	
Earthworm	<i>Rhododrilus robustus</i>		
Earthworm	<i>Rhododrilus rosae</i>		
Earthworm	<i>Rhododrilus sexpapillatus</i> Dyne, 1980		
Earthworm	<i>Rhododrilus similis</i>		
Earthworm	<i>Rhododrilus subtilis</i>	OL	
Earthworm	<i>Rhododrilus sutherlandi</i>		
Earthworm	<i>Rhododrilus tetratbeaca</i>	OL	
Lygaeid bug	<i>Rhypodes brevipilis</i> Eyles, 1990		
Spider—Hahniidae	<i>Rinawa bola</i> Forster, 1970		
Spider—Hahniidae	<i>Rinawa cantuaria</i> Forster, 1970		
Spider—Hahniidae	<i>Rinawa otagoensis</i> Forster, 1970		
Spider—Hahniidae	<i>Rinawa pula</i> Forster, 1970		
Spider—Amphinectidae	<i>Rorea otagoensis</i> Forster & Wilton, 1973	OL	
Nematode	<i>Rotylenchus kabikateae</i> Ryss & Wouts, 1997	OL	
Nematode	<i>Rotylenchus labiodiscus</i> Wouts & Sturhan, 1999	OL	
Nematode	<i>Rotylenchus nelsonensis</i> Ryss & Wouts, 1997	OL	
Nematode	<i>Ruamowhitia orae</i> Yeates, 1967	OL	
Spider—Synotaxidae	<i>Runga akaroa</i> Forster, 1990		
Spider—Synotaxidae	<i>Runga flora</i> Forster, 1990		
Spider—Synotaxidae	<i>Runga moana</i> Forster, 1990		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Data Deficient—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Spider—Salticidae	<i>Salticus albobarbatus</i> Powell, 1873		
Spider—Salticidae	<i>Salticus albopalpis</i> Urquhart, 1885		
Spider—Salticidae	<i>Salticus alpinus</i> Urquhart, 1885		
Spider—Salticidae	<i>Salticus appressus</i> Powell, 1873		
Spider—Salticidae	<i>Salticus atratus</i> Powell, 1873		
Spider—Salticidae	<i>Salticus curvus</i> Urquhart, 1885		
Spider—Salticidae	<i>Salticus fumosus</i> Powell, 1873		
Spider—Salticidae	<i>Salticus furvus</i> Urquhart, 1885		
Spider—Salticidae	<i>Salticus mustelinus</i> Powell, 1873		
Spider—Salticidae	<i>Salticus pbrinoides</i> Walckanaer, 1837		
Spider—Salticidae	<i>Salticus tabinus</i> Urquhart, 1885		
Spider—Salticidae	<i>Salticus tenebricus</i> Urquhart, 1885		
Spider—Salticidae	<i>Salticus vnotatus</i> Powell, 1873		
Spider—Salticidae	<i>Salticus zanthofrontalis</i> Urquhart, 1886		
Beetle	<i>Sciacharis yakasensis</i> Franz, 1985		
Nematode	<i>Sciadiocara tarapunga</i> Clark, 1978		
Spider—Hahniidae	<i>Scotospilus divisa</i> (Forster, 1970)	OL	
Spider—Hahniidae	<i>Scotospilus nelsonensis</i> (Forster, 1970)	OL	
Spider—Hahniidae	<i>Scotospilus plena</i> (Forster, 1970)		
Spider—Hahniidae	<i>Scotospilus westlandica</i> (Forster, 1970)	OL	
Nematode	<i>Sectonema sica</i> Clark, 1964	OL	
Chafer beetle	<i>Sericospilus minor</i>	OL	
Spider—Thomisidae	<i>Sidymella benhami</i> (Hogg, 1911)		
Earthworm	<i>Spenceriella argillae</i>	OL	
Earthworm	<i>Spenceriella gigantea</i>		
Earthworm	<i>Spenceriella pallida</i>	OL	
Earthworm	<i>Spenceriella shakespearei</i>		
Spider—Nemesiidae	<i>Stanwellia bipectinata</i> (Todd, 1945)	OL	
Spider—Nemesiidae	<i>Stanwellia hollowayi</i> (Forster, 1968)		
Spider—Nemesiidae	<i>Stanwellia boubora</i> (Forster, 1968)	OL	
Spider—Nemesiidae	<i>Stanwellia kaituna</i> (Forster, 1968)		
Spider—Nemesiidae	<i>Stanwellia media</i> (Forster, 1968)		
Spider—Nemesiidae	<i>Stanwellia regia</i> (Forster, 1968)	OL	
Moth	<i>Stathmopoda endotherma</i> Meyrick, 1931		
Spider—Theridiidae	<i>Steatoda nobilosa</i> Urquhart, 1893 (incertae sedis)	OL	
Chafer beetle	<i>Stethaspis convexa</i> (Given, 1952)		
Chafer beetle	<i>Stethaspis pulchra</i> (Given, 1952)		
Moth	<i>Stigmella progamma</i> (Meyrick, 1924)	OL	
Moth	<i>Stigmella propalaea</i> (Meyrick, 1889)	OL	
Bug	<i>Stizocephalus brevirostris</i> Eyles, 1970	SO	
Spider—Orsolobidae	<i>Subantarctia centralis</i> Forster & Platnick, 1985		
Spider—Orsolobidae	<i>Subantarctia dugdalei</i> Forster, 1956		
Spider—Orsolobidae	<i>Subantarctia fiordensis</i> Forster, 1956		
Spider—Orsolobidae	<i>Subantarctia florae</i> Forster, 1956		
Spider—Orsolobidae	<i>Subantarctia muka</i> Forster & Platnick, 1985	OL	
Spider—Orsolobidae	<i>Subantarctia penara</i> Forster & Platnick, 1985	OL	
Spider—Orsolobidae	<i>Subantarctia stewartensis</i> Forster, 1956	OL	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Data Deficient—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Spider—Orsolobidae	<i>Subantarctia trina</i> Forster & Platnick, 1985		
Earthworm	<i>Sylovodrilus gravus</i>		
Fly	<i>Sympycnus alchymicus</i> (Parent, 1933)	OL	
Spider—Thomisidae	<i>Synema suteri</i> Dahl, 1907		
Spider—Gnaphosidae	<i>Taieria erekus</i> (L. Koch, 1873)		
Spider—Gnaphosidae	<i>Taieria kaituna</i> Forster, 1979		
Spider—Gnaphosidae	<i>Taieria miranda</i> Forster, 1979		
Spider—Gnaphosidae	<i>Taieria obtusa</i> Forster, 1979		
Nematode	<i>Takamangai waenga</i> Yeates, 1967	OL	
Spider—Orsolobidae	<i>Tangata alpina</i> (Forster, 1956)		
Spider—Orsolobidae	<i>Tangata furcata</i> Forster & Platnick, 1985		
Spider—Orsolobidae	<i>Tangata kohuka</i> Forster & Platnick, 1985	OL	
Spider—Orsolobidae	<i>Tangata muribiku</i> Forster & Platnick, 1985		
Spider—Orsolobidae	<i>Tangata nigra</i> Forster & Platnick, 1985		
Spider—Orsolobidae	<i>Tangata orepukiensis</i> (Forster, 1956)		
Spider—Orsolobidae	<i>Tangata otago</i> Forster & Platnick, 1985	OL	
Spider—Orsolobidae	<i>Tangata parafurcata</i> Forster & Platnick, 1985	OL	
Spider—Orsolobidae	<i>Tangata plena</i> (Forster, 1956)		
Spider—Orsolobidae	<i>Tangata pouaka</i> Forster & Platnick, 1985	OL	
Spider—Orsolobidae	<i>Tangata rakiura</i> (Forster, 1956)		
Spider—Orsolobidae	<i>Tangata stewartensis</i> (Forster, 1956)	OL	
Spider—Orsolobidae	<i>Tangata sylvester</i> Forster & Platnick, 1985		
Spider—Orsolobidae	<i>Tangata tautuku</i> Forster & Platnick, 1985	OL	
Spider—Orsolobidae	<i>Tangata townsendi</i> Forster & Platnick, 1985	OL	
Spider—Orsolobidae	<i>Tangata waipoua</i> Forster & Platnick, 1985		
Spider—Agelenidae	<i>Tararua celeripes</i> (Urquhart, 1891)		
Spider—Agelenidae	<i>Tararua clara</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Tararua diversa</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Tararua foordi</i> Forster & Wilton, 1973	OL	
Spider—Agelenidae	<i>Tararua puna</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Tararua ratuma</i> Forster & Wilton, 1973		
Spider—Agelenidae	<i>Tararua versuta</i> Forster & Wilton, 1973		
Spider—Orsolobidae	<i>Tautukua isolata</i> Forster & Platnick, 1985	OL	
Spider—Cyatholipidae	<i>Tekella bisetosa</i> Forster, 1988		
Spider—Cyatholipidae	<i>Tekella unisetosa</i> Forster, 1988		
Spider—Tetragnathidae	<i>Tetragnatha flavigula</i> Urquhart, 1891		
Spider—Tetragnathidae	<i>Tetragnatha multipunctata</i> Urquhart, 1891		
Spider—Tetragnathidae	<i>Tetragnatha nigricans</i> Dalmas, 1918		
Nematode	<i>Tetrameres tarapungae</i> Clark, 1978		
Spider—Micropholcommatidae	<i>Textricella antipoda</i> Forster, 1959		
Spider—Micropholcommatidae	<i>Textricella aucklandica</i> Forster, 1955		
Spider—Micropholcommatidae	<i>Textricella mcfarlanei</i> Forster, 1959	OL	
Spider—Micropholcommatidae	<i>Textricella nigra</i> Forster, 1959		
Spider—Micropholcommatidae	<i>Textricella propinqua</i> Forster, 1959		
Spider—Micropholcommatidae	<i>Textricella pusilla</i> Forster, 1959		
Spider—Micropholcommatidae	<i>Textricella signata</i> Forster, 1959		
Moth	<i>Thectophila acmotypa</i> Meyrick, 1927	OL	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Data Deficient—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Nematode	<i>Thelastoma meadsi</i> Clark, 1978		
Spider—Theridiidae	<i>Theridion albocinctum</i> Urquhart, 1892	OL	
Spider—Theridiidae	<i>Theridion ampliatum</i> Urquhart, 1892		
Spider—Theridiidae	<i>Theridion argentatum</i> (Roewer, 1942)		
Spider—Theridiidae	<i>Theridion cruciferum</i> Urquhart, 1886		
Spider—Theridiidae	<i>Theridion flabelliferum</i> Urquhart, 1887		
Spider—Theridiidae	<i>Theridion gibbosa</i> Urquhart, 1894		
Spider—Theridiidae	<i>Theridion gracilipes</i> Urquhart, 1889		
Spider—Theridiidae	<i>Theridion longicrure</i> Marples, 1956		
Spider—Theridiidae	<i>Theridion prophryeticum</i> Urquhart, 1889		
Spider—Theridiidae	<i>Theridion pumilio</i> Urquhart, 1886		
Spider—Theridiidae	<i>Theridion punicapunctatum</i> Urquhart, 1891		
Spider—Theridiidae	<i>Theridion rufipes</i> (Lucas, 1846)		
Spider—Theridiidae	<i>Theridion squalidæ</i> Urquhart, 1886		
Spider—Theridiidae	<i>Theridion viridana</i> Urquhart, 1887		
Moth	"Tinea" <i>texta</i> Meyrick, 1931		
Moth	<i>Tingena loxotis</i> (Meyrick, 1905)		
Snail	<i>Tornatellides subperforatus kermadecensis</i> Pilsbry & Cooke, 1915	OL	
Snail	<i>Tornatellinops iredalei</i> (Pilsbry & Cooke, 1915)	OL	
Spider—Cycloctenidae	<i>Toxopsiella alpina</i> Forster, 1964		
Spider—Cycloctenidae	<i>Toxopsiella australis</i> Forster, 1964		
Spider—Cycloctenidae	<i>Toxopsiella centralis</i> Forster, 1964		
Spider—Cycloctenidae	<i>Toxopsiella dugdalei</i> Forster, 1964		
Spider—Cycloctenidae	<i>Toxopsiella laurencei</i> Forster, 1964		
Spider—Cycloctenidae	<i>Toxopsiella medialis</i> Forster, 1964		
Spider—Cycloctenidae	<i>Toxopsiella minuta</i> Forster, 1964		
Spider—Cycloctenidae	<i>Toxopsiella nelsonensis</i> Forster, 1979		
Spider—Cycloctenidae	<i>Toxopsiella orientalis</i> Forster, 1964		
Spider—Cycloctenidae	<i>Toxopsiella parrotti</i> Forster, 1964		
Spider—Cycloctenidae	<i>Toxopsiella perplexa</i> Forster, 1964		
Spider—Desidae	<i>Toxopsoides buttoni</i> Forster & Wilton, 1973	OL	
Moth	<i>Trachypepla nimbosa</i> Philpott, 1930	OL	
Nematode	<i>Trissonchulus littoralis</i> Yeates, 1967	OL	
Spider—Salticidae	<i>Trite herbigrada</i> (Urquhart, 1889)		
Spider—Salticidae	<i>Trite mustilina</i> (Powell, 1873)		
Spider—Salticidae	<i>Trite parvula</i> Bryant, 1935		
Spider—Salticidae	<i>Trite urvillei</i> (Dalmas, 1917)		
Spider—Desidae	<i>Tuakana mirada</i> Forster, 1971		
Spider—Desidae	<i>Tuakana wiltoni</i> Forster, 1971	OL	
Snail	<i>Tubuata raoulensis</i> (Pilsbry & Cooke, 1915)	OL	
Spider—Orsolobidae	<i>Turretia dugdalei</i> Forster & Platnick, 1986	OL	
Soft scale	<i>Umbonichiton adelus</i>	OL	
Soft scale	<i>Umbonichiton jubatus</i>	OL	
Soft scale	<i>Umbonichiton pellaspis</i>	OL	
Weevil	<i>Unas piceus</i> Broun, 1909		
Nematode	<i>Viktorocara torea</i> Clark, 1978		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

Data Deficient—continued

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Spider—Dictynidae	<i>Viridictyna australis</i> Forster, 1971		
Spider—Dictynidae	<i>Viridictyna kikkawai</i> Forster, 1971		
Spider—Dictynidae	<i>Viridictyna nelsonensis</i> Forster, 1971		
Spider—Dictynidae	<i>Viridictyna parva</i> Forster, 1971		
Spider—Dictynidae	<i>Viridictyna picata</i> Forster, 1971		
Spider—Orsolobidae	<i>Waiporia algida</i> (Forster, 1956)		
Spider—Orsolobidae	<i>Waiporia egmont</i> Forster & Platnick, 1986	OL	
Spider—Orsolobidae	<i>Waiporia extensa</i> (Forster, 1956)		
Spider—Orsolobidae	<i>Waiporia haavea</i> Forster & Platnick, 1986	OL	
Spider—Orsolobidae	<i>Waiporia hornabrooki</i> (Forster, 1956)		
Spider—Orsolobidae	<i>Waiporia mensa</i> (Forster, 1956)		
Spider—Orsolobidae	<i>Waiporia modica</i> (Forster, 1956)		
Spider—Orsolobidae	<i>Waiporia owaka</i> Forster & Platnick, 1986		
Spider—Orsolobidae	<i>Waiporia ruabine</i> Forster & Platnick, 1986	OL	
Spider—Orsolobidae	<i>Waiporia tuata</i> Forster & Platnick, 1986	OL	
Spider—Orsolobidae	<i>Waiporia willtoni</i> Forster & Platnick, 1986	OL	
Spider—Orsolobidae	<i>Waipoua bila</i> Forster & Platnick, 1986		
Spider—Orsolobidae	<i>Waipoua insula</i> Forster & Platnick, 1986	OL	
Spider—Orsolobidae	<i>Waipoua montana</i> Forster & Platnick, 1986	OL	
Spider—Orsolobidae	<i>Waipoua otiana</i> Forster & Platnick, 1986	OL	
Spider—Orsolobidae	<i>Waipoua ponanga</i> Forster & Platnick, 1986		
Spider—Orsolobidae	<i>Waipoua toronui</i> Forster & Platnick, 1986		
Spider—Orsolobidae	<i>Waipoua totara</i> (Forster, 1956)	OL	
Spider—Synotaxidae	<i>Wairua reinga</i> Forster, 1991	OL	
Spider—Synotaxidae	<i>Wairua waikanae</i> Forster, 1991		
Spider—Amaurobiidae	<i>Waitetola buttoni</i> Forster & Wilton, 1974		
Spider—Amphinectidae	<i>Waterea cornigera</i> Forster & Wilton, 1974	OL	
Spider—Orsolobidae	<i>Wiltonia elongata</i> Forster & Platnick, 1986	OL	
Spider—Orsolobidae	<i>Wiltonia eylesi</i> Forster & Platnick, 1986	OL	
Spider—Orsolobidae	<i>Wiltonia fiordensis</i> Forster & Platnick, 1986	OL	
Spider—Orsolobidae	<i>Wiltonia graminicola</i> Forster & Platnick, 1986		
Spider—Orsolobidae	<i>Wiltonia lima</i> Forster & Platnick, 1986		
Spider—Orsolobidae	<i>Wiltonia nelsonensis</i> Forster & Platnick, 1986	OL	
Spider—Orsolobidae	<i>Wiltonia pecki</i> Forster & Platnick, 1986	OL	
Spider—Orsolobidae	<i>Wiltonia porina</i> Forster & Platnick, 1986	OL	
Spider—Orsolobidae	<i>Wiltonia rottoi</i> Forster & Platnick, 1986	OL	
Nematode	<i>Xiphinema waimungui</i> Yeates, Boag & Brown, 1997	OL	
Spider—Anapidae	<i>Zealanapis conica</i> (Forster, 1951)		
Spider—Anapidae	<i>Zealanapis insula</i> Platnick & Forster, 1989	OL	
Spider—Anapidae	<i>Zealanapis kuscheli</i> Platnick & Forster, 1989		
Spider—Anapidae	<i>Zealanapis matua</i> Platnick & Forster, 1989		
Spider—Anapidae	<i>Zealanapis montana</i> Platnick & Forster, 1989		
Spider—Anapidae	<i>Zealanapis otago</i> Platnick & Forster, 1989		
Spider—Anapidae	<i>Zealanapis punta</i> Platnick & Forster, 1989		
Spider—Anapidae	<i>Zealanapis waipoua</i> Platnick & Forster, 1989		
Spider—Mecysmauchenidae	<i>Zearchaea fiordensis</i> Forster, 1956		
Snail	<i>Zelandiscus worthyi</i> Climo, 1989	OL	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Nematode	<i>Zeldia punua</i> Yeates, 1967		
Robber fly	<i>Zosteria novazealandica</i> Daniels, 1988		
<b>Taxonomically indeterminate</b>			
Spider—Huttoniidae	About 20 species awaiting description		
Spider—Miturgidae	About 20 undescribed miturgid species		
Snail	<i>Allodiscus</i> "aff. <i>planulatus</i> " (NMNZ M.82155)		
Snail	<i>Allodiscus</i> "aff. <i>turbotti</i> " (NMNZ M.100283)		
Snail	<i>Allodiscus</i> "minute white" (NMNZ M.154802)		
Snail	<i>Allodiscus</i> "Mount Peel" (NMNZ M.84431)		
Weevil	<i>Anagotus</i> sp. "Moehau"		OL
Aphid	<i>Aphis</i> sp. 2		
Moth	<i>Archyala</i> sp. "Mataroa"		
Moth	<i>Archyala</i> sp. "Mount Greenland"		
Slug	<i>Athoracophorus</i> sp. 7 (NMNZ M.151433) "Warawara 2"		
Spider—Theridiosomatidae	<i>Baalzebub</i> sp.		
Snail	Charopidae sp. 2 (NMNZ M.151451)		OL
Snail	Charopidae sp. 10 (NMNZ M.85650)		
Snail	Charopidae sp. 18		OL
Snail	Charopidae sp. 21 (NMNZ M.36721)		
Snail	Charopidae sp. 26 (NMNZ M.82966)		
Snail	Charopidae sp. 28 (NMNZ M.98308)		OL
Snail	Charopidae sp. 50 (NMNZ M.22563)		
Snail	Charopidae sp. 69 (NMNZ M.128746)		
Snail	Charopidae sp. 72 (NMNZ M.99415)		OL
Snail	Charopidae sp. 76 (NMNZ M.127900)		OL
Snail	Charopidae sp. 78 (NMNZ M.28634)		
Snail	Charopidae sp. 81 (NMNZ M.127831)		
Snail	Charopidae sp. 82 (NMNZ M.99974)		
Snail	Charopidae sp. 83 (NMNZ M.79298)		
Snail	Charopidae sp. 84 (NMNZ M.127823)		
Snail	Charopidae sp. 85 (NMNZ M.127799)		
Snail	Charopidae sp. 86 (NMNZ M.125428)		
Snail	Charopidae sp. 88 (NMNZ M.127813)		
Snail	Charopidae sp. 90 (NMNZ M.127809)		
Snail	Charopidae sp. 95 (NMNZ M.127825)		
Snail	Charopidae sp. 102 (NMNZ M.116417)		
Snail	Charopidae sp. 109 (NMNZ M.155124)		
Snail	Charopidae sp. 110 (NMNZ M.30389)		
Snail	Charopidae sp. 117 (NMNZ M.102002)		
Snail	Charopidae sp. 118 (NMNZ M.123272)		
Snail	Charopidae sp. 129 (NMNZ M.127913)		
Snail	Charopidae sp. 131 (NMNZ M.127915)		
Snail	Charopidae sp. 139 (NMNZ M.32395)		
Snail	Charopidae sp. 140 (NMNZ M.70610)		
Snail	Charopidae sp. 141 (NMNZ M.23044)		
Snail	Charopidae sp. 142 (NMNZ M.72506)		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

*Data Deficient—continued*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Snail	Charopidae sp. 143 (NMNZ M.32915)		
Snail	Charopidae sp. 144 (NMNZ M.29527)		
Snail	Charopidae sp. 145 (NMNZ M.92795)		
Snail	Charopidae sp. 146 (NMNZ M.29932)		
Snail	Charopidae sp. 148 (NMNZ M.99636)		
Snail	Charopidae sp. 151 (NMNZ M.127919)		
Snail	Charopidae sp. 155 (NMNZ M.79509)		
Snail	Charopidae sp. 172 (NMNZ M.76144)		
Snail	Charopidae sp. 175 (NMNZ M.86590)		
Snail	Charopidae sp. 176 (NMNZ M.88962)		
Snail	Charopidae sp. 178 (NMNZ M.89920)		
Snail	Charopidae sp. 180 (NMNZ M.100098)		
Snail	Charopidae sp. 181 (NMNZ M.103775)		
Snail	Charopidae sp. 183 (NMNZ M.120284)		
Snail	Charopidae sp. 186 (NMNZ M.28762)		
Snail	Charopidae sp. 188 (NMNZ M.30514)		
Snail	Charopidae sp. 194 (NMNZ M.115991)		
Snail	Charopidae sp. 200 (NMNZ M.86260)		
Snail	Charopidae sp. 203 (NMNZ M.97140)		
Snail	Charopidae sp. 217 (NMNZ M.29839)		
Snipe-fly	<i>Chrysopilus</i> sp. "Paparoa Range"		
Moth	<i>Clepsicosma</i> sp. "Titirangi"		
Moth	<i>Coridomorpha</i> "long palpi"		OL
Cave centipede	<i>Cryptops</i> n. sp.		
Rove beetle	<i>Cyparium</i> sp. "Fiordland"		
Rove beetle	<i>Cyparium</i> sp. "Mangamuka"		
Moth	<i>Ericodesma</i> sp. "mingimingi"		OL
Moth	<i>Eudonta linealis</i> (Walker, 1866)		OL
Snail	<i>Flammulina</i> "constanceae/feredayi" (NMNZ M.87559)		OL
Moth	<i>Gadira</i> "black brown EGW"		
Cave weta	Genus aff. <i>Talitropsis</i> sp. "Poor Knights"		
Cranefly	<i>Gynoplistia</i> "Arthurs"		
Cave centipede	<i>Haasiella</i> sp.		
Ground weta	<i>Hemiandrus</i> "Dodsons"		OL
Ground weta	<i>Hemiandrus</i> "Hapuku"		OL
Ground weta	<i>Hemiandrus</i> "Longwood Range"		OL
Ground weta	<i>Hemiandrus</i> "Mt George"		OL
Ground weta	<i>Hemiandrus</i> "Nokomai"		OL
Ground weta	<i>Hemiandrus</i> "Pureora 1"		OL
Ground weta	<i>Hemiandrus</i> "Pureora 2"		OL
Ground weta	<i>Hemiandrus</i> "Redhills"		OL
Ground weta	<i>Hemiandrus</i> "Richmond"		
Ground weta	<i>Hemiandrus</i> "Rocklands"		OL
Ground weta	<i>Hemiandrus</i> "Staveley"		OL
Ground weta	<i>Hemiandrus</i> "Tapuaenuku"		OL
Ground weta	<i>Hemiandrus</i> "Timaru"		OL

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

*Continued on next page*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Ground weta	<i>Hemiandrus</i> "Waimakariri"	OL	
Moth	<i>Hierodoris</i> "clear wing"		
Moth	<i>Hierodorts</i> "silver banded"	DP, ST	
Moth	<i>Hierodoris</i> "tiger stripes RJBH"	OL	
Snail	"Hyalolaoma" "Kaukapakapa" (NMNZ M.82946)	OL	
Snail	"Hyalolaoma" "Maungatapere" (NMNZ M.78786)	OL	
Moth	<i>Hydriomena iolanthe</i> Hudson, 1939		
Native bee, manuscript name "murihiku"	<i>Hylaeus</i> "murihiku"		
Native bee	<i>Hylaeus</i> sp. "Fiordland yellow-faced bee"		
Snail	<i>Iotula</i> "kessneri" (NMNZ M.69540)		
Moth	<i>Izatha griseata</i> Hudson, 1939		
Moth	<i>Izatha</i> sp. "big brown"		
Moth	<i>Izatha</i> sp. "mini-metadelta"		
Moth	<i>Izatha</i> sp. "small grey"		
Moth	<i>Izatha</i> sp. "tufted disc"		
Moth	<i>Izatha</i> sp. "whorled antennae"		
Small fungus beetle	Leiodidae n. gen. "Karamea"		
Small fungus beetle	Leiodidae n. gen. "Mangamuka Range"		
Native bee	<i>Leioproctus</i> sp. 3 "coastal"		
Snail	<i>Liarea</i> sp. "Bream Head" (NMNZ M.158257)	OL	
Earthworm	<i>Maoridrilus ?intermedius</i>	OL	
Earthworm	<i>Maoridrilus ?mauiensis</i>	OL	
Earthworm	<i>Maoridrilus ?thomsoni</i>	OL	
Beetle	<i>Maorinus</i> sp.		
Ground beetle	<i>Megadromus</i> sp. 9 "Millers Flat"	OL	
Click beetle	<i>Metablax</i> sp.	OL	
Chironomid fly	<i>Microtendipes</i> (Winterbourn et. al., 2000) "Blue Lake"		
Chafer beetle	<i>Odontria</i> sp. "Kirkliston Range"	OL	
Moth	<i>Palaeomicra</i> sp. "Percy"		
Cranefly	Pedeciini n. gen., n. sp. "Banks Peninsula"		
Flax snail (pupuharakeke)	<i>Placostylus (Maoristylus) ambagiosus</i> "Reinga"	HI, OL	VU B1+2abcde (for <i>P. ambagiosus</i> s.l.)
Snail	"Poripotaka" "aff. ramsayi" (NMNZ M.105949)	OL	
Snail	"Powellaoma" "stewartensis" (NMNZ M.79415)	OL	
Large land snail	<i>Powelliphanta</i> "Waitotara"		
Spider—Malkaridae	Present in NZ, but are an unrevised group		
Spider—Mysmenidae	Present in NZ, but family is unworked		
Scarab/chafer beetle	<i>Prodontria</i> sp. "Ben Ohau"	OL	
Scarab/chafer beetle	<i>Prodontria</i> sp. "Five Rivers"	OL	
Scarab/chafer beetle	<i>Prodontria</i> sp. "Nevis"		
Snail	"Pseudophrixgnathus" "aff. elevata" (NMNZ M.151439)	OL	
Snail	Punctidae sp. 4 (NMNZ M.88026)	OL	
Snail	Punctidae sp. 13 (NMNZ M.151444)	OL	
Snail	Punctidae sp. 14 (NMNZ M.96422)		
Snail	Punctidae sp. 19 (NMNZ M.151440)	OL	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

*Data Deficient—continued*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
Snail	Punctidae sp. 26 (NMNZ M.78582)	OL	
Snail	Punctidae sp. 30 (NMNZ M.151445)	OL	
Snail	Punctidae sp. 31 (NMNZ M.151447)	OL	
Snail	Punctidae sp. 32 (NMNZ M.38683)	OL	
Snail	Punctidae sp. 33 (NMNZ M.58191)	OL	
Snail	Punctidae sp. 35 (NMNZ M.151448)	OL	
Snail	Punctidae sp. 36 (NMNZ M.151446)	OL	
Snail	Punctidae sp. 37 (NMNZ M.123669)		
Snail	Punctidae sp. 44 (NMNZ M.88276)		
Snail	Punctidae sp. 66 (NMNZ M.125724)		
Snail	Punctidae sp. 111 (NMNZ M.115948)		
Snail	Punctidae sp. 170 (NMNZ M.99071)		
Snail	Punctidae sp. 209 (NMNZ M.151452)		
Snail	Punctidae sp. 218 (NMNZ M.38666)		
Snail	Punctidae sp. 233 (NMNZ M.77777)		
Snail	Punctidae sp. 248 (NMNZ M.107891)		
Snail	Punctidae sp. 249 (NMNZ M.126644)		
Snail	Punctidae sp. 268 (NMNZ M.127753)		
Snail	Punctidae sp. 269 (NMNZ M.116424)		
Snail	Punctidae sp. 270 (NMNZ M.127742)		
Moth	<i>Pyroderces</i> sp. “yellow”	OL	
Snail	<i>Ranfurlya</i> “meadsi” (NMNZ M.151455)	OL	
Cave weta	Rhaphidophoridae sp. “Poor Knights”		
Earthworm	<i>Rhododrilus</i> ? <i>monticola</i>	OL	
Snail	“Roscoepunctum” “elworthyi” (NMNZ M.83481)	OL	
Snail	“Roscoepunctum” “Mt Duppera” (NMNZ M.106034)	OL	
Snail	“Roscoepunctum” “sutherlandi” (NMNZ M.83482)	OL	
Snail	“Roscoepunctum” “Takaka Hill” (NMNZ M.126713)	OL	
Snail	<i>Rotadiscus</i> “aff. jamiesoni” (NMNZ M.89806)	OL	
Robber fly	<i>Saropogon</i> “rereke”		
Snail	<i>Sinployea</i> “Ohinepoutea” (NMNZ M.78239)	OL	
Snail	<i>Sinployea</i> “parvaformis” (NMNZ M.76163)	OL	
Moth	<i>Stigmella</i> sp. “blue”	OL	
Moth	<i>Stigmella</i> sp. “giant”	OL	
Moth	<i>Stigmella</i> sp. “October”	OL	
Weevil	<i>Sympedius</i> sp. “Chathams”		
Moth	<i>Theoxena</i> sp. “non-pectinate”		
Snail	<i>Therasta</i> sp. (NMNZ M.73237)	OL	
Snail	<i>Therastiella</i> “narrow umbilicus” (NMNZ M.79608)	OL	
Moth	Tineidae n. sp. “Hicks Bay stripe”		
Moth	Tineidae n. sp. “Matuku monster”		
Moth	<i>Trachypepla</i> sp. “Cloudy Bay”		
Kamo carabid	Undescribed genus	OL	
Snail	“Vicariodiscus” “cresswelli” (NMNZ M.113794)		
Snail	<i>Wainutia</i> “Umbrella Range” (NMNZ M.127657)	OL	
Snail	<i>Zelandiscus</i> “innubilis” (NMNZ M.89834)	OL	
Ground beetle	<i>Zeopoecilus</i> “D’Urville Island-Squally Cove”		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

*Continued on next page*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
<b>VASCULAR PLANTS</b>			
<b>Taxonomically determinate</b>			
	<i>Cardamine bilobata</i> Kirk		
	<i>Carex allanti</i> Hamlin	DP	
	<i>Carex carsei</i> Petrie		
	<i>Carex druceana</i> Hamlin		
	<i>Carex kirkii</i> var. <i>elatior</i> Kük. (CHR 73151)	OL	
	<i>Carex raoulii</i> Boott		
	<i>Celmisia cordatifolia</i> var. <i>brockettii</i> Martin (WELT 4496)		
	<i>Celmisia cordatifolia</i> var. <i>similis</i> Martin (CHR 365726)		
	<i>Centipeda aotearoana</i> N.G. Walsh		
	<i>Chenopodium detestans</i> Kirk	TO	
	<i>Chenopodium pusillum</i> Hook.f.		
	<i>Colobanthus brevisepalus</i> Kirk		
	<i>Corallospantium crassicaule</i> var. <i>racemosum</i> Kirk (CHR 141532)		
	<i>Crassula mataikona</i> A.P. Druce		
	<i>Deschampsia pusilla</i> Petrie		
	<i>Elymus tenuis</i> (Buchanan) Å. Löve et Connor		
	<i>Epilobium hirtigerum</i> A. Cunn.	DP, SO, HI	
	<i>Epilobium pictum</i> Petrie		
	<i>Epilobium vernicosum</i> Cheeseman		
	<i>Euchiton polylepis</i> (D.G. Drury) Breitw. et J.M. Ward		
	<i>Euphrasia integrifolia</i> Petrie		
	<i>Geranium microphyllum</i> Hook.f. s.s.		
	<i>Geum albiflorum</i> (Hook.f.) Cheeseman		
	<i>Grammitis gunnii</i> Parris	SO, DP	
	<i>Gunnera densiflora</i> Hook.f.		
	<i>Hebe matthewsii</i> (Cheeseman) Cockayne		
	<i>Koeleria rigidior</i> Edgar et Gibb		
	<i>Lachnagrostis uda</i> Edgar		
	<i>Lagenifera montana</i> Hook.f.	DP	
	<i>Lepidium desvauxii</i> Thell.	SO	
	<i>Lepilaena bilocularis</i> Kirk	SO	
	<i>Libertia cranwelliae</i> Blanchon, B.G. Murray et Braggins	DP	
	<i>Myosotis glabrescens</i> L.B. Moore		
	<i>Myosotis laingii</i> Cheeseman		
	<i>Myosotis petiolata</i> var. <i>pottsiiana</i> L.B. Moore		
	<i>Myosotis suavis</i> Petrie		
	<i>Myosotis uniflora</i> Hook.f.		
	<i>Nematoceras rivularis</i> Hook.f. s.s.		
	<i>Neopaxia linearifolia</i> Heenan		
	<i>Olearia angulata</i> Kirk		DD
	<i>Pachystegia minor</i> (Cheeseman) Molloy		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continued on next page

*Data Deficient—continued*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
	<i>Polygonum plebeium</i> R. Br.	SO	
	<i>Pterostylis auriculata</i> Colenso		
	<i>Pterostylis irwintii</i> D.L. Jones, Molloy et M.A. Clem.		
	<i>Pterostylis porrecta</i> D.L. Jones, Molloy et M.A. Clem.		
	<i>Schizeilema allani</i> Cheeseman		
	<i>Stenostachys laevis</i> (Petrie) Connor		
	<i>Vittadinia australis</i> Hook.f.		
	<i>Zotovia acicularis</i> Edgar et Connor		
<b>Taxonomically indeterminate</b>			
	<i>Agrostis</i> (a) (CHR 402485; Dunstan Range)	OL	
	<i>Brachyglossis cockaynei</i> (G. Simpson et J.S. Thomson) B. Nord. (AK 253995)		
	<i>Brachyglossis rotundifolia</i> var. <i>ambigua</i> (Cheeseman) B. Nord. (AK 251870)	OL	
	<i>Cardamine</i> aff. <i>bilobata</i> (CHR 511915; eastern South Island)		
	<i>Celmisia</i> aff. <i>discolor</i> (CHR 197967; Fiordland)		
	<i>Celmisia</i> aff. <i>gracilenta</i> (a) (CHR 282958; Te Mata Peak)	OL	
	<i>Dracophyllum</i> (Venter 13745; Mt Rochfort)	DP	
	<i>Elymus</i> aff. <i>solandri</i> (a) (AK 222754; “channel”)		
	<i>Elymus</i> aff. <i>solandri</i> (b) (CHR 1613; South Marlborough)		
	<i>Galtum</i> aff. <i>perpusillum</i> (CHR 476063; Kaitorete)		
	<i>Geranium</i> (c) (CHR 546319; Von)	OL	
	<i>Grammitis</i> aff. <i>rawlingsii</i> (a) (CHR 420132; Great Barrier)		
	<i>Grammitis</i> aff. <i>rawlingsii</i> (b) (AK 236942; Mt Williams)	OL	
	<i>Hebe</i> aff. <i>brevifolia</i> (AK 235669; Surville Cliffs)	OL, CD	
	<i>Hebe</i> aff. <i>treadwellii</i> (CHR 394533; Bald Knob Ridge)	OL	
	<i>Leptinella intermedia</i> (D.G. Lloyd) D.G. Lloyd et C. Webb (CANU 17225)		
	<i>Libertia</i> aff. <i>ixioides</i> (a) (CHR 469712; “large capsule”)		
	<i>Libertia</i> aff. <i>ixioides</i> (b) (CHR 174779; Omaha)		
	<i>Limosella</i> (a) (CHR 222625; Opunake)		
	<i>Melicytus</i> aff. <i>alpinus</i> (a) (CHR 541565; Rangipo)		
	<i>Melicytus</i> aff. <i>alpinus</i> (b) (CHR 541568; Otago)		
	<i>Melicytus</i> aff. <i>alpinus</i> (c) (CHR 541567; “dark”)		
	<i>Melicytus</i> aff. <i>alpinus</i> (d) (CHR 530143; “Brockie”)	OL	
	<i>Melicytus</i> aff. <i>alpinus</i> (e) (CHR 514919B; Livingstone)		
	<i>Melicytus</i> aff. <i>obovatus</i> (c) (CHR 393733; Mt Owen)	OL	
	<i>Myosotis</i> (c) (CHR 198630; Fiordland)		
	<i>Myosotis</i> aff. <i>pulvinaris</i> (CHR 431563; Umbrella)		
	<i>Myosotis</i> aff. <i>tenericaulis</i> (AK 7570; Garvie)		
	<i>Nematoceras</i> aff. <i>rivularis</i> (AK 251833; Kaitarakih)	OL	

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

*Continued on next page*

*Data Deficient—continued*

COMMON NAME	TAXON (2004)	QUALIFIER <sup>a</sup>	IUCN <sup>b</sup>
	<i>Nematoceras</i> aff. <i>rivularis</i> (CHR 518025; Kaimai)		
	<i>Nematoceras</i> aff. <i>rivularis</i> (CHR 518313; “whiskers”)		
	<i>Nematoceras</i> aff. <i>rivularis</i> (CHR 534752; “rest area”)		
	<i>Nematoceras</i> aff. <i>trilobus</i> (CHR 518304; “pygmy”)		
	<i>Nematoceras</i> aff. <i>trilobus</i> (CHR 534742; Trotters)		
	<i>Nematoceras</i> aff. <i>trilobus</i> (CHR 537604; Rimutaka)		
	<i>Pachystegia</i> aff. <i>insignis</i> (CHR 565298; Lowry)		
	<i>Pimelea</i> (f) (AK 189577; Maunganui Bluff)	OL	
	<i>Pimelea</i> aff. <i>prostrata</i> (CHR 257898; Kaitorete)	OL	
	<i>Poa</i> aff. <i>sublimis</i> (CHR 402510; Eyre Mountains)	OL	
	<i>Pratia</i> aff. <i>macrodon</i> (AK 255606; Old Man Range)	OL	
	<i>Pterostylis</i> aff. <i>graminea</i> (CHR 513330; “sphagnum”)		
	<i>Ranunculus</i> (c) (CHR 472008; Garvie Range)		
	<i>Ranunculus</i> aff. <i>reflexus</i> (CHR 394270; Mt Peel)		
	<i>Raoulia</i> (a) (CHR 79537; “K”)		
	<i>Rytidosperma</i> <i>tenue</i> (Petrie) Connor et Edgar (WELT 39920)		
	<i>Spiranthes</i> aff. <i>novae-zelandiae</i> (CHR 518297; Motutangi)	HI, EF	
	<i>Thelymitra</i> aff. <i>longifolia</i> (CHR 537579; Whakapapa)		

<sup>a</sup> Qualifiers: CD—Conservation Dependent; DP—Data Poor; EF—Extreme Fluctuations; EW—Extinct in the wild; HI—Human Induced; OL—One Location; RC—Recovering; RF—Recruitment Failure; SO—Secure Overseas; ST—Stable; TO—Threatened Overseas.

<sup>b</sup> See [www.redlist.org/info/categories\\_criteria.html](http://www.redlist.org/info/categories_criteria.html) for an explanation of these IUCN categories.

Continue to next file: Part 2