Further information:

Motukarara Conservation Nursery

Waihora Domain, Motukarara, RD 2, Christchurch 7672

PHONE: 03 329 7846

EMAIL: motukarara@doc.govt.nz

www.doc.govt.nz/ conservationnursery

www.doc.govt.nz

Native plants natural to Canterbury Plains





Cover photo: Kōwhai

Published by

Department of Conservation Mahaanui/Sockburn Office PO Box 11089, Sockburn 844 Christchurch, New Zealand

Editing and design: DOC Creative Service Conservation House Wellington

May 202

This publication is produced using paper sourced from well-managed renewable and legally logged fores

New Zealand Government



The history of the native vegetation of the Canterbury Plains is one of sucession, destruction and regeneration. For many millions of years prior to European settlement the patterns of native vegetation were defined by shifting rivers and creeks, floods, shingle build-up, alluvial deposits, soil damage and fire.

The plant life varied between:

- 1. Prostrate herbs and grasses on gravels.
- 2. Tussocks and small shrubs on younger terraces.
- Woodland vegetation of kānuka, cabbage tree, olearia and kōwhai on higher, older terraces.
- 4. Dense woodlands of broad-leaf hardwoods such as karamū, lancewood, *Hoheria angustifolia, Lophomyrtus obcordata* etc. on deeper soils.
- 5. Podocarps on fingers of deep, moist soils.

Today's landscape has suffered 150 years of intensive manipulation, not only for economic reasons but also to transform it into one that is English in character. This has been achieved by the introduction of exotic trees, shrubs and grasses, and grazing, drainage and cultivation of the land.

Planting species represented in these display gardens will eventually provide a link between the native remnants of the Canterbury Plains, (kānuka forests of Eyrewell and Bankside reserves), the foothills (beech forest of the Oxford-Ashley area) and the tall podocarp/hardwood forests of Riccarton Bush in Christchurch and Port Hills reserves.

R = rare

T = threatened

Aciphylla subflabellata, kuri kuri, spear grass

Anemanthele lessoniana (R), hunangāmoho, wind grass,

bamboo grass

Austroderia richardii, toetoe

Brachyscome pinnata (T)

Carex buchananii, matirewa, cutty grass

Carex comans, tussock grass

Carex flagellifera, mānia, tussock grass

Carmichaelia grandiflora

Carmichaelia kirkii (T)

Clematis marata

Coprosma areolata

Coprosma crassifolia

Coprosma intertexta (R)

Coprosma lucida, shining karamū

Coprosma obconica (R/T)

Coprosma sp. 'Taylorii'

Coprosma propinqua, mingimingi

Coprosma robusta, karamū

Coprosma rotundifolia, round-leaved coprosma

Coprosma rugosa

Coprosma pedicellata (R/T)

Coprosma virescens

Cordyline australis, tī kōuka, cabbage tree

Corokia cotoneaster, korokio, corokia

Elaeocarpus hookerianus, põkākā

Festuca novae-zelandiae, fescue grass

Griselinia littoralis, pāpāuma, broadleaf

Helichrysum lanceolatum, niniao

Hypericum gramineum, rolled-leaf hypericum

Kunzea serotina, kānuka

Leptinella filiformis (T)

Leptinella squalida

Leptospermum scoparium, mānuka, tea tree

Lophomyrtus obcordata, rõhutu, NZ myrtle

Melicytus alpinus, porcupine shrub

Muehlenbeckia astonii (T), shrubby tororaro

Myrsine australis, māpou, matipou

Myrsine divaricata, weeping māpou

Olearia adenocarpa (T), scented tree-daisy

Ozothamnus leptophyllus, tauhinu, golden cottonwood

Phormium tenax, harakeke, New Zealand flax

Pittosporum eugenioides, tarata, lemonwood

Pittosporum tenuifolium, kōhūhu

Plagianthus regius, mānatu, ribbonwood

Poa cita, wī, silver tussock

Podocarpus totara, tōtara

Pomaderris phylicifolia var. ericifolia (T), tauhinu


Prumnopitys taxifolia, mataī, black pine

Raukaua anomalus, whauwhaupaku

Sophora microphylla, kōwhai

Sophora prostrata, dwarf kōwhai

Teucridium parvifolium (T)



Higher, older terraces

Open woodland vegetation Dense broadleaf hardwood

Young terraces

Tussocks and small shrubs

Gravels

Prostrate herbs and grasses