Figure 3: Hut Valley 1980–2000



Hut Valley 1980. Low grassy turf and scattered club sedge. Turf maintained by rabbits.



Hut Valley 1985. Dense sward of low kanuka scrub.



Hut Valley 1988. Dense kanuka scrub.



Hut Valley 2000. Dense tall kanuka scrub.

Figure 4: Photopoint 1—Duneland at Te Ratahi (McEwan's Bay), 1979–2002




Figure 5: Photopoint 2—Eastern Side of Motu Hara (Main Summit), 1966–2002

Major changes from grassland to pohutukawa treeland and then to pohutukawa forest.







1990



1996



2001



Figure 6: Photopoint 3—Eastern end of island, adjacent to Te Ratahi (McEwan's Bay), 1955–2002

Major change from grassland to dense pohutukawa forest.




Figure 7: Photopoint 8—Hut Valley, 1955–2002 Change from grassland to kanuka forest.









1990 1996





Figure 8: Photopoint 9E—Lower southern slopes of Raetihi (Pa Hill), 1952–2002






Figure 9: Photopoint 10A—Southern slopes of Motu Hara (Main Summit), 1964–2002









1996 2001



Figure 10: Photopoint 11—Saddle at the head of Onepu (Sulphur Bay) Valley, 1955–2002
Change from grassland to dense windshorn kanuka scrub.




Figure 11: Photopoint 13—Head of Hut Valley, 1990–2002 Change from grassland to pohutukawa treeland and forest.









Figure 12: Photopoint HH1—lower southern slopes of Raetihi (Pa Hill), 1990–2002

Vegetation has changed from grassland to pohutukawa-kanuka/grassland to pohutukawa forest and treeland.





1990 1996





Figure 13: Photopoint 14—Southern and western slopes of Motu Hara (Main Summit), 1955–2001






Figure 14: Photopoint 15—Head of Hut Valley, eastern side, 1990–2002









Figure 15: Photopoint 16—Dunes and flat land adjacent to Oneroa (Boulder Bay), 1955–2002









1996 2001

