

# Key facts about Moehau rat and possum control 2017

The Department of Conservation is undertaking pest control over approximately 4,500 hectares in the Moehau ecological area.

# Why are we undertaking pest control in Moehau?

The Moehau ecological area contains large areas of mixed podocarp-broadleaf forest, with remnant kauri, coastal pohutakawa forest and manuka and kanuka shrublands. These places are valued for their diverse native flora and fauna, and scenic landscapes and a wide range of recreation opportunities.

High rat numbers have negative impacts through predation of native lizards, eggs, birds, insects and seeds. They compete with native animals for food and prevent seedling growth. Therefore the primary outcome of the operation is low rat numbers.

Possums also consume large amounts of native vegetation which significantly impacts the health of the forest. They also raid nests for eggs and fledglings and compete with native animals for food. Controlling possums allows forest habitats to recover; therefore the secondary outcome of the operation is low possum numbers.

### What has been decided?

Cereal pellets containing bio-degradable sodium fluoroacetate (known as 1080) are being used to control pests. Helicopters are

being used to distribute the pellets through the operational area.

This operation is being conducted as part of a wider Coromandel pest control operation. Two other operations in the Papakai and Otahu ecological areas will be conducted within the same timeframe. Ground control techniques are also under discussion for surrounding land areas.

# Timing of the operation

The planned operation timeframe is between July and October 2017. The exact timing of the operation will be weather dependent and will consist of a non-toxic pre-feed, followed by a toxin drop.

Pest operations are timed for late winter and early spring when food sources for target pests are at their lowest. This increases the take up of bait and the success of the operation.

In addition hitting predators at this time reduces predation on native species through at least one and possibly two breeding seasons to improve nesting survival and increase populations.

Why are we using these methods?

Produced by: Department of Conservation Whitianga Field Base PO Box 276 Whitianga 3510 February 2017



Department of Conservation *Te Papa Atawbai* 

Aerial 1080 is the most efficient and cost effective method currently available to reduce rat, possum and stoat (through secondary poisoning) populations to low levels. This is especially true in large and difficult to access areas.

Baits are applied by helicopter using GPS satellite navigation technology. This results in an even coverage and allows the pilot to be sure of the exact location of the operation boundaries.

Spreading the bait evenly increases the exposure to rats and possums and it is expected the majority will encounter and consume the bait.

### Operational planning

The use of toxins for this work requires the consent of the Medical Officer of Health and resource consent.

DOC managers follow procedures and apply conditions to ensure that all legal and policy requirements are met, and that any potential risks are mitigated or managed.

## Key facts

The pesticides we use are poisonous to most animals, including to humans and domestic animals. Poisoning can occur through eating baits or poisoned animals. The risk to dogs with pesticide in carcasses will remain until they have decomposed, which can take from four to nine months.

These risks can be eliminated by following these simple rules:

- DO NOT touch bait
- WATCH CHILDREN at all times if you are visiting this area during the operational period
- DO NOT EAT animals from this area
- Remember, poison baits or carcasses are DEADLY to DOGS

Observe these rules whenever you see warning signs about pesticides. Warning signs indicate that pesticide residues may still be present in baits or animals. When signs are removed, this means that you can resume activities as normal in the area. If in doubt, check with DOC.

### For more information


Visit the following website: www.doc.govt.nz/1080

If you would like more information contact:


Liz Whitwell
Department of Conservation
Whitianga Field Base
Whitianga

T: 07 869 5637

E: coromandelenquiries@doc.govt.nz








1.5 Kilometre

Scale at A4 = 1:50,000
NZGD 2000 New Zealand Transverse Mercator
Not for navigation
Crown Copyright Reserved
Basemap: LINZ Topo250 / 50 BA36 Whangamata
Date Produced: 23/03/2017
DOC, Geospatial Services

Moehau

Proposed 1080 Aerial Pest Control Map 2017





10 Kilometre

Scale at A4 = 1:400,000
NZGD 2000 New Zealand Transverse Mercator
Not for navigation
Crown Copyright Reserved
Basemap: LINZ Topo250 / 50 BA36 Whangamata
Date Produced: 21/03/2017
DOC, Geospatial Services

Moehau / Papakai / Otahu / Mataora

Proposed 1080 Aerial Pest Control Map 2017

